

# Conjugation in Semigroups

João Araújo\*

Universidade Aberta, R. Escola Politécnica, 147  
1269-001 Lisboa, Portugal

&

Centro de Álgebra da Universidade de Lisboa  
1649-003 Lisboa, Portugal, mjoao@lmc.fc.ul.pt

Janusz Konieczny†

Department of Mathematics, University of Mary Washington  
Fredericksburg, VA 22401, jkoniecz@umw.edu

António Malheiro

Departamento de Matemática, Faculdade de Ciências e Tecnologia,  
Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

&

Centro de Álgebra da Universidade de Lisboa  
1649-003 Lisboa, Portugal, malheiro@cii.fc.ul.pt

## Abstract

The action of any group on itself by conjugation and the corresponding conjugacy relation play an important role in group theory. There have been several attempts to extend the notion of conjugacy to semigroups. In this paper, we present a new definition of conjugacy that can be applied to an arbitrary semigroup and it does not reduce to the universal relation in semigroups with a zero. We compare the new notion of conjugacy with existing definitions, characterize the conjugacy in various semigroups of transformations on a set, and count the number of conjugacy classes in these semigroups when the set is infinite.

2010 *Mathematics Subject Classification.* 20M07, 20M20, 20M15.

## 1 Introduction

Let  $G$  be a group. For elements  $a, b \in G$ , we say that  $a$  is *conjugate* to  $b$  if there exists  $g \in G$  such that  $b = g^{-1}ag$ . It is clear that this relation is an equivalence on  $G$  and that  $a$  is conjugate to  $b$  if and only if there exists  $g \in G$  such that  $ag = gb$ . Using the latter formulation, one may try to extend the notion of conjugacy to semigroups in the following way: define a relation  $\sim_l$  on a semigroup  $S$  by

$$a \sim_l b \Leftrightarrow \exists_{g \in S^1} ag = gb, \quad (1.1)$$

where  $S^1$  is  $S$  with an identity adjoined. If  $a \sim_l b$ , we say that  $a$  is *left conjugate* to  $b$  [34, 39, 40]. (We will write “ $\sim$ ” with various subscripts for possible definitions of conjugacy in semigroups.

---

\*Partially supported by FCT through the following projects: PEst-OE/MAT/UI1043/2011, Strategic Project of Centro de Álgebra da Universidade de Lisboa; and PTDC/MAT/101993/2008, Project Computations in groups and semigroups.

†Supported by the 2011–12 University of Mary Washington Faculty Research Grant.

The subscript in  $\sim_l$  comes from the name “left conjugate.”) In a general semigroup  $S$ , the relation  $\sim_l$  is reflexive and transitive, but not symmetric. If  $S$  has a zero, then  $\sim_l$  is the universal relation  $S \times S$ . The relation  $\sim_l$  is an equivalence in any free semigroup. Lallement [28] has defined the conjugate elements of a free semigroup  $S$  as those related by  $\sim_l$  and showed that  $\sim_l$  is equal to the following equivalence on the free semigroup  $S$ :

$$a \sim_p b \Leftrightarrow \exists_{u,v \in S^1} a = uv \text{ and } b = vu. \quad (1.2)$$

In a general semigroup  $S$ , the relation  $\sim_p$  is reflexive and symmetric, but not transitive. If  $a \sim_p b$  in a general semigroup, we say that  $a$  and  $b$  are *primarily conjugate* [27] (hence the subscript in  $\sim_p$ ). Kudryavtseva and Mazorchuk [26, 27] have defined the transitive closure  $\sim_p^*$  of  $\sim_p$  as a conjugacy relation in a general semigroup. (See also [18].)

Otto [34] has studied the relations  $\sim_l$  and  $\sim_p$  in the monoids  $S$  presented by finite Thue systems, and introduced a new definition of conjugate elements in such an  $S$ :

$$a \sim_o b \Leftrightarrow \exists_{g,h \in S^1} ag = gb \text{ and } bh = ha. \quad (1.3)$$

(Since  $S$  is a monoid,  $S^1 = S$ . However, we wanted to write the definition of  $\sim_o$  so that it would apply to any semigroup.) For any semigroup  $S$ ,  $\sim_o$  is an equivalence on  $S$ , and so it provides another possible definition of conjugacy in a general semigroup. However, this definition is not useful for semigroups  $S$  with zero since for every such  $S$ , we have  $\sim_o = S \times S$ . Note that  $\sim_o$  the largest equivalence contained in  $\sim_l$  and that  $\sim_p \subseteq \sim_o$  since if  $a = uv$  and  $b = vu$ , then  $au = ub$  and  $bv = va$ .

The aim of this paper is to introduce a new definition of conjugacy in an arbitrary semigroup, avoiding the problems of the notions listed above. (That is,  $\sim_l$  is not symmetric; both  $\sim_l$  and  $\sim_o$  reduce to the universal relation in semigroups with zero; and  $\sim_p$  is not transitive and so it requires taking the transitive closure.) Our conjugacy will be an equivalence relation  $\sim_c$  on any semigroup  $S$ , it will not reduce to the universal relation even when  $S$  has a zero, and it will be such that  $\sim_c \subseteq \sim_o \subseteq \sim_l$  in every semigroup  $S$ ,  $\sim_c = \sim_o$  if  $S$  is a semigroup without zero, and  $\sim_c = \sim_l = \sim_p = \sim_o$  if  $S$  is a group or a free semigroup.

In Section 2 we introduce the new notion of conjugacy and prove some basic results. The following four sections are devoted to the study of  $\sim_c$  in several transformation semigroups on a finite or infinite set  $X$ . The tools we use in this study are the characterization of  $\sim_c$  in transformation semigroups in terms of certain partial homomorphisms of directed graphs (Section 3) and the concept of a connected partial transformation (Section 4). Conjugacy classes in the partial transformation monoid  $P(X)$  are characterized (for any  $X$ ) and counted (for an infinite  $X$ ) in Section 5; conjugacy in the full transformation monoid  $T(X)$  is treated in Section 6; and Section 7 deals with the monoid  $\Gamma(X)$  of full injective transformations. The paper ends with a number of problems for experts in combinatorics, symbolic dynamics, set theory, semigroups, and matrix theory (Section 8).

## 2 Definition of Conjugacy

We briefly describe the motivation of our new concept of conjugacy. The starting point was the relation  $\sim_o$  introduced by Otto [34]. As we have already pointed out, the relation  $\sim_o$  is the universal relation  $S \times S$  if a semigroup  $S$  has a zero. Our goal has been to retain Otto’s concept for semigroups without zero but modify his definition in such a way that the resulting conjugacy would not reduce to triviality for semigroups with zero.

To find a suitable definition, we considered the semigroup  $P(X)$  of partial transformations on  $X$ , that is, the set of all mappings whose domain and image are included in  $X$ , with function composition as multiplication. This semigroup has a zero, namely the transformation whose domain is empty. Let  $\alpha, \beta \in P(X)$ . Then  $\alpha \sim_o \beta$  if and only if  $\alpha\phi = \phi\beta$  and  $\beta\psi = \psi\alpha$  for some  $\phi, \psi \in P(X)$ . (We write functions on the right and compose from left to right.) Of course, the

last two equalities hold for  $\phi = \psi = 0$ . We could insist that  $\phi$  and  $\psi$  should not be 0 but this would not solve the problem since then the resulting relation would not be transitive.

The solution is this. In the composition  $\alpha\phi$ , it only matters how  $\phi$  is defined on the elements of  $\text{im}(\alpha)$  (the image of  $\alpha$ ). We insist that  $\phi$  be defined for *all* elements of  $\text{im}(\alpha)$ , that is, that  $\text{im}(\alpha) \subseteq \text{dom}(\phi)$ . With the requirement that the transformations  $\phi$  and  $\psi$  come from the sets  $\{\phi \in P(X) : \text{im}(\alpha) \subseteq \text{dom}(\phi)\}$  and  $\{\psi \in P(X) : \text{im}(\beta) \subseteq \text{dom}(\psi)\}$ , the relation is an equivalence. Moreover, we will verify that for  $\alpha \neq 0$ ,  $\text{im}(\alpha) \subseteq \text{dom}(\phi)$  if and only if  $(\gamma\alpha)\phi \neq 0$  for every  $\gamma\alpha \in P(X)\alpha \setminus \{0\}$ , where  $P(X)\alpha \setminus \{0\}$  is the left principal ideal generated by  $\alpha$  with 0 removed. Therefore, the requirement that  $\phi$  and  $\psi$  have “large” domains can be expressed in an abstract semigroup. These considerations motivate the definition below.

Let  $S$  be a semigroup with zero. For  $a \in S$  with  $a \neq 0$ , consider  $S^1a \setminus \{0\}$ , the left principal ideal generated by  $a$  with zero removed. We will denote by  $\mathbb{P}(a)$  the set of all elements  $g \in S$  such that  $(ma)g \neq 0$  for all  $ma \in S^1a \setminus \{0\}$ . We define  $\mathbb{P}(0)$  to be  $\{0\}$ . If  $S$  has no zero, we agree that  $\mathbb{P}(a) = S$  for every  $a \in S$ . We will write  $\mathbb{P}^1(a)$  for  $\mathbb{P}(a) \cup \{1\}$ , where 1 is the identity in  $S^1$ .

**Definition 2.1.** Define a relation  $\sim_c$  on a semigroup  $S$  by

$$a \sim_c b \Leftrightarrow \exists_{g \in \mathbb{P}^1(a)} \exists_{h \in \mathbb{P}^1(b)} ag = gb \text{ and } bh = ha. \quad (2.1)$$

If  $a \sim_c b$ , we say that  $a$  is *conjugate* to  $b$ .

The relation  $\sim_c$  will be called the *conjugacy* on  $S$ , which is justified by the following theorem.

**Theorem 2.2.** *Let  $S$  be a semigroup. Then:*

- (1) *the relation  $\sim_c$  is an equivalence relation on  $S$ ;*
- (2) *if  $\sim_l$ ,  $\sim_p$ , and  $\sim_o$  are relations on  $S$  defined by (1.1), (1.2), and (1.3), respectively, then:*
  - (a)  $\sim_c \subseteq \sim_o \subseteq \sim_l$ ,
  - (b) *if  $S$  is a semigroup without zero, then  $\sim_c = \sim_o$ , and*
  - (c) *if  $S$  is a group or a free semigroup, then  $\sim_c = \sim_l = \sim_p = \sim_o$ .*

*Proof.* It is clear that  $\sim_c$  is reflexive and symmetric. Suppose  $a \sim_c b$  and  $b \sim_c c$ . Then there are  $g_1 \in \mathbb{P}^1(a)$  and  $g_2 \in \mathbb{P}^1(b)$  such that  $ag_1 = g_1b$  and  $bg_2 = g_2c$ . Thus  $a(g_1g_2) = (ag_1)g_2 = (g_1b)g_2 = g_1(bg_2) = g_1(g_2c) = (g_1g_2)c$ . Let  $ma \in S^1a \setminus \{0\}$ . Since  $g_1 \in \mathbb{P}^1(a)$ , we have  $(mg_1)b = m(ag_1) = (ma)g_1 \neq 0$ . Thus  $(mg_1)b \in S^1b \setminus \{0\}$ , and so, since  $g_2 \in \mathbb{P}^1(b)$ , we have  $(ma)(g_1g_2) = m(ag_1)g_2 = m(g_1b)g_2 = ((mg_1)b)g_2 \neq 0$ . Hence  $g_1g_2 \in \mathbb{P}^1(a)$ . Similarly, there is  $h \in \mathbb{P}^1(c)$  such that  $ch = ha$ . Hence  $a \sim_c c$ , and so  $\sim_c$  is transitive. We have proved (1).

Statements 2(a) and 2(b) follow immediately from the definitions of  $\sim_l$ ,  $\sim_o$ , and  $\sim_c$ . Statement 2(c) is clearly true if  $S$  is a group. Let  $S$  be a free semigroup. Then  $\sim_l = \sim_p$  by [28, Corollary 5.2]. Thus, by 2(a) and 2(b),  $\sim_c = \sim_o \subseteq \sim_l = \sim_p \subseteq \sim_o$ , which implies  $\sim_c = \sim_o = \sim_l = \sim_p$ .  $\square$

For an element  $a$  of a semigroup  $S$ , the equivalence class of  $a$  with respect to  $\sim_c$  will be called the *conjugacy class* of  $a$  and denoted  $[a]_c$ .

Let  $S$  be a semigroup with 0. In contrast with the fact that  $\sim_o = S \times S$ , the conjugacy class of 0 with respect to  $\sim_c$  is  $\{0\}$ , so we always have  $\sim_c \neq S \times S$  unless  $S = \{0\}$ . Indeed, suppose  $a \sim_c 0$ . Then  $ag = g0 = 0$  for some  $g \in \mathbb{P}^1(a)$ . If  $a \neq 0$ , then  $ag \neq 0$  (since  $a \in S^1a \setminus \{0\}$ ). But  $ag = 0$ , and so it follows that  $a = 0$ . Hence we have the following lemma.

**Lemma 2.3.** *If  $S$  is a semigroup with 0 then  $[0]_c = \{0\}$ .*

For a set  $A$ , we denote by  $\Delta_A$  (or  $\Delta$  if  $A$  is understood) the identity relation on  $A$ , that is  $\Delta_A = \{(a, a) : a \in A\}$ . Recall that in any group  $G$ , the relation  $\sim_c$  is the usual group conjugacy, that is  $a \sim_c b$  if and only if  $g^{-1}ag = b$  for some  $g \in G$ . It follows that in any group  $G$ , we have  $\sim_c = \Delta$  if and only if  $G$  is commutative. This result extends to semigroups as follows.

**Theorem 2.4.** *Let  $S$  be a semigroup without zero. Then  $\sim_c = \Delta$  if and only if  $S$  is commutative and cancellative.*

*Proof.* Since  $S$  has no zero,  $\sim_c = \sim_o$ . It is clear that if  $S$  is commutative and cancellative, then  $\sim_c = \Delta$ . Conversely, suppose that  $\sim_c = \Delta$ . Let  $a, b \in S$ . Since  $(ab)a = a(ba)$  and  $(ba)b = b(ab)$ , we have  $(ab) \sim_c (ba)$ , and hence  $ab = ba$ . We have proved that  $S$  is commutative. Let  $a, b, c \in S$  be such that  $ac = bc$ . Since  $S$  is commutative,  $ac = bc$  implies  $a \sim_c b$ , which in turn implies  $a = b$ . It follows that  $S$  is cancellative.  $\square$

Theorem 2.4 is not true for semigroups with zero. For example, let  $S = \{a, 0\}$  be a 2-element semigroup with  $aa = 0$ . Then  $S$  is not cancellative but we already know that  $[0]_c = \{0\}$ , so  $\sim_c = \Delta$ .

### 3 Restrictive Partial Homomorphisms of Digraphs

The remainder of the paper is devoted to the study of the conjugacy  $\sim_c$  in several important semigroups of transformations on a set  $X$  (finite or infinite). The main tool in our study will be the characterization of  $\sim_c$  in terms of certain partial homomorphisms of directed graphs (see Theorem 3.8 and Corollary 3.9).

A *directed graph* (or a *digraph*) is a pair  $\Gamma = (X, R)$  where  $X$  is a non-empty set (not necessarily finite) and  $R$  is a binary relation on  $X$ . Any element  $x \in X$  is called a *vertex* of  $\Gamma$ , and any pair  $(x, y) \in R$  is called an *arc* of  $\Gamma$ . We will call a vertex  $x$  *terminal* if there is no  $y \in X$  such that  $(x, y) \in R$ .

For any mapping  $f : A \rightarrow B$ , we will denote the domain of  $f$  by  $\text{dom}(f)$  and the image of  $f$  by  $\text{im}(f)$ . For  $A_1 \subseteq A$ , we denote by  $f|_{A_1}$  the restriction of  $f$  to  $A_1$ , and by  $A_1 f$  the image of  $A_1$  under  $f$ .

Let  $\Gamma_1 = (X_1, R_1)$  and  $\Gamma_2 = (X_2, R_2)$  be digraphs. A mapping  $\phi : X_1 \rightarrow X_2$  is called a *homomorphism* from  $\Gamma_1$  to  $\Gamma_2$  if for all  $x, y \in X_1$ , if  $(x, y) \in R_1$ , then  $(x\phi, y\phi) \in R_2$  [17]. Generalizing, a partial mapping  $\phi$  from  $X_1$  to  $X_2$  (that is, a mapping  $\phi$  from some subset of  $X_1$  to  $X_2$ ) is called a *partial homomorphism* from  $\Gamma_1$  to  $\Gamma_2$  if for all  $x, y \in X_1$ , if  $(x, y) \in R_1$  and  $x, y \in \text{dom}(\phi)$ , then  $(x\phi, y\phi) \in R_2$ .

**Definition 3.1.** Let  $\Gamma_1 = (X_1, R_1)$  and  $\Gamma_2 = (X_2, R_2)$  be digraphs. A partial mapping  $\phi$  from  $X_1$  to  $X_2$  is called a *restrictive partial homomorphism* (or an *rp-homomorphism*) from  $\Gamma_1$  to  $\Gamma_2$  if it satisfies the following conditions for all  $x, y \in X_1$ :

- (a) if  $(x, y) \in R_1$ , then  $x, y \in \text{dom}(\phi)$  and  $(x\phi, y\phi) \in R_2$ ;
- (b) if  $x$  is a terminal vertex in  $\Gamma_1$  and  $x \in \text{dom}(\phi)$ , then  $x\phi$  is a terminal vertex in  $\Gamma_2$ .

We say that  $\Gamma_1$  is *rp-homomorphic* to  $\Gamma_2$  if there is an rp-homomorphism from  $\Gamma_1$  to  $\Gamma_2$ .

Clearly, every rp-homomorphism from  $\Gamma_1$  to  $\Gamma_2$  is a partial homomorphism from  $\Gamma_1$  to  $\Gamma_2$ . It is also clear that the composition of rp-homomorphisms is an rp-homomorphism.

**Remark 3.2.** Call a vertex  $x$  of a digraph  $\Gamma = (X, R)$  *isolated* if there is no  $y \in X$  such that  $(x, y) \in R$  or  $(y, x) \in R$ . Let  $\phi$  be an rp-homomorphism from  $\Gamma_1 = (X_1, R_1)$  to  $\Gamma_2 = (X_2, R_2)$ . Denote by  $X'_1$  the set of all vertices in  $\Gamma_1$  that are not isolated. Then  $\phi' = \phi|_{X'_1}$  is also an rp-homomorphism from  $\Gamma_1$  to  $\Gamma_2$ .

In picturing directed graphs, we will adopt the convention that the arrows will be deleted with the understanding that the arrow goes up along the edge, to the right if the edge is horizontal, and the arrows go counter-clockwise around a cycle. For example, consider the digraphs  $\Gamma_1 = (X_1, R_1)$ , where  $X_1 = \{1, 2, 3, 4\}$  and  $R_1 = \{(2, 3), (3, 4)\}$ , and  $\Gamma_2 = (X_2, R_2)$ , where  $X_2 = \{a, b, c, d\}$  and  $R_2 = \{(a, b), (b, d), (c, d)\}$ . Then a mapping presented in Figure 3.1 is a partial




Figure 3.1: A partial homomorphism from  $\Gamma_1$  to  $\Gamma_2$ .




Figure 3.2: An rp-homomorphism from  $\Gamma_1$  to  $\Gamma_2$ .

homomorphism from  $\Gamma_1$  to  $\Gamma_2$  (but not a restrictive partial homomorphism), and a mapping from Figure 3.2 is an rp-homomorphism from  $\Gamma_1$  to  $\Gamma_2$ .

Let  $\alpha \in P(X)$ . Then  $\alpha$  can be represented by the digraph  $\Gamma(\alpha) = (X, R_\alpha)$ , where for all  $x, y \in X$ ,  $(x, y) \in R_\alpha$  if and only if  $x \in \text{dom}(\alpha)$  and  $x\alpha = y$  [15, Section 1.2]. If  $x \in \text{dom}(\alpha)$  and  $x\alpha = y$ , we will write  $x \xrightarrow{\alpha} y$  (or  $x \rightarrow y$  if no ambiguity arises). For  $\alpha \in P(X)$ , the set  $\text{dom}(\alpha) \cup \text{im}(\alpha)$  will be called the *span* of  $\alpha$  and denoted  $\text{span}(\alpha)$ .

For example, the digraph in Figure 3.3 represents the transformation

$$\alpha = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \dots \\ 2 & 3 & 1 & 1 & 1 & 5 & 8 & 9 & 10 & \dots \end{pmatrix} \in T(X),$$

where  $X = \{1, 2, 3, \dots\}$  and  $T(X)$  is the semigroup of all  $\alpha \in P(X)$  such that  $\text{dom}(\alpha) = X$ .




Figure 3.3: The digraph of a transformation.

**Definition 3.3.** Any  $\alpha \in P(X)$  with  $\text{im}(\alpha) = \{x\}$  will be called a *constant*. A subsemigroup  $S$  of  $P(X)$  will be called *constant rich* if for every  $x \in X$ , there is  $\alpha \in S$  such that  $\text{im}(\alpha) = \{x\}$ .

Among the constant rich subsemigroups of  $P(X)$ , we have  $P(X)$  itself (an all its nonzero ideals), the full transformation semigroup  $T(X)$  (and all its ideals), and the symmetric inverse semigroup  $\mathcal{I}(X)$  of all injective  $\alpha \in P(X)$  (and all its nonzero ideals).

**Notation 3.4.** From now on, we will fix a nonempty set  $X$  and an element  $\diamond \notin X$ . For  $\alpha \in P(X)$  and  $x \in X$ , we will write  $x\alpha = \diamond$  if and only if  $x \notin \text{dom}(\alpha)$ . We will also assume that  $\diamond\alpha = \diamond$ . With this notation, it will make sense to write  $x\alpha = y\beta$  or  $x\alpha \neq y\beta$  ( $\alpha, \beta \in P(X)$ ,  $x, y \in X$ ) even when  $x \notin \text{dom}(\alpha)$  or  $y \notin \text{dom}(\beta)$ .

We will also denote by  $\mathbb{Z}$ ,  $\mathbb{Z}_+$ , and  $\mathbb{N}$  the set of integers, positive integers, and nonnegative integers, respectively, and for semigroups  $S$  and  $T$ , write  $S \leq T$  to mean that  $S$  is a subsemigroup of  $T$ .

**Lemma 3.5.** Let  $S \leq P(X)$  such that  $S$  is constant rich, let  $\alpha \in S$  with  $\alpha \neq 0$ , and  $\phi \in S^1$ .

- (1)  $\phi \in \mathbb{P}^1(\alpha)$  if and only if  $\text{im}(\alpha) \subseteq \text{dom}(\phi)$ .
- (2) If  $\phi \in \mathbb{P}^1(\alpha)$  and  $\alpha\phi = \phi\beta$  for some  $\beta \in S$ , then  $\text{span}(\alpha) \subseteq \text{dom}(\phi)$  and for all  $x, y \in X$ ,  $x \xrightarrow{\alpha} y$  implies  $x\phi \xrightarrow{\beta} y\phi$ .

*Proof.* Let  $S$  be constant rich. Suppose  $\phi \in \mathbb{P}^1(\alpha)$ . Let  $y \in \text{im}(\alpha)$ , that is,  $y = x\alpha$  for some  $x \in \text{dom}(\alpha)$ . Since  $S$  is constant rich, there is  $\gamma \in S$  with  $\text{im}(\gamma) = \{x\}$ . Then  $\text{im}(\gamma\alpha) = \{y\}$ , and so  $\gamma\alpha \in S^1\alpha \setminus \{0\}$ . Thus  $(\gamma\alpha)\phi \neq 0$  (since  $\phi \in \mathbb{P}^1(\alpha)$ ), which is only possible when  $y \in \text{dom}(\phi)$ . Hence  $\text{im}(\alpha) \subseteq \text{dom}(\phi)$ .

Conversely, suppose  $\text{im}(\alpha) \subseteq \text{dom}(\phi)$ . Let  $\mu\alpha \in S^1\alpha \setminus \{0\}$ . Since  $\mu\alpha \neq 0$ , there is  $x \in X$  such that  $x(\mu\alpha) \neq 0$ . But then  $x(\mu\alpha) = (x\mu)\alpha \in \text{im}(\alpha) \subseteq \text{dom}(\phi)$ , and so  $x \in \text{dom}((\mu\alpha)\phi)$ . Thus  $(\mu\alpha)\phi \neq 0$ , and so  $\phi \in \mathbb{P}^1(\alpha)$ . We have proved (1).

To prove (2), suppose  $\phi \in \mathbb{P}^1(\alpha)$  and  $\alpha\phi = \phi\beta$  for some  $\beta \in S$ . Let  $x, y \in X$  and suppose that  $x \xrightarrow{\alpha} y$ . Then, since  $\alpha\phi = \phi\beta$ , we have

$$(x\phi)\beta = x(\phi\beta) = x(\alpha\phi) = (x\alpha)\phi = y\phi. \quad (3.1)$$

By (1),  $\text{im}(\alpha) \subseteq \text{dom}(\phi)$ , and so  $y = x\alpha \in \text{dom}(\phi)$ . Then, by (3.1),  $x\phi \neq \diamond$ , which implies  $x \in \text{dom}(\phi)$ . It follows that  $\text{span}(\alpha) \subseteq \text{dom}(\phi)$ . Moreover, by (3.1) again,  $(x\phi)\beta = y\phi \neq \diamond$ , and so  $x\phi \xrightarrow{\beta} y\phi$ .  $\square$

**Lemma 3.6.** Let  $\alpha, \beta \in P(X)$  and let  $\phi$  be an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . Then  $\text{span}(\alpha) \subseteq \text{dom}(\phi)$ .

*Proof.* Let  $x \in \text{span}(\alpha)$ . If  $x \in \text{dom}(\alpha)$ , then  $x \xrightarrow{\alpha} x\alpha$ , and so  $x, x\alpha \in \text{dom}(\phi)$  by Definition 3.1. If  $x \in \text{im}(\alpha)$ , then  $x \xrightarrow{\alpha} x$  for some  $z \in \text{dom}(\alpha)$ , and so  $z, x \in \text{dom}(\phi)$ . Hence  $\text{span}(\alpha) \subseteq \text{dom}(\phi)$ .  $\square$

**Lemma 3.7.** Let  $S \leq P(X)$  such that  $S$  is constant rich, let  $\alpha, \beta \in S$  with  $\alpha \neq 0$ , and  $\phi \in S^1$ . Then  $\alpha\phi = \phi\beta$  with  $\phi \in \mathbb{P}^1(\alpha)$  if and only if  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ .

*Proof.* Suppose  $\alpha\phi = \phi\beta$  with  $\phi \in \mathbb{P}^1(\alpha)$ . Let  $x, y \in X$  and suppose that  $x \xrightarrow{\alpha} y$ . Then  $x\phi \xrightarrow{\beta} y\phi$  by Lemma 3.5, and so  $\phi$  satisfies (a) of Definition 3.1. Suppose that  $x$  is a terminal vertex of  $\Gamma(\alpha)$  and  $x \in \text{dom}(\phi)$ . Then  $x\phi \in X$  and  $x\alpha = \diamond$ . Since  $\alpha\phi = \phi\beta$ , we have  $(x\phi)\beta = (x\alpha)\phi = \diamond\phi = \diamond$ , and so  $x\phi$  is a terminal vertex in  $\Gamma(\beta)$ . Hence  $\phi$  satisfies (b) of Definition 3.1. Thus  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ .

Conversely, suppose that  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . Let  $x \in X$ . Suppose  $x \notin \text{dom}(\alpha)$ . Then  $x(\alpha\phi) = (x\alpha)\phi = \diamond\phi = \diamond$ . If  $x \notin \text{dom}(\phi)$ , then  $x(\phi\beta) = (x\phi)\beta = \diamond\beta = \diamond$ .

If  $x \in \text{dom}(\phi)$ , then, by (b) of Definition 3.1,  $x\phi$  is a terminal vertex in  $\Gamma(\beta)$ , and so  $x(\phi\beta) = (x\phi)\beta = \diamond$ . Hence, in both cases,  $x(\alpha\phi) = x(\phi\beta)$ .

Suppose  $x \in \text{dom}(\alpha)$  and let  $y = x\alpha \in X$ . Then  $x \xrightarrow{\alpha} y$ , and so, by Definition 3.1,  $x, y \in \text{dom}(\phi)$  and  $x\phi \xrightarrow{\beta} y\phi$ . Hence  $x(\alpha\phi) = (x\alpha)\phi = y\phi$  and  $x(\phi\beta) = (x\phi)\beta = y\phi$ , and so  $x(\alpha\phi) = x(\phi\beta)$ . We have proved that  $\alpha\phi = \phi\beta$ . Finally, since  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ , we have that  $\text{span}(\alpha) \subseteq \text{dom}(\phi)$  by Lemma 3.6, and so  $\phi \in \mathbb{P}^1(\alpha)$  by Lemma 3.5.  $\square$

**Theorem 3.8.** *Let  $S \leq P(X)$  such that  $S$  is constant rich, let  $\alpha, \beta \in S$ . Then  $\alpha \sim_c \beta$  in  $S$  if and only if there are  $\phi, \psi \in S^1$  such that  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$  and  $\psi$  is an rp-homomorphism from  $\Gamma(\beta)$  to  $\Gamma(\alpha)$ .*

*Proof.* Suppose  $\alpha \sim_c \beta$ . If  $\alpha = 0$ , then  $\beta = 0$  (since  $[0]_c = \{0\}$ ), and so  $\phi = \text{id}_X \in S^1$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . Suppose  $\alpha \neq 0$ . Since  $\alpha \sim_c \beta$ , there is  $\phi \in \mathbb{P}^1(\alpha)$  such that  $\alpha\phi = \phi\beta$ , and so  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$  by Lemma 3.7. A desired  $\psi \in S^1$  exists by symmetry.

Conversely, suppose that desired  $\phi$  and  $\psi$  exist. If  $x \xrightarrow{\alpha} y$  then  $x\phi \xrightarrow{\beta} y\phi$ , and if  $x \xrightarrow{\beta} y$  then  $x\psi \xrightarrow{\alpha} y\psi$ . It follows that either  $\alpha = \beta = 0$  or  $\alpha, \beta \neq 0$ . In the former case, we clearly have  $\alpha \sim_c \beta$ . Suppose  $\alpha, \beta \neq 0$ . Then, by Lemma 3.7,  $\alpha\phi = \phi\beta$  with  $\phi \in \mathbb{P}^1(\alpha)$  and  $\beta\psi = \psi\alpha$  with  $\psi \in \mathbb{P}^1(\beta)$ , which implies  $\alpha \sim_c \beta$ .  $\square$

Let  $\alpha, \beta \in T(X)$ . Then the graph  $\Gamma(\alpha)$  has no terminal vertices (if  $x \in X$ , then  $x \xrightarrow{\alpha} x\alpha$ ), and so every homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$  is restrictive. This observation and Theorem 3.8 give us the following corollary.

**Corollary 3.9.** *Let  $S \leq T(X)$  such that  $S$  contains all constants, and let  $\alpha, \beta \in S$ . Then  $\alpha \sim_c \beta$  in  $S$  if and only if there are  $\phi, \psi \in S^1$  such that  $\phi$  is a homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$  and  $\psi$  is a homomorphism from  $\Gamma(\beta)$  to  $\Gamma(\alpha)$ .*

## 4 Connected Partial Transformations

In this section, we introduce the concept of connected partial transformation. The definitions and results of this section will be crucial in characterizing conjugacy in various semigroups of transformations.

**Definition 4.1.** Let  $\dots, x_{-2}, x_{-1}, x_0, x_1, x_2, \dots$  be pairwise distinct elements of  $X$ . The following elements of  $P(X)$  will be called *basic* partial transformations on  $X$ .

- (1) A *cycle* of length  $k$  ( $k \geq 1$ ), written  $(x_0 x_1 \dots x_{k-1})$ , is an element of  $P(X)$  defined by the digraph  $x_0 \rightarrow x_1 \rightarrow \dots \rightarrow x_{k-1} \rightarrow x_0$ .
- (2) A *right ray*, written  $[x_0 x_1 x_2 \dots]$ , is an element of  $P(X)$  defined by the digraph

$$x_0 \rightarrow x_1 \rightarrow x_2 \rightarrow \dots$$

- (3) A *double ray*, written  $\langle \dots x_{-1} x_0 x_1 \dots \rangle$ , is an element of  $P(X)$  defined by the digraph

$$\dots \rightarrow x_{-1} \rightarrow x_0 \rightarrow x_1 \rightarrow \dots$$

- (4) A *left ray*, written  $\langle \dots x_2 x_1 x_0 \rangle$ , is an element of  $P(X)$  defined by the digraph

$$\dots \rightarrow x_2 \rightarrow x_1 \rightarrow x_0.$$

- (5) A *chain* of length  $k$  ( $k \geq 1$ ), written  $[x_0 x_1 \dots x_k]$ , is an element of  $P(X)$  defined by the digraph  $x_0 \rightarrow x_1 \rightarrow \dots \rightarrow x_k$ .

By a *ray* we will mean a double, right, or left ray.

We note the following.

- (i) All basic partial transformations are injective.
- (ii) The span of a basic partial transformation is exhibited by the notation. For example, the span of the right ray  $[1\ 2\ 3\ \dots]$  is  $\{1, 2, 3, \dots\}$ .
- (iii) The left bracket in “ $\sigma = [x\ \dots]$ ” indicates that  $x \notin \text{im}(\sigma)$ ; while the right bracket in “ $\sigma = \dots x]$ ” indicates that  $x \notin \text{dom}(\sigma)$ . For example, for the chain  $\sigma = [1\ 2\ 3\ 4]$ ,  $\text{dom}(\sigma) = \{1, 2, 3\}$  and  $\text{im}(\sigma) = \{2, 3, 4\}$ .
- (iv) A cycle  $(x_0\ x_1\ \dots\ x_{k-1})$  differs from the corresponding cycle in the symmetric group of permutations on  $X$  in that the former is undefined for every  $x \in X \setminus \{x_0, x_1, \dots, x_{k-1}\}$  while the latter is defined on and fixes every such  $x$ .

**Definition 4.2.** An element  $\gamma \in P(X)$  is called *connected* if  $\gamma \neq 0$  and for all  $x, y \in \text{span}(\gamma)$ ,  $x\gamma^k = y\gamma^m \neq \diamond$  for some integers  $k, m \geq 0$  (where  $\gamma^0 = \text{id}_X$ ).

We note that a nonzero  $\gamma \in P(X)$  is connected if and only if the underlying undirected graph of the digraph  $\Gamma^0(\gamma)$  is connected, where  $\Gamma^0(\gamma)$  is the digraph  $\Gamma(\gamma)$  with the isolated vertices removed, and that all basic partial transformations are connected.

**Definition 4.3.** Let  $\alpha, \beta \in P(X)$ . We say that  $\beta$  is *contained* in  $\alpha$  (or  $\alpha$  *contains* or *has*  $\beta$ ), and write  $\beta \sqsubset \alpha$ , if  $\text{dom}(\beta) \subseteq \text{dom}(\alpha)$  and  $x\beta = x\alpha$  for every  $x \in \text{dom}(\beta)$ . In other words,  $\beta \sqsubset \alpha$  iff  $\beta = \varepsilon\alpha$  where  $\varepsilon$  is the identity on the domain of  $\beta$ . We say that  $\alpha$  and  $\beta$  are *disjoint* if  $\text{dom}(\alpha) \cap \text{dom}(\beta) = \emptyset$ ; they are *completely disjoint* if  $\text{span}(\alpha) \cap \text{span}(\beta) = \emptyset$ .

For example, the right ray  $[3\ 4\ 5\ 6\ \dots]$  and chain  $[0\ 1\ 2\ 5]$  in  $P(\mathbb{Z})$  are disjoint but not completely disjoint. Their join  $[3\ 4\ 5\ 6\ \dots] \sqcup [0\ 1\ 2\ 5]$  (see Definition 4.4 below) is connected.

**Definition 4.4.** Let  $C$  be a set of pairwise disjoint elements of  $P(X)$ . The *join* of the elements of  $C$ , denoted  $\bigsqcup_{\gamma \in C} \gamma$ , is an element of  $P(X)$  defined by

$$x \left( \bigsqcup_{\gamma \in C} \gamma \right) = \begin{cases} x\gamma & \text{if } x \in \text{dom}(\gamma) \text{ for some } \gamma \in C, \\ \diamond & \text{otherwise.} \end{cases}$$

If  $C = \{\gamma_1, \gamma_2, \dots, \gamma_k\}$  is finite, we may write  $\bigsqcup_{\gamma \in C} \gamma$  as  $\gamma_1 \sqcup \gamma_2 \sqcup \dots \sqcup \gamma_k$ .

**Proposition 4.5.** Let  $\alpha \in P(X)$  with  $\alpha \neq 0$ . Then there exists a unique set  $C$  of pairwise completely disjoint, connected transformations contained in  $\alpha$  such that  $\alpha = \bigsqcup_{\gamma \in C} \gamma$ .

*Proof.* Define a relation  $R$  on  $\text{dom}(\alpha)$  by:  $(x, y) \in R$  if  $x\alpha^k = y\alpha^m \neq \diamond$  for some integers  $k, m \geq 0$ . It is clear that  $R$  is an equivalence relation on  $\text{dom}(\alpha)$ . Let  $J$  be a complete set of representatives of the equivalence classes of  $R$ . For every  $x \in J$ , let  $\gamma_x = \alpha|_{xR}$ , where  $xR$  is the  $R$ -equivalence class of  $x$ . By the definition of  $R$ , each such  $\gamma_x$  is connected, and  $\gamma_x$  and  $\gamma_y$  are completely disjoint for all  $x, y \in J$  with  $x \neq y$ . Then the set  $C = \{\gamma_x : x \in J\}$  consists of pairwise completely disjoint, connected transformations contained in  $\alpha$ , and  $\alpha = \bigsqcup_{\gamma \in C} \gamma$ .

Suppose  $D$  is any set of pairwise completely disjoint, connected transformations contained in  $\alpha$  such that  $\alpha = \bigsqcup_{\delta \in D} \delta$ . Let  $\delta \in D$  and let  $y \in \text{dom}(\delta)$ . Then  $y \in xR$  for some  $x \in J$ . We want to prove that  $\delta = \gamma_x$ . Let  $z \in \text{dom}(\delta)$ . Since  $\delta$  is connected,  $y\delta^k = z\delta^m \neq \diamond$  for some  $k, m \geq 0$ . But then, since  $\delta$  is contained in  $\alpha$ , we have  $y\alpha^k = z\alpha^m \neq \diamond$ . Hence  $(y, z) \in R$ , and so  $z \in yR = xR = \text{dom}(\gamma_x)$ . We have proved that  $\text{dom}(\delta) \subseteq \text{dom}(\gamma_x)$ .

Suppose to the contrary that  $\text{dom}(\gamma_x)$  is not included in  $\text{dom}(\delta)$ , that is, that there is  $w \in \text{dom}(\gamma_x)$  such that  $w \notin \text{dom}(\delta)$ . Since  $\gamma_x$  is connected,  $w\gamma_x^p = y\gamma_x^q \neq \diamond$  for some  $p, q \geq 0$ . Let



$y_i = y\gamma_x^i = y\alpha^i$  and  $w_j = w\gamma_x^j = w\alpha^j$  for  $i = 0, 1, \dots, p$  and  $j = 0, 1, \dots, q$ . Then  $y_p = w_q$  and let  $u = y_p = w_q$ . With this notation, in the digraph  $\Gamma(\alpha)$ , we have

$$y = y_0 \rightarrow y_1 \rightarrow \dots \rightarrow y_p = u \text{ and } w = w_0 \rightarrow w_1 \rightarrow \dots \rightarrow w_q = u.$$

We claim that  $\{y_0, y_1, \dots, y_{p-1}\} \subseteq \text{dom}(\delta)$ . If not, then, since  $y_0 = y \in \text{dom}(\delta)$ , there would be  $i \in \{0, \dots, p-2\}$  such that  $y_i \in \text{dom}(\delta)$  and  $y_{i+1} \notin \text{dom}(\delta)$ . But  $y_{i+1} \in \text{dom}(\alpha)$ , and so  $y_{i+1} \in \text{dom}(\delta_1)$  for some  $\delta_1 \in D$ . We would then have  $\delta \neq \delta_1$  and  $y_{i+1} \in \text{span}(\delta) \cap \text{span}(\delta_1)$ , which is impossible since  $\delta$  and  $\delta_1$  are completely disjoint. The claim has been proved. Since  $w \in \text{dom}(\gamma_x) \subseteq \text{dom}(\alpha)$ , there is  $\delta_2 \in D$  such that  $w \in \text{dom}(\delta_2)$ . By the foregoing argument applied to  $\delta_2$  and  $\{w_0, w_1, \dots, w_{q-1}\}$ , we obtain  $\{w_0, w_1, \dots, w_{q-1}\} \subseteq \text{dom}(\delta_2)$ . Thus

$$y_{p-1}\delta = y_{p-1}\alpha = y_p = u = w_q = w_{q-1}\alpha = w_{q-1}\delta_2.$$

Thus we have  $\delta \neq \delta_2$  with  $u \in \text{im}(\delta) \cap \text{im}(\delta_2)$ , which is a contradiction since  $\delta$  and  $\delta_2$  are completely disjoint. We have proved that  $\text{dom}(\gamma_x) \subseteq \text{dom}(\delta)$ , and so  $\text{dom}(\delta) = \text{dom}(\gamma_x)$ . Now for all  $v \in \text{dom}(\delta) = \text{dom}(\gamma_x)$ , we have  $v\delta = v\alpha = v\gamma_x$ , and so  $\delta = \gamma_x \in C$ . We have proved that  $D \subseteq C$ .

For the reverse inclusion, let  $\gamma_x$  be an arbitrary element of  $C$ . Select  $y \in \text{dom}(\gamma_x)$ . Then, there is  $\delta \in D$  such that  $y \in \text{dom}(\delta)$ . By the foregoing argument, we have  $\delta = \gamma_x$ , and so  $\gamma_x \in D$ . Hence  $C \subseteq D$ , and so  $D = C$ . We have proved that the set  $C$  is unique, which completes the proof.  $\square$

Any element of the set  $C$  from Proposition 4.5 will be called a *connected component* of  $\alpha$ . We note that the connected components of  $\alpha$  correspond to the connected components of the underlying undirected graph of  $\Gamma(\alpha)$  that are not isolated vertices.

**Definition 4.6.** Let  $\alpha \in P(X)$  and let  $\mu$  be a basic partial transformation contained in  $\alpha$ . We say that  $\mu$  is *maximal* in  $\alpha$  if for every  $x \in \text{span}(\mu)$ ,  $x \notin \text{dom}(\mu)$  implies  $x \notin \text{dom}(\alpha)$ , and  $x \notin \text{im}(\mu)$  implies  $x \notin \text{im}(\alpha)$ . Note that if  $\mu$  is a cycle or a double ray, then  $\mu$  is always maximal in  $\alpha$ .

For example, consider  $\alpha = [3456\dots] \sqcup [0125] \in P(\mathbb{Z})$ . Then  $\alpha$  contains infinitely many right rays, for example  $[2567\dots]$ , but only two of them, namely  $[3456\dots]$  and  $[012567\dots]$  are maximal. Also,  $\alpha$  contains infinitely many chains, for example  $[3456]$ , but none of them is maximal.

We will now establish which combinations of basic partial transformations can occur in a connected element of  $P(X)$ .

**Lemma 4.7.** *Let  $\gamma \in P(X)$  be connected.*

- (1) *If  $\gamma$  has a cycle  $(x_0 x_1 \dots x_{k-1})$ , then for every  $x \in \text{dom}(\gamma)$ ,  $x\gamma^m = x_0$  for some  $m \geq 0$ .*
- (2) *If  $\gamma$  has a right ray  $[x_0 x_1 x_2 \dots]$  or a double ray  $\langle \dots x_{-1} x_0 x_1 \dots \rangle$ , then for every  $x \in \text{dom}(\gamma)$ ,  $x\gamma^m = x_i$  for some  $m, i \geq 0$ .*
- (3) *If  $\gamma$  has a maximal chain  $[x_k \dots x_1 x_0]$  or a maximal left ray  $\langle \dots x_2 x_1 x_0 \rangle$ , then for every  $x \in \text{span}(\gamma)$ ,  $x\gamma^m = x_0$  for some  $m \geq 0$ .*

*Proof.* Suppose  $\gamma$  has a cycle  $(x_0 x_1 \dots x_{k-1})$  and let  $x \in \text{dom}(\gamma)$ . Since  $\gamma$  is connected,  $x\gamma^p = x_0\gamma^q$  for some  $p, q \geq 0$ . Since  $x_0$  lies on the cycle  $(x_0 x_1 \dots x_{k-1})$ , we may assume that  $0 \leq q \leq k-1$ . Thus for  $m = p + k - q$ , we have

$$x\gamma^m = x\gamma^{p+k-q} = (x\gamma^p)\gamma^{k-q} = (x_0\gamma^q)\gamma^{k-q} = x_0\gamma^{k-q} = x_0.$$

Suppose  $\gamma$  has a right ray  $[x_0 x_1 x_2 \dots]$  and let  $x \in \text{dom}(\gamma)$ . Since  $\gamma$  is connected,  $x\gamma^m = x_0\gamma^i = x_i$  for some  $m, i \geq 0$ . A proof in the case of a double ray is the same.

Suppose  $\gamma$  has a chain  $[x_k \dots x_1 x_0]$  and let  $x \in \text{span}(\gamma)$ . Since  $\gamma$  is connected,  $x\gamma^p = x_0\gamma^q \neq \diamond$  for some  $p, q \geq 0$ . Note that  $q$  must be 0 (since  $x_0\gamma^q = \diamond$  for every  $q \geq 1$ ). Thus  $x\gamma^p = x_0\gamma^0 = x_0$ . The proof in the case of a maximal left ray is the same.  $\square$

**Proposition 4.8.** *Let  $\gamma \in P(X)$  be connected.*

- (1) *If  $\gamma$  has a cycle, then the cycle is unique and  $\gamma$  does not have any double rays or right rays or maximal chains or maximal left rays.*
- (2) *If  $\gamma$  has a double ray, then it does not have any maximal chains or maximal left rays.*
- (3) *If  $\gamma$  has a right ray, but no double rays, then it has a maximal right ray and it does not have any left rays or maximal chains.*
- (4) *If  $\gamma$  has a chain, but no cycles or rays, then it has a maximal chain.*
- (5) *If  $\gamma$  has a left ray, but no cycles or double rays, then it has a maximal left ray.*

*Proof.* Suppose that  $\gamma$  has a cycle, and let  $\theta$  and  $\vartheta$  be cycles in  $\gamma$ , say  $\theta = (x_0 x_1 \dots x_{k-1})$ . Let  $y \in \text{dom}(\vartheta)$ . By Lemma 4.7,  $y\gamma^p = x_0$  for some  $p \geq 0$ . Thus  $x_0$  lies on  $\vartheta$ , so we may write  $\vartheta = (y_0 y_1 \dots y_{m-1})$  with  $y_0 = x_0$ . We may assume that  $k \leq m$ . But then  $x_i = x_0\gamma^i = y_0\gamma^i = y_i$  for every  $i \in \{0, \dots, k-1\}$  and  $y_{k-1}\gamma = x_{k-1}\gamma = x_0 = y_0$ , that is,  $\theta = \vartheta$ .

Suppose that  $\gamma$  with a cycle  $(x_0 x_1 \dots x_{k-1})$  also has a double ray, say  $\langle \dots y_{-1} y_0 y_1 \dots \rangle$ . By Lemma 4.7,  $y_0\gamma^m = x_0$  for some  $m \geq 0$ . But then  $y_0\gamma^{m+k} = (y_0\gamma^m)\gamma^k = x_0\gamma^k = x_0 = y_0$ , which is a contradiction since  $y_0\gamma^{m+k} = y_{m+k} \neq y_0$  (since  $m \geq 0$  and  $k \geq 1$ ). Thus  $\gamma$  does not have a double ray. This completes the proof of (1) since a connected  $\gamma$  with a cycle cannot have any terminal vertices, and hence cannot have any maximal chains or maximal left rays. Statement (2) also follows since a connected  $\gamma$  with a double ray cannot have terminal vertices either.

Let  $\eta = [x_0 x_1 x_2 \dots]$  be a right ray in  $\alpha$ . If  $\eta$  is not maximal, then  $x_{-1}\gamma = x_0$  for some  $x_{-1} \in X \setminus \{x_0, x_1, \dots\}$ . (If  $x_{-1} = x_i$  for some  $i \geq 0$ , then  $\gamma$  would have a cycle, which is impossible by (1).) Thus  $\eta_1 = [x_{-1} x_0 x_1 x_2 \dots]$  is a right ray in  $\alpha$ . If  $\eta_1$  is not maximal, then  $x_{-2}\gamma = x_{-1}$  for some  $x_{-2} \in X \setminus \{x_{-1}, x_0, x_1, \dots\}$ , and so  $\eta_2 = [x_{-2} x_{-1} x_0 x_1 x_2 \dots]$  is a right ray in  $\alpha$ . Continuing this way, we must arrive at a maximal right ray in  $\alpha$  (after finitely many steps) since otherwise  $\alpha$  would have a double ray. This completes the proof of (3) since a connected  $\gamma$  with a right ray cannot have any terminal vertices.

To prove (4), let  $\lambda = [x_0 x_1 \dots x_k]$  be a chain in  $\alpha$ . If  $x_0 \in \text{im}(\alpha)$ , then, since  $\alpha$  has no left rays, we can use the argument as in the proof of (3) for a right ray to extend  $\lambda$  to a chain  $\lambda' = [x_{-m} \dots x_{-1} x_0 x_1 \dots x_k]$  such that  $x_{-m} \notin \text{im}(\alpha)$ . Similarly, since  $\alpha$  has no right rays or cycles, we can extend  $\lambda'$  to a chain  $\lambda'' = [x_{-m} x_{-m+1} \dots x_{-1} x_0 x_1 \dots x_k x_{k+1} \dots x_{k+p}]$  such that  $x_{k+p} \notin \text{dom}(\alpha)$ . Then  $\lambda''$  is a maximal chain in  $\alpha$ . We have proved (4). The proof of (5) is similar.  $\square$

**Remark 4.9.** It follows from Proposition 4.8 that as far as the types of basic transformations go, a connected  $\gamma \in P(X)$  can contain one of the following.

- (1) A single cycle and no double rays or right rays or maximal chains or maximal left rays (see Figure 4.1);
- (2) A double ray but no cycles or maximal chains or maximal left rays (see Figure 4.2);
- (3) A maximal right ray but no cycles or double rays or left rays or maximal chains (see Figure 4.3);
- (4) A maximal left ray but no cycles or double rays or right rays (see Figure 4.4 and Definition 4.10);
- (5) A maximal chain but no cycles or rays (see Figure 4.5 and Definition 4.10).

We note that the uniqueness applies only to a cycle. A connected  $\gamma$  can have any number (finite or infinite) of (maximal) chains or (maximal) rays of any type.




Figure 4.1: A connected partial transformation with a cycle.




Figure 4.2: A connected partial transformation with a double ray.

For our purposes, it will not be necessary to distinguish connected partial transformations that have double rays only or left rays only. (In other words, if a connected  $\gamma \in P(X)$  has a double ray, then it will not matter whether it has a maximal right ray as well; similarly, if it has a maximal left ray, then it will not matter whether it has a maximal chain as well.) However, we will need to distinguish connected transformations that have right rays only, and connected transformations that have chains only.

**Definition 4.10.** Let  $\gamma \in P(X)$  be connected. If  $\gamma$  satisfies (3) of Remark 4.9, we will say that  $\gamma$  is of (or has) *type rro* (“right rays only”). If  $\gamma$  satisfies (5) of Remark 4.9, we will say that  $\gamma$  is of *type cho* (“chains only”).

**Lemma 4.11.** *Let  $\gamma \in P(X)$  be connected such that  $\gamma$  contains a maximal left ray or it is of type cho. Then  $\gamma$  contains a unique terminal vertex.*

*Proof.* Since  $\gamma$  contains a maximal left ray or a maximal chain, it contains a terminal vertex. Suppose  $x$  and  $y$  are terminal vertices in  $\gamma$ . Since  $\gamma$  is connected,  $x\gamma^k = y\gamma^m \neq \diamond$  for some  $k, m \geq 0$ . But since  $x$  and  $y$  are terminal, this is only possible when  $k = m = 0$ . Thus  $x = y$ .  $\square$




Figure 4.3: A connected partial transformation of type *rro*.




Figure 4.4: A connected partial transformation with a maximal left ray.

**Definition 4.12.** Let  $\gamma \in P(X)$  be connected such that  $\gamma$  has a maximal left ray or is of type *cho*. The unique terminal vertex of  $\gamma$  established by Lemma 4.11 will be called the *root* of  $\gamma$ .

For integers  $a$  and  $b$ , we write  $a \mid b$  if  $a$  divides  $b$ , that is, if  $b = ak$  for some integer  $k$ . For integers  $a$  and  $n$  with  $n \geq 1$ , we denote by  $\text{mod}(a, n)$  the unique integer  $r$  in  $\{0, 1, \dots, n-1\}$  such that  $a \equiv r \pmod{n}$ . We note that

$$\text{mod}(a+1, n) = \begin{cases} \text{mod}(a, n) + 1 & \text{if } \text{mod}(a, n) \neq n-1, \\ 0 & \text{if } \text{mod}(a, n) = n-1. \end{cases} \quad (4.1)$$

**Proposition 4.13.** Let  $\gamma, \delta \in P(X)$  be connected such that  $\gamma$  has a cycle  $(x_0 x_1 \dots x_{k-1})$ . Then  $\Gamma(\gamma)$  is *rp-homomorphic* to  $\Gamma(\delta)$  if and only if  $\delta$  has a cycle  $(y_0 y_1 \dots y_{m-1})$  such that  $m \mid k$ .

*Proof.* Suppose there is an *rp-homomorphism*  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ . Let  $y_i = x_i \phi$  for  $i = 0, 1, \dots, k-1$ . Then  $y_0 \xrightarrow{\delta} y_1 \xrightarrow{\delta} \dots \xrightarrow{\delta} y_{k-1} \xrightarrow{\delta} y_0$ , and so  $y_0 \delta^k = y_0$ . Let  $m$  be the smallest integer in  $\{1, 2, \dots, k\}$  such that  $y_0 \delta^m = y_0$ . Then  $(y_0 y_1 \dots y_{m-1})$  is a cycle in  $\delta$ . By the




Figure 4.5: A connected partial transformation of type *cho*.

Division Algorithm,  $k = mq + r$  for some  $q, r \in \mathbb{N}$  with  $0 \leq r < m$ . Since  $y_0 \delta^m = y_0$ , we have  $y_0 \delta^{mq} = y_0$ , and so  $y_0 = y_0 \delta^k = (y_0 \delta^{mq}) \delta^r = y_0 \delta^r$ . Thus  $r = 0$  by the definition of  $m$ , and so  $k = mq$ , that is,  $m \mid k$ .

Conversely suppose that  $\delta$  has a desired cycle. We will define an rp-homomorphism  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$  such that  $\text{dom}(\phi) = \text{dom}(\gamma)$  and  $\text{im}(\phi) = \{y_0, y_1, \dots, y_{m-1}\}$ . (Note that  $\text{dom}(\gamma) = \text{span}(\gamma)$  since  $\gamma$  has a cycle.) For  $x \in \text{dom}(\gamma)$ , let  $p_x$  be the smallest nonnegative integer such that  $x\gamma^{p_x} = x_0$  (such  $p_x$  exists by Lemma 4.7), and let  $q_x = \text{mod}(-p_x, m)$ . Define  $\phi$  on  $\text{dom}(\gamma)$  by  $x\phi = y_{q_x}$ . Suppose  $x \xrightarrow{\gamma} z$ . We consider two possible cases.

**Case 1.**  $x = x_0$ .

Then  $p_x = 0$ ,  $z = x\gamma = x_0\gamma = x_1$ , and  $p_z = k - 1$ . Thus  $q_x = \text{mod}(0, m) = 0$  and  $q_z = \text{mod}(-k + 1, m) = 1$  (since  $m \mid k$ , and so  $-k \equiv 0 \pmod{m}$ ). Hence  $x\phi = y_0 \xrightarrow{\delta} y_1 = z\phi$ .

**Case 2.**  $x \neq x_0$ .

Then, since  $x \xrightarrow{\gamma} z$ , we have  $p_z = p_x - 1$ , and so

$$q_z = \text{mod}(-p_z, m) = \text{mod}(-p_x + 1, m). \quad (4.2)$$

Suppose  $q_x = \text{mod}(-p_x, m) \neq m - 1$ . Then, by (4.1) and (4.2),  $q_z = \text{mod}(-p_x + 1, m) = \text{mod}(-p_x, m) + 1 = q_x + 1$ , and so  $x\phi = y_{q_x} \xrightarrow{\delta} y_{q_x+1} = y_{q_z} = z\phi$ .

Suppose  $q_x = \text{mod}(-p_x, m) = m - 1$ . Then  $-p_x \equiv -1 \pmod{m}$ , and so  $p_x \equiv 1 \pmod{m}$ . Thus  $p_x = tm + 1$  for some integer  $t$ , and so  $p_z = p_x - 1 = tm$ . Hence  $q_z = \text{mod}(-p_z, m) = \text{mod}(-tm, m) = 0$ , and so  $x\phi = y_{q_x} = y_{m-1} \xrightarrow{\delta} y_0 = y_{q_z} = z\phi$ .

Thus, in both cases,  $x\phi \xrightarrow{\delta} x\phi$ , and so  $\phi$  is an rp-homomorphism. (Condition (b) of Definition 3.1 is satisfied since  $\Gamma(\gamma)$  does not have any terminal vertices.)  $\square$

**Lemma 4.14.** *Let  $\gamma, \delta \in P(X)$  be connected such that  $\delta$  has a cycle  $(y_0 y_1 \dots y_{m-1})$ . Suppose  $\gamma$  has a double ray or  $\gamma$  is of type rro. Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$ .*

*Proof.* Suppose  $\gamma$  has a double ray  $\mu = \langle \dots x_{-1} x_0 x_1 \dots \rangle$ . We will define an rp-homomorphism  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$  such that  $\text{dom}(\phi) = \text{dom}(\gamma)$  and  $\text{im}(\phi) = \{y_0, y_1, \dots, y_{m-1}\}$ . For  $x \in \text{dom}(\gamma)$ , let  $p_x$  be the smallest nonnegative integer such that  $x\gamma^{p_x} = x_i$  for some  $i$  (such  $p_x$  exists by Lemma 4.7), and let  $q_x = \text{mod}(i - p_x, m)$ . Define  $\phi$  on  $\text{dom}(\gamma)$  by  $x\phi = y_{q_x}$ . Suppose  $x \xrightarrow{\gamma} z$ . We consider two possible cases.

**Case 1.**  $x = x_i$  for some  $i \in \mathbb{Z}$ .

Then  $p_x = 0$ ,  $z = x\gamma = x_i\gamma = x_{i+1}$ , and  $p_z = 0$ . Thus  $q_x = \text{mod}(i, m)$  and  $q_z = \text{mod}(i+1, m)$ . If  $q_x \neq m-1$ , then  $q_z = q_x + 1$ , and so  $x\phi = y_{q_x} \xrightarrow{\delta} y_{q_x+1} = y_{q_z} = z\phi$ . If  $q_x = m-1$ , then  $q_z = 0$ , and so  $x\phi = y_{q_x} = y_{m-1} \xrightarrow{\delta} y_0 = y_{q_z} = z\phi$ .

**Case 2.**  $x \neq x_i$  for every  $i \in \mathbb{Z}$ .

Then, since  $x \xrightarrow{\gamma} z$ , we have  $p_z = p_x - 1$  with  $x\gamma^{p_x} = z\gamma^{p_z} = i$ , and so

$$q_z = \text{mod}(i - p_z, m) = \text{mod}(i - p_x + 1, m). \quad (4.3)$$

If  $q_x \neq m-1$ , then, by (4.1) and (4.3),  $q_z = \text{mod}(i - p_x + 1, m) = \text{mod}(i - p_x, m) + 1 = q_x + 1$ , and so  $x\phi = y_{q_x} \xrightarrow{\delta} y_{q_x+1} = y_{q_z} = z\phi$ . If  $q_x = m-1$ , then  $q_z = 0$ , and so again  $x\phi \xrightarrow{\delta} z\phi$ .

Hence, since  $\Gamma(\gamma)$  has no terminal vertices,  $\phi$  is an rp-homomorphism. The proof in the case when  $\gamma$  has type *rro* is similar.  $\square$

**Lemma 4.15.** *Let  $\gamma, \delta \in P(X)$  be connected. Suppose that  $\delta$  has a double ray and  $\gamma$  either has a double ray or has type *rro*. Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$ .*

*Proof.* Suppose  $\gamma$  has a double ray, say  $\langle \dots x_{-1} x_0 x_1 \dots \rangle$ , and let  $\langle \dots y_{-1} y_0 y_1 \dots \rangle$  be a double ray in  $\delta$ . We will define an rp-homomorphism  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$  such that  $\text{dom}(\phi) = \text{dom}(\gamma)$  and  $\text{im}(\phi) = \{\dots, y_{-1}, y_0, y_1, \dots\}$ . (Note that  $\text{dom}(\gamma) = \text{span}(\gamma)$  since  $\gamma$  has a double ray or it is of type *rro*.) For  $x \in \text{dom}(\gamma)$ , let  $p_x$  be the smallest nonnegative integer such that  $x\gamma^{p_x} = x_i$  for some integer  $i$ . Define  $\phi$  on  $\text{dom}(\gamma)$  by  $x\phi = y_{i-p_x}$  where  $x\gamma^{p_x} = x_i$ . Suppose  $x \xrightarrow{\gamma} z$ . We consider two possible cases.

**Case 1.**  $x = x_i$  for some integer  $i$ .

Then  $p_x = 0$ ,  $z = x\gamma = x_i\gamma = x_{i+1}$ , and  $p_z = 0$ . Thus

$$x\phi = y_{i-p_x} = y_i \xrightarrow{\delta} y_{i+1} = y_{i+1-p_z} = z\phi.$$

**Case 2.**  $x \neq x_i$  for every integer  $i$ .

Then, since  $x \xrightarrow{\gamma} z$ , we have  $p_z = p_x - 1$  and  $x\gamma^{p_x} = z\gamma^{p_z} = x_i$  for some  $i$ . Thus

$$x\phi = y_{i-p_x} \xrightarrow{\delta} y_{i-p_x+1} = y_{i-p_z} = z\phi.$$

Thus, in both cases,  $x\phi \xrightarrow{\delta} z\phi$ , and so  $\phi$  is an rp-homomorphism since  $\Gamma(\gamma)$  does not have any terminal vertices. The proof in the case when  $\gamma$  has type *rro* is similar.  $\square$

**Lemma 4.16.** *Let  $\gamma, \delta \in P(X)$  be connected. Suppose that  $\delta$  has a maximal left ray and  $\gamma$  either has a maximal left ray or is of type *cho*. Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$ .*

*Proof.* Let  $\langle \dots y_2 y_1 y_0 \rangle$  be a maximal left ray in  $\delta$ . Note that  $y_0$  is the root of  $\delta$ . Let  $x_0$  be the root of  $\gamma$ . We will define an rp-homomorphism  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$  such that  $\text{dom}(\phi) = \text{span}(\gamma)$  and  $\text{im}(\phi) \subseteq \{\dots, y_2, y_1, y_0\}$ . For  $x \in \text{span}(\gamma)$ , let  $p_x$  be the smallest nonnegative integer such that  $x\gamma^{p_x} = x_0$  (such  $p_x$  exists by Lemma 4.7). Define  $\phi$  on  $\text{span}(\gamma)$  by  $x\phi = y_{p_x}$ . If  $x \xrightarrow{\gamma} z$ , then  $p_z = p_x - 1$ , and so  $x\phi = y_{p_x} \xrightarrow{\delta} y_{p_x-1} = y_{p_z} = z\phi$ . Further, the only terminal vertex in  $\Gamma(\gamma)$  is  $x_0$  and  $x_0\phi = y_0$  (since  $p_{x_0} = 0$ ), which is a terminal vertex in  $\Gamma(\delta)$ . Hence  $\phi$  is an rp-homomorphism.  $\square$

**Lemma 4.17.** *Let  $\gamma, \delta \in P(X)$  be connected such that  $\gamma$  is of type *rro*. Suppose  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$ . Then  $\delta$  cannot have a maximal left ray or be of type *cho*.*

*Proof.* Let  $\phi$  be an rp-homomorphism from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ . Select a right ray  $[x_0 x_1 x_2 \dots]$  in  $\gamma$ . Suppose to the contrary that  $\delta$  has a maximal left ray or is of type *cho*. Let  $y_0$  be the root of  $\delta$ . By Lemma 4.7,  $(x_0\phi)\delta^k = y_0$  for some integer  $k \geq 0$ . By Lemma 3.7,  $\gamma\phi = \phi\delta$ , and so  $(x_0\phi)\delta^{k+1} = (x_0\gamma^{k+1})\phi = x_{k+1}\phi$ . But  $(x_0\phi)\delta^{k+1} = (x_0\phi)\delta^k\delta = y_0\delta = \diamond$ , and so  $x_{k+1}\phi = \diamond$ , which is a contradiction. The result follows.  $\square$

**Proposition 4.18.** *Let  $S \leq P(X)$  such that  $S$  is constant rich, and let  $\alpha, \beta \in S$  with  $\alpha \neq 0$ . Then there is an rp-homomorphism  $\phi \in S^1$  from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$  with  $\text{dom}(\phi) = \text{span}(\alpha)$  if and only if*

- (a) *for every connected component  $\gamma$  of  $\alpha$ , there exist a connected component  $\delta$  of  $\beta$  and an rp-homomorphism  $\phi_\gamma \in P(X)$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$  with  $\text{dom}(\phi_\gamma) = \text{span}(\gamma)$ ; and*
- (b)  *$\bigsqcup_{\gamma \in C} \phi_\gamma \in S^1$ , where  $C$  is the collection of connected components of  $\alpha$ .*

*Proof.* Suppose there is an rp-homomorphism  $\phi \in S^1$  from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$  such that  $\text{dom}(\phi) = \text{span}(\alpha)$ . Let  $\gamma$  be a connected component of  $\alpha$  and let  $x \in \text{span}(\gamma)$ . Then, by Proposition 4.5,  $x\phi \in \delta$  for some connected component  $\delta$  of  $\beta$ . We claim that  $(\text{span}(\gamma))\phi \subseteq \text{span}(\delta)$ . Let  $z \in \text{span}(\gamma)$ . Since  $\gamma$  is connected,  $x\alpha^k = x\gamma^k = z\gamma^m = z\alpha^m \neq \diamond$  for some integers  $k, m \geq 0$ . By Lemma 3.7, we have  $\alpha\phi = \phi\beta$ , and so  $(z\phi)\beta^m = (z\alpha^m)\phi = (x\alpha^k)\phi = (x\phi)\beta^k \neq \diamond$ , which implies that  $z\phi$  and  $x\phi$  are in the span of the same connected component of  $\beta$ , that is,  $z\phi \in \text{span}(\delta)$ . The claim has been proved. Let  $\phi_\gamma = \phi|_{\text{span}(\gamma)}$ . Then  $\phi_\gamma$  is an rp-homomorphism from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$  (by the claim and the fact that  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ ),  $\text{dom}(\phi_\gamma) = \text{span}(\gamma)$  (by the definition of  $\phi_\gamma$ ), and  $\bigsqcup_{\gamma \in C} \phi_\gamma = \phi \in S^1$  (by the definition of  $\phi_\gamma$  and the fact that  $\text{dom}(\phi) = \text{span}(\alpha)$ ).

Conversely, suppose that (a) and (b) are satisfied. Let  $\phi = \bigsqcup_{\gamma \in C} \phi_\gamma$ . Note that  $\phi$  is well defined since  $\phi_\gamma$  and  $\phi_{\gamma'}$  are disjoint if  $\gamma \neq \gamma'$ . Suppose  $y \xrightarrow{\alpha} z$ . Then  $y, z \in \text{span}(\gamma)$  for some connected component  $\gamma$  of  $\alpha$ . Thus  $y, z \in \text{dom}(\phi_\gamma)$  and  $y\phi = y\phi_\gamma \xrightarrow{\delta} z\phi_\gamma = z\phi$ , implying  $y\phi \xrightarrow{\beta} z\phi$ . Suppose  $y$  is a terminal vertex in  $\Gamma(\alpha)$  and  $y \in \text{dom}(\phi)$ . Then, there is a unique connected component  $\gamma$  of  $\alpha$  such that  $y$  is a terminal vertex in  $\Gamma(\gamma)$ . Then  $y\phi = y\phi_\gamma$  is a terminal vertex in  $\Gamma(\delta)$ , and so a terminal vertex in  $\Gamma(\beta)$ . Hence  $\phi$  is an rp-homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . Moreover,  $\text{dom}(\phi) = \text{span}(\alpha)$  (by the definition of  $\phi$  and  $\phi \in S^1$  (by (b))).  $\square$

**Lemma 4.19.** *Let  $\alpha, \beta \in P(X)$  be such that  $\Gamma(\alpha)$  is rp-homomorphic to  $\Gamma(\beta)$ . If  $\alpha$  has a cycle of length  $k$ , then  $\beta$  has a cycle of length  $m$  such that  $m \mid k$ .*

*Proof.* It follows immediately from Propositions 4.13 and 4.18.  $\square$

A binary relation  $R$  on a set  $A$  is called *well founded* if every nonempty subset  $B \subseteq A$  contains an  $R$ -minimal element; that is,  $a \in B$  exists such that there is no  $y \in B$  with  $(y, a) \in R$  [21, page 25]. Let  $R$  be a well-founded relation on  $A$ . Then there is a unique function  $\rho$  defined on  $A$  with ordinals as values such that for every  $x \in A$ ,

$$\rho(x) = \sup\{\rho(y) + 1 : (y, x) \in R\}. \quad (4.4)$$

The ordinal  $\rho(x)$  is called the *rank* of  $x$  in  $\langle A, R \rangle$  [21, Theorem 2.27].

For a mapping  $f : A \rightarrow B$  and  $b \in B$ , we denote by  $bf^{-1}$  the preimage of  $b$  under  $f$ .

**Definition 4.20.** Let  $\gamma \in P(X)$  be connected of type *rro* or *cho*. Recall that  $R_\gamma$  is a binary relation on  $\text{span}(\gamma)$  defined by  $(y, x) \in R_\gamma$  if  $y\gamma = x$ . (Note that  $(y, x) \in R_\gamma \Leftrightarrow y \xrightarrow{\gamma} x \Leftrightarrow y \in x\gamma^{-1}$ ). The relation  $R_\gamma$  is well founded since there is no sequence  $\langle x_0, x_1, x_2, \dots \rangle$  such that  $\dots \xrightarrow{\gamma} x_2 \xrightarrow{\gamma} x_1 \xrightarrow{\gamma} x_0$ . (See [21, Lemma 5.5].) For  $x \in \text{span}(\gamma)$ , we will denote the rank of  $x$  in  $\langle \text{span}(\gamma), R_\gamma \rangle$  by  $\rho_\gamma(x)$  (or  $\rho(x)$  if  $\gamma$  is clear from the context).

It follows from (4.4) that for every  $x \in \text{span}(\gamma)$  with  $\rho(x) > 0$ , we have  $\rho(y) < \rho(x)$  for every  $y \in x\gamma^{-1}$ , and if  $\nu = \sup\{\rho(y) : y \in x\gamma^{-1}\}$  then

$$\rho(x) = \begin{cases} \nu + 1 & \text{if } \rho(y) = \nu \text{ for some } y \in x\gamma^{-1}, \\ \nu & \text{if } \rho(y) < \nu \text{ for every } y \in x\gamma^{-1}. \end{cases} \quad (4.5)$$

**Example 4.21.** Let  $X = \{x_0, x_1, x_2, \dots, y_0, y_1, y_2, \dots\}$  and let

$$\gamma = [x_0 x_1 x_2 x_3 \dots] \sqcup [y_0 x_2] \sqcup [y_1 y_2 x_2] \sqcup [y_3 y_4 y_5 x_2] \sqcup [y_6 y_7 y_8 y_9 x_2] \sqcup \dots \in P(X).$$

Then  $\gamma$  is connected of type *rro* and we have:  $\rho(x_0) = 0$ ,  $\rho(x_1) = 1$ , and  $\rho(x_{2+i}) = \omega + i$  for every  $i \geq 0$ , where  $\omega$  is the smallest infinite ordinal. We also have:  $\rho(y_6) = 0$ ,  $\rho(y_7) = 1$ ,  $\rho(y_8) = 2$ , and  $\rho(y_9) = 3$ .

**Example 4.22.** Let  $X = \{y_0, y_1, y_2, \dots\} \cup \bigcup_{i=0}^{\infty} \{z_0^i, z_1^i, z_2^i, \dots\}$ . For every integer  $i \geq 0$ , let

$$\delta_i = [z_1^i z_0^i] \sqcup [z_2^i z_3^i z_0^i] \sqcup [z_4^i z_5^i z_6^i z_0^i] \sqcup [z_7^i z_8^i z_9^i z_{10}^i z_0^i] \sqcup \dots \in P(X).$$

Then each  $\delta_i$  is connected of type *cho* and  $\rho_{\delta_i}(z_0^i) = \omega$ . Further, let

$$\gamma = (\delta_0 \sqcup [z_0^0 y_0]) \sqcup (\delta_1 \sqcup [z_0^1 y_1 y_0]) \sqcup (\delta_2 \sqcup [z_0^2 y_2 y_3 y_0]) \sqcup (\delta_3 \sqcup [z_0^3 y_4 y_5 y_6 y_0]) \sqcup \dots \in P(X).$$

Then  $\gamma$  is connected of type *cho* and  $\rho_\gamma(y_0) = \omega + \omega = 2\omega$ .

We will need the following lemma from the theory of well-founded relations [22, Appendix B].

**Lemma 4.23.** *Let  $R_1$  and  $R_2$  be well-founded relations on  $A_1$  and  $A_2$ , respectively. Suppose a function  $f : A_1 \rightarrow A_2$  is such that for all  $x, y \in A_1$ , if  $(x, y) \in R_1$  then  $(xf, yf) \in R_2$ . Then for all  $x \in A_1$ ,  $\rho(x) \leq \rho(xf)$ .*

**Notation 4.24.** Let  $\gamma \in P(X)$  be connected and let  $x \in \text{span}(\gamma)$ . We denote by  $\downarrow x$  the set of all  $y \in \text{span}(\gamma)$  such that  $x = y\gamma^m$  for some  $m \geq 0$ . If  $x \in \text{im}(\gamma)$ , we denote by  $\gamma_x$  the restriction of  $\gamma$  to  $\downarrow x \setminus \{x\}$ . Note that  $\gamma_x$  is connected and it either contains a maximal left ray or is of type *cho*, and that, in either case,  $x$  is the root of  $\gamma_x$ .

**Lemma 4.25.** *Let  $\gamma, \delta \in P(X)$  be connected such that  $\gamma$  is of type *rro* or *cho* and  $\delta$  is contained in  $\gamma$ . Then for every  $x \in \text{span}(\delta)$ :*

- (1)  $\rho_\delta(x) \leq \rho_\gamma(x)$ ;
- (2) if  $\delta = \gamma_z$  for some  $z \in \text{im}(\gamma)$ , then  $\rho_\delta(x) = \rho_\gamma(x)$ .

*Proof.* First note that  $\delta$  must be of type *rro* or *cho*. Statement (1) follows from Lemma 4.23 with  $f : \text{span}(\delta) \rightarrow \text{span}(\gamma)$  defined by  $xf = x$ . To prove (2), we suppose  $\delta = \gamma_z$  and proceed by well-founded induction [21, Theorem 2.6]. Let  $x \in \text{span}(\delta)$ . The result is true if  $x$  is  $R_\delta$ -minimal since then  $x$  is also  $R_\gamma$ -minimal. Suppose  $\rho_\delta(y) = \rho_\gamma(y)$  for all  $y \in \text{span}(\delta)$  such that  $(y, x) \in R_\delta$ . Then

$$\rho_\delta(x) = \sup\{\rho_\delta(y) + 1 : (y, x) \in R_\delta\} = \sup\{\rho_\gamma(y) + 1 : (y, x) \in R_\gamma\} = \rho_\gamma(x),$$

where the last but one equality follows from the inductive hypothesis and the fact that for  $\delta = \gamma_z$ ,  $x\delta^{-1} = x\gamma^{-1}$  for all  $x \in \text{span}(\delta)$ .  $\square$

**Proposition 4.26.** *Let  $\gamma, \delta \in P(X)$  be connected of type *cho* with roots  $x_0$  and  $y_0$ , respectively. Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  if and only if  $\rho(x_0) \leq \rho(y_0)$ .*

*Proof.* Suppose there is an rp-homomorphism  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ . Then  $\phi : \text{span}(\gamma) \rightarrow \text{span}(\delta)$  with  $x_0\phi = y_0$ , and for all  $x, y \in \text{span}(\gamma)$ , if  $x \xrightarrow{\gamma} y$  then  $x \xrightarrow{\delta} y$ . Thus  $\rho(x_0) \leq \rho(y_0)$  by Lemma 4.23.

Conversely, suppose  $\rho(x_0) \leq \rho(y_0)$ . We will prove that  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by transfinite induction on  $\rho(x_0)$ . Let  $\rho(x_0) = 1$ . Then for every  $z \in \text{dom}(\gamma)$ , we have  $z \xrightarrow{\gamma} x_0$ . Since  $\rho(y_0) \geq \rho(x_0) = 1$ , there is some  $w \in \text{dom}(\delta)$  such that  $w \xrightarrow{\delta} y_0$ . Define  $\phi$  on  $\text{span}(\gamma)$  by:  $x_0\phi = y_0$  and  $z\phi = w$  for every  $z \in \text{dom}(\gamma)$ . Then clearly  $\phi$  is an rp-homomorphism from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ .



Let  $\rho(x_0) = \mu > 1$  and suppose that for all connected  $\gamma_1, \delta_1 \in P(X)$  of type *cho* with roots  $z$  and  $w$ , respectively, if  $\rho(z) < \mu$  and  $\rho(z) \leq \rho(w)$ , then  $\Gamma(\gamma_1)$  is rp-homomorphic to  $\Gamma(\delta_1)$ .

Let  $z \in x_0\gamma^{-1}$  and note that  $\rho(z) < \mu$ . Since  $\rho(y_0) \geq \mu$ , there is  $w_z \in y_0\delta^{-1}$  such that  $\rho(z) \leq \rho(w_z)$ . If  $z \in \text{im}(\gamma)$ , then  $\gamma_z$  and  $\delta_{w_z}$  are connected with  $\rho_{\gamma_z}(z) = \rho(z) \leq \rho(w_z) = \rho_{\delta_{w_z}}(w_z)$ , and so, by the inductive hypothesis, there is an rp-homomorphism  $\phi_z$  from  $\Gamma(\gamma_z)$  to  $\Gamma(\delta_{w_z})$ . If  $z \notin \text{im}(\gamma)$  (that is, if  $\downarrow z = \{z\}$ ), we define  $\phi_z$  on  $\downarrow z = \{z\}$  by  $z\phi_z = w_z$ .

Define  $\phi$  on  $\text{span}(\gamma)$  by:  $x_0\phi = y_0$  and  $u\phi = u\phi_z$  if  $u \in \downarrow z$  for some  $z \in x_0\gamma^{-1}$ . Then  $\phi$  is well-defined since the collection  $\{\downarrow z\}_{z \in x_0\gamma^{-1}}$  is a partition of  $\text{dom}(\gamma) (= \text{span}(\gamma) \setminus \{x_0\})$ . Suppose  $u \xrightarrow{\gamma} v$ . If  $v \in \downarrow z$  for some  $z \in x_0\gamma^{-1}$ , then  $u \in \downarrow z$  as well, and so  $u\phi = u\phi_z \xrightarrow{\delta} v\phi_z = v\phi$ . If  $v = x_0$ , then  $u = z \in x_0\gamma^{-1}$ , and so  $u\phi = z\phi = z\phi_z = w_z \xrightarrow{\delta} y_0 = x_0\phi = v\phi$ . Hence, since  $x_0\phi = y_0$  and  $x_0$  is the unique terminal vertex of  $\Gamma(\gamma)$ ,  $\phi$  is an rp-homomorphism from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ .  $\square$

**Definition 4.27.** Let  $\langle a_n \rangle_{n \geq 0}$  and  $\langle b_n \rangle_{n \geq 0}$  be sequences of ordinals (indexed by nonnegative integers  $n$ ). We say that  $\langle b_n \rangle$  *dominates*  $\langle a_n \rangle$  if there is  $k \geq 0$  such that

$$b_{k+n} \geq a_n \text{ for every } n \geq 0.$$

**Notation 4.28.** Let  $\gamma \in P(X)$  be connected of type *rro* and let  $\eta = [x_0 x_1 x_2 \dots]$  be a maximal right ray in  $\gamma$ . We denote by  $\langle \eta_n^\gamma \rangle_{n \geq 0}$  the sequence of ordinals such that

$$\eta_n^\gamma = \rho_\gamma(x_n) \text{ for every } n \geq 0.$$

For example, for  $\gamma$  from Example 4.21 and the right ray  $\eta = [x_0 x_1 x_2 \dots]$  in  $\gamma$ , the sequence  $\langle \eta_n^\gamma \rangle$  is  $\langle 0, 1, \omega, \omega + 1, \omega + 2, \omega + 3, \dots \rangle$ .

**Proposition 4.29.** Let  $\gamma, \delta \in P(X)$  be connected of type *rro*. Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  if and only if there are maximal right rays  $\eta$  in  $\gamma$  and  $\xi$  in  $\delta$  such that  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$ .

*Proof.* Suppose there is an rp-homomorphism  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ . Select a maximal right ray  $\eta = [x_0 x_1 x_2 \dots]$  in  $\gamma$  (possible by Proposition 4.8.) Then  $x_0\phi \xrightarrow{\delta} x_1\phi \xrightarrow{\delta} x_2\phi \xrightarrow{\delta} \dots$ , and so, since  $\delta$  does not have any double rays, there is  $w \in \text{dom}(\delta) - \text{im}(\delta)$  such that  $w\delta^k = x_0\phi$  for some  $k \geq 0$ . Thus

$$\xi = [y_0 = w \ y_1 = w\delta \ \dots \ y_{k-1} = w\delta^{k-1} \ y_k = w\delta^k = x_0\phi \ y_{k+1} = x_1\phi \ y_{k+2} = x_2\phi \ \dots]$$

is a maximal right ray in  $\delta$ . For every  $n \geq 0$ , the mapping  $\phi|_{\downarrow x_n}$  is an rp-homomorphism from  $\Gamma(\gamma_{x_n})$  to  $\Gamma(\delta_{y_{k+n}})$  (see Notation 4.24). Thus for every  $n \geq 0$ , we have  $\rho_{\gamma_{x_n}}(x_n) \leq \rho_{\delta_{y_{k+n}}}(y_{k+n})$  by Proposition 4.26, and so  $\rho(x_n) \leq \rho(y_{k+n})$  by Lemma 4.25. Hence  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$ .

Conversely, suppose there are maximal right rays  $\eta = [x_0 x_1 x_2 \dots]$  in  $\gamma$  and  $\xi = [y_0 y_1 y_2 \dots]$  in  $\delta$  such that  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$ , that is, there is  $k \geq 0$  such that  $\xi_{k+n}^\delta \geq \eta_n^\gamma$  for every  $n \geq 0$ . We define a collection  $\{B_n\}_{n \geq 0}$  of subsets of  $\text{span}(\gamma)$  by

$$B_0 = \{x_0\}, \quad B_n = \downarrow x_n - \downarrow x_{n-1} \text{ for } n \geq 1.$$

Since  $\gamma$  is connected,  $\{B_n\}_{n \geq 0}$  is a partition of  $\text{span}(\gamma)$ .

We will now define an rp-homomorphism  $\phi$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$  by defining  $\phi$  on  $B_n$  for every  $n \geq 0$ . First, we set  $x_0\phi = y_k$ . Let  $n \geq 1$ . If  $B_n = \{x_n\}$ , we set  $x_n\phi = y_{k+n}$ . Suppose  $|B_n| \geq 2$ . Let  $\gamma_n = \gamma|_{B_n \setminus \{x_n\}}$  and  $\delta_n = \delta_{y_{k+n}}$ . Then  $\gamma_n$  and  $\delta_n$  are connected of type *cho* with roots  $x_n$  and  $y_{k+n}$ , respectively. By Lemma 4.25,

$$\rho_{\gamma_n}(x_n) \leq \rho_\gamma(x_n) = \eta_n^\gamma \leq \xi_{k+n}^\delta = \rho_\delta(y_{k+n}) = \rho_{\delta_n}(y_{k+n}).$$

Thus, by Proposition 4.26, there is an rp-homomorphism  $\phi_n$  from  $\Gamma(\gamma_n)$  to  $\Gamma(\delta_n)$ . Note that  $x_n\phi_n = y_{k+n}$ . We define  $\phi$  on  $B_n$  by  $x\phi = x\phi_n$ .

Suppose  $x \xrightarrow{\gamma} z$ . Then  $z \in B_n$  for some  $n \geq 0$ . If  $x \in B_n$ , then  $x\phi = x\phi_n \xrightarrow{\delta} z\phi_n = z\phi$  since  $\phi_n$  is an rp-homomorphism from  $\Gamma(\gamma_n)$  to  $\Gamma(\delta_n)$ . If  $x \notin B_n$ , then we must have  $x = x_{n-1}$  and  $z = x_n$ , and so  $x\phi = x_{n-1}\phi = y_{k+n-1} \xrightarrow{\delta} y_{k+n} = x_n\phi = z\phi$ .

Hence, in all cases, if  $x \xrightarrow{\gamma} z$  then  $x\phi \xrightarrow{\delta} z\phi$ . Thus, since  $\Gamma(\gamma)$  does not have any terminal vertices,  $\phi$  is an rp-homomorphism from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ .  $\square$

The following lemma will be needed in the next section.

**Lemma 4.30.** *Let  $\gamma, \delta \in P(X)$  be of type rro. Let  $\eta$  be a maximal right ray in  $\gamma$  and  $\xi$  be a maximal right ray in  $\delta$  such that  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$ . Then for every maximal right ray  $\eta_1$  in  $\gamma$  and every maximal right ray  $\xi_1$  in  $\delta$   $\langle (\xi_1)_n^\delta \rangle$  dominates  $\langle (\eta_1)_n^\gamma \rangle$ .*

*Proof.* Since  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$ , there is an integer  $k \geq 0$  such that

$$\xi_{k+n}^\delta \geq \eta_n^\gamma \text{ for every } n \geq 0.$$

Let  $\eta = [x_0 x_1 x_2 \dots]$  and  $\xi = [y_0 y_1 y_2 \dots]$ . Let  $\eta_1 = [w_0 w_1 w_2 \dots]$  and  $\xi_1 = [z_0 z_1 z_2 \dots]$  be arbitrary maximal right rays in  $\gamma$  and  $\delta$ , respectively. Since  $\gamma$  and  $\delta$  are connected, there are integers  $l, q, m, p \geq 0$  such that  $x_l = x_0 \gamma^l = w_0 \gamma^q = w_q$  and  $y_m = y_0 \delta^m = z_0 \delta^p = z_p$ . We may assume that  $m \geq k$ . Then for every  $n \geq 0$ ,

$$\begin{aligned} (\xi_1)_{(p+l)+n}^\delta &= \rho_\delta(z_{p+(l+n)}) = \rho_\delta(y_{m+(l+n)}) \geq \rho_\delta(y_{k+(l+n)}) = \xi_{k+(l+n)}^\delta \geq \eta_{l+n}^\gamma, \text{ and} \\ \eta_{l+n}^\gamma &= \rho_\gamma(x_{l+n}) = \rho_\gamma(w_{q+n}) \geq \rho_\gamma(w_n) = (\eta_1)_n^\gamma. \end{aligned}$$

Hence  $\langle (\xi_1)_n^\delta \rangle$  dominates  $\langle (\eta_1)_n^\gamma \rangle$ .  $\square$

## 5 Conjugacy in $P(X)$

In this section we characterize the conjugacy  $\sim_c$  in the semigroup  $P(X)$  of partial transformations on any nonempty set  $X$  (finite or infinite).

In  $P(X)$  and, more generally, in any a constant rich subsemigroup  $S$  of  $P(X)$ , the conjugacy relation  $\sim_c$  can be reformulated, as a consequence of Lemma 3.5(1), in the following way: given any  $\alpha, \beta \in S$ , we have  $\alpha \sim_c \beta$  in  $S$  if and only if there exist  $\phi, \psi \in S^1$  such that  $\alpha\phi = \phi\beta$  and  $\beta\psi = \psi\alpha$ , with  $\text{dom}(\alpha\phi) = \text{dom}(\alpha)$  and  $\text{dom}(\beta\psi) = \text{dom}(\beta)$ . Notice that the semigroup  $P(X)$  can be regarded as a left restriction semigroup with respect to the set of partial identities  $E = \{\text{id}_Y : Y \subseteq X\}$  (see [19] for a survey). Hence  $P(X)$  is equipped with a unary operation  $+$  assigning to any  $\alpha \in P(X)$  the element  $\alpha^+ = \text{id}_{\text{dom}(\alpha)}$ . Any subsemigroup  $S$  of  $P(X)$  closed under  $+$  is called a left restriction semigroup. If  $S$  is a left restriction semigroup that is also constant rich, then for all  $\alpha, \beta \in S$ ,

$$\alpha \sim_c \beta \Leftrightarrow \exists \phi, \psi \in S^1 : \alpha\phi = \phi\beta \text{ and } \beta\psi = \psi\alpha, \text{ with } (\alpha\phi)^+ = \alpha^+ \text{ and } (\beta\psi)^+ = \beta^+.$$

We now proceed to characterize the conjugacy relation  $\sim_c$  in  $P(X)$  in terms of the basic partial transformations.

**Definition 5.1.** Let  $M$  be a nonempty subset of the set  $\mathbb{Z}_+$  of positive integers. Then  $M$  is partially ordered by the relation  $|$  (divides). Order the elements of  $M$  according to the usual “less than” relation:  $m_1 < m_2 < m_3 < \dots$ . We define a subset  $\text{sac}(M)$  of  $M$  as follows: for every integer  $n$ ,  $1 \leq n < |M| + 1$ ,

$$m_n \in \text{sac}(M) \Leftrightarrow (\forall_{i < n}) m_n \text{ is not a multiple of } m_i.$$

The set  $\text{sac}(M)$  is a maximal antichain of the poset  $(M, |)$ . We will call  $\text{sac}(M)$  the *standard antichain* of  $M$ .

For example, if  $M = \{4, 6, 8, 10, 18\}$  then  $\text{sac}(M) = \{4, 6, 10\}$ ; if  $M = \{1, 2, 4, 8, 16, 32, \dots\}$  then  $\text{sac}(M) = \{1\}$ .

**Definition 5.2.** Let  $\alpha \in P(X)$  such that  $\alpha$  contains a cycle. Let

$$M = \{n \in \mathbb{Z}_+ : (\exists x \in \text{dom}(\alpha)) \ x\alpha^n = x \text{ and } x\alpha^i \neq x \text{ for every } i, 1 \leq i < n\}.$$

Note that  $M$  is the set of the lengths of cycles in  $\alpha$ . The standard antichain of  $(M, |)$  will be called the *cycle set* of  $\alpha$  and denoted by  $\text{cs}(\alpha)$ . We agree that  $\text{cs}(\alpha) = \emptyset$  if  $\alpha$  has no cycles.

**Theorem 5.3.** Let  $\alpha, \beta \in P(X)$ . Then  $\alpha \sim_c \beta$  in  $P(X)$  if and only if  $\alpha = \beta = 0$  or  $\alpha, \beta \neq 0$  and the following conditions are satisfied:

- (1)  $\text{cs}(\alpha) = \text{cs}(\beta)$ ;
- (2)  $\alpha$  has a double ray but not a cycle  $\Leftrightarrow \beta$  has a double ray but not a cycle;
- (3a) if  $\alpha$  has a connected component  $\gamma$  of type *rro*, but no cycles or double rays, then  $\beta$  has a connected component  $\delta$  of type *rro*, but no cycles or double rays, and  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$  for some maximal right rays  $\eta$  in  $\gamma$  and  $\xi$  in  $\delta$ ;
- (3b) if  $\beta$  has a connected component  $\delta$  of type *rro*, but no cycles or double rays, then  $\alpha$  has a connected component  $\gamma$  of type *rro*, but no cycles or double rays, and  $\langle \eta_n^\gamma \rangle$  dominates  $\langle \xi_n^\delta \rangle$  for some maximal right rays  $\xi$  in  $\delta$  and  $\eta$  in  $\gamma$ ;
- (4)  $\alpha$  has a maximal left ray  $\Leftrightarrow \beta$  has a maximal left ray;
- (5a) if  $\alpha$  has a connected component  $\gamma$  of type *cho* with root  $x_0$ , but no maximal left rays, then  $\beta$  has a connected component  $\delta$  of type *cho* with root  $y_0$ , but no maximal left rays, and  $\rho_\gamma(x_0) \leq \rho_\delta(y_0)$ ;
- (5b) if  $\beta$  has a connected component  $\delta$  of type *cho* with root  $y_0$ , but no maximal left rays, then  $\alpha$  has a connected component  $\gamma$  of type *cho* with root  $x_0$ , but no maximal left rays, and  $\rho_\delta(y_0) \leq \rho_\gamma(x_0)$ .

*Proof.* Suppose  $\alpha \sim_c \beta$ . Then, since  $[0]_{\sim_c} = \{0\}$  in every semigroup with 0, either  $\alpha = \beta = 0$  or  $\alpha, \beta \neq 0$ . Suppose  $\alpha, \beta \neq 0$ . Then, by Theorem 3.8, there is an rp-homomorphism  $\phi$  from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . We may assume that  $\text{dom}(\phi) = \text{span}(\alpha)$  (see Remark 3.2).

Suppose  $\alpha$  has a cycle. Then, by Lemma 4.19,  $\beta$  also has a cycle. Let  $n \in \text{cs}(\alpha)$ . Then  $\alpha$  has a cycle of length  $n$ , and so  $\beta$  has a cycle of length  $m$  such that  $m | n$ . By the definition of  $\text{cs}(\beta)$ , there is  $m_1 \in \text{cs}(\beta)$  such that  $m_1 | m$ . Thus  $\beta$  has a cycle of length  $m_1$ , and so  $\alpha$  has a cycle of length  $n_1$  such that  $n_1 | m_1$ , so  $n_1 | m_1 | m | n$ . Since  $\text{cs}(\alpha)$  is an antichain,  $n_1 | n$  and  $n \in \text{cs}(\alpha)$  implies  $n_1 = n$ . Thus  $n = m_1$ , and so  $n \in \text{cs}(\beta)$ . We have proved that  $\text{cs}(\alpha) \subseteq \text{cs}(\beta)$ . Similarly,  $\text{cs}(\beta) \subseteq \text{cs}(\alpha)$ , and so  $\text{cs}(\alpha) = \text{cs}(\beta)$ . By symmetry, if  $\beta$  has a cycle, then  $\alpha$  also has a cycle and  $\text{cs}(\beta) = \text{cs}(\alpha)$ . If neither  $\alpha$  nor  $\beta$  has a cycle, then  $\text{cs}(\alpha) = \text{cs}(\beta) = \emptyset$ . We have proved (1).

Suppose  $\alpha$  has a double ray, say  $\langle \dots x_{-1} x_0 x_1 \dots \rangle$ , but no cycles. Then  $\beta$  does not have a cycle either by Lemma 4.19, and  $\dots \xrightarrow{\beta} x_{-1}\phi \xrightarrow{\beta} x_0\phi \xrightarrow{\beta} x_1\phi \xrightarrow{\beta} \dots$ , where  $\phi$  is a homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . The elements  $\dots, x_{-1}\phi, x_0\phi, x_1\phi, \dots$  are pairwise disjoint (since otherwise  $\beta$  would have a cycle), and so  $\langle \dots x_{-1}\phi x_0\phi x_1\phi \dots \rangle$  is a double ray in  $\beta$ . The converse is true by symmetry. This proves (2).

Suppose that  $\alpha$  has a connected component  $\gamma$  of type *rro*, but neither a cycle nor a double ray. By Proposition 4.18, there is a connected component  $\delta$  of  $\beta$  such that  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$ . By (1) and (2),  $\delta$  does not have a cycle or a double ray. By Lemma 4.17,  $\delta$  does not have a maximal left ray and it is not of type *cho*. Hence  $\delta$  has type *rro*. By Proposition 4.29, there are maximal right rays  $\eta$  in  $\gamma$  and  $\xi$  in  $\delta$  such that  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$ . We have proved (3a). Condition (3b) holds by symmetry.

Suppose  $\alpha$  has a maximal left ray, say  $\langle \dots x_2 x_1 x_0 \rangle$ . Then  $\dots \xrightarrow{\beta} x_2 \phi \xrightarrow{\beta} x_1 \phi \xrightarrow{\beta} x_0 \phi$  and  $x_0 \phi$  is a terminal vertex in  $\Gamma(\beta)$ , which implies that  $\langle \dots x_2 \phi x_1 \phi x_0 \phi \rangle$  is a maximal left ray in  $\beta$ . The converse is true by symmetry. This proves (4).

Suppose  $\alpha$  has a connected component  $\gamma$  of type *cho* with root  $x_0$ , but not a maximal left ray. By Proposition 4.18 and its proof, there is a connected component  $\delta$  of  $\beta$  such that  $\phi_\gamma = \phi|_{\text{span}(\gamma)}$  is an rp-homomorphism from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ . Since  $x_0$  is a terminal vertex in  $\gamma$ ,  $y_0 = x_0 \phi_\gamma$  is a terminal vertex in  $\delta$ . Since  $\beta$  has no maximal left ray (by (3)),  $\delta$  is of type *cho* and  $y_0$  is the root of  $\delta$ . By Proposition 4.26,  $\rho_\gamma(x_0) \leq \rho_\delta(y_0)$ . We have proved (5a). Condition (5b) holds by symmetry.

Conversely, if  $\alpha = \beta = 0$  then  $\alpha \sim_c \beta$ . Suppose that  $\alpha, \beta \neq 0$  and that (1)–(5b) hold. Let  $\gamma$  be a connected component of  $\alpha$ . We will prove that  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  for some connected component  $\delta$  of  $\beta$ .

Suppose  $\gamma$  has a cycle of length  $k$ . Since, by (1),  $\text{cs}(\alpha) = \text{cs}(\beta)$ ,  $\beta$  has a cycle  $\vartheta$  of length  $m$  such that  $m \mid k$ . Let  $\delta$  be the connected component of  $\beta$  containing  $\vartheta$ . Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Proposition 4.13.

Suppose  $\gamma$  has a double ray. If some connected component  $\delta$  of  $\beta$  has a cycle, then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Lemma 4.14. Suppose  $\beta$  does not have a cycle. Then, by (1) and (2), both  $\alpha$  and  $\beta$  have a double ray but not a cycle. Let  $\delta$  be a connected component of  $\beta$  containing a double ray. Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Lemma 4.15.

Suppose  $\gamma$  is of type *rro*. If  $\beta$  has some connected component  $\delta$  with a cycle or a double ray, then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Lemmas 4.14 and 4.15. Suppose  $\beta$  does not have a cycle or a double ray. Then, by (3a), there is a connected component  $\delta$  in  $\beta$  of type *rro* such that  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$  for some maximal right rays  $\eta$  in  $\gamma$  and  $\xi$  in  $\delta$ . Hence  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Proposition 4.29.

Suppose  $\gamma$  has a maximal left ray. Then, by (4), some connected component  $\delta$  of  $\beta$  has a maximal left ray. Then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Lemma 4.16.

Suppose  $\gamma$  is of type *cho* with root  $x_0$ . If  $\beta$  has some connected component  $\delta$  with a maximal left ray, then  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Lemma 4.16. Suppose  $\beta$  does not have a maximal left ray. Then, by (4),  $\alpha$  does not have a maximal left ray, and so, by (5a), there is a connected component  $\delta$  in  $\beta$  of type *cho* with root  $y_0$  such that  $\rho_\gamma(x_0) \leq \rho_\delta(y_0)$ . Hence  $\Gamma(\gamma)$  is rp-homomorphic to  $\Gamma(\delta)$  by Proposition 4.26.

We have proved that for every connected component  $\gamma$  of  $\alpha$ , there exists a connected component  $\delta$  of  $\beta$  and an rp-homomorphism  $\phi_\gamma \in P(X)$  from  $\Gamma(\gamma)$  to  $\Gamma(\delta)$ . We may assume that for every  $\gamma \in C(\alpha)$ ,  $\text{dom}(\phi_\gamma) = \text{span}(\gamma)$ . Hence  $\Gamma(\alpha)$  is rp-homomorphic to  $\Gamma(\beta)$  by Proposition 4.18. By symmetry,  $\Gamma(\beta)$  is rp-homomorphic to  $\Gamma(\alpha)$ , and so  $\alpha \sim_c \beta$  by Theorem 3.8.  $\square$

**Example 5.4.** Let  $X$  be an infinite set containing  $x_0, y_1, y_2, y_3, \dots$  and let  $\alpha, \beta \in P(X)$  be the partial transformations whose digraphs are presented in Figure 5.1. Then  $\alpha$  is connected of type *cho* with root  $x_0$ , and  $\beta = \delta_1 \sqcup \delta_2 \sqcup \delta_3 \sqcup \delta_4 \sqcup \dots$ , where  $\delta_i$  is a chain with root  $y_i$ . We have  $\rho_\gamma(x_0) = \omega$ , where  $\gamma = \alpha$ , and for every integer  $i \geq 1$ ,  $\rho_{\delta_i}(y_i) = i$ . Hence  $\alpha$  and  $\beta$  are not conjugate by (5a) of Theorem 5.3.

**Definition 5.5.** For  $\alpha \in P(X)$ , we define

$$s(\alpha) = \sup\{\rho_\gamma(x_0) : \gamma \text{ is a connected component of } \alpha \text{ of type } \textit{cho} \text{ with root } x_0\},$$

where we agree that  $s(\alpha) = 0$  if  $\alpha$  has no connected component of type *cho*.

Suppose  $\alpha, \beta \in P(X)$  have a connected component of type *cho*, but no cycles or rays. Then, by Theorem 5.3, if  $\alpha \sim_c \beta$  then  $s(\alpha) = s(\beta)$ . However, the converse is not true. Indeed, consider  $\alpha, \beta \in P(X)$  from Example 5.4 (see Figure 5.1). Then  $\alpha$  is connected of type *cho* with the root of order  $\omega$ , and  $\beta$  is a join of connected components of type *cho* (chains) whose roots have orders  $1, 2, 3, 4, \dots$ . Thus  $s(\alpha) = s(\beta) = \omega$ , but  $(\alpha, \beta) \notin \sim_c$  by (5a) of Theorem 5.3. However, if  $X$  is finite and  $\alpha, \beta \in P(X)$  have no cycles, then  $s(\alpha) = s(\beta)$  does imply  $\alpha \sim_c \beta$ .




Figure 5.1: The digraphs of  $\alpha$  and  $\beta$  from Example 5.4.

The transformations of a finite  $P(X)$  have no rays. Hence, Theorem 5.3 gives us the following corollary.

**Corollary 5.6.** *Let  $X$  be finite, and let  $\alpha, \beta \in P(X)$ . Then  $\alpha \sim_c \beta$  if and only if  $\text{cs}(\alpha) = \text{cs}(\beta)$  and  $s(\alpha) = s(\beta)$ .*

**Example 5.7.** Let  $\alpha$  and  $\beta$  be partial transformations whose digraphs are presented in Figures 5.2 and 5.3, respectively. Then  $\text{cs}(\alpha) = \text{cs}(\beta) = \{2, 3\}$  and  $s(\alpha) = s(\beta) = 3$ . Thus  $\alpha \sim_c \beta$  by Corollary 5.6.




Figure 5.2: The digraph of  $\alpha$  from Example 5.7.




Figure 5.3: The digraph of  $\beta$  from Example 5.7.

Using Theorem 5.3, we will count the conjugacy classes in  $P(X)$  for an infinite set  $X$  (Theorem 5.16). We will use the aleph notation for the infinite cardinals, that is, for an ordinal  $\varepsilon$ , we will write  $\aleph_\varepsilon$  for the cardinal indexed by  $\varepsilon$ . If  $\aleph_\varepsilon$  is viewed as an ordinal, we will consistently write  $\omega_\varepsilon$ . This is important because we will need to distinguish between ordinal and cardinal

arithmetic. For example,  $\omega_0 < \omega_0 + 1$  (ordinal arithmetic) but  $\aleph_0 = \aleph_0 + 1$  (cardinal arithmetic). It will be always clear from the context which arithmetic is used.

A cardinal  $\aleph_\varepsilon$  is called *singular* if there is a limit ordinal  $\vartheta < \omega_\varepsilon$  and there is an increasing transfinite sequence  $\langle \lambda_\nu \rangle_{\nu < \vartheta}$  of ordinals  $\lambda_\nu < \omega_\varepsilon$  such that  $\omega_\varepsilon = \sup\{\lambda_\nu : \nu < \vartheta\}$  [20, page 160, Definition 2.1]. (As in [20], “increasing” means “strictly increasing.”) If  $\aleph_\varepsilon$  is not singular, then it is called *regular*.

For any cardinal  $\aleph_\varepsilon$ , the cardinal  $\aleph_{\varepsilon+1}$  is called the *successor* cardinal of  $\aleph_\varepsilon$ . Every successor cardinal is regular [20, page 162, Theorem 2.4]. The following lemma follows immediately from this fact and the definition of a regular cardinal.

**Lemma 5.8.** *Let  $\aleph_{\varepsilon+1}$  be a successor cardinal and let  $A$  be a set of ordinals such that  $|A| < \aleph_{\varepsilon+1}$  and  $\lambda < \omega_{\varepsilon+1}$  for every  $\lambda \in A$ . Then  $\sup\{\lambda : \lambda \in A\} < \omega_{\varepsilon+1}$ .*

To prove the counting theorem, we need a series of lemmas.

**Lemma 5.9.** *Let  $|X| = \aleph_\varepsilon$  and let  $\gamma \in P(X)$  be of type *cho* with root  $x_0$ . Then  $\rho(x_0) < \omega_{\varepsilon+1}$ .*

*Proof.* Let  $x \in \text{span}(\gamma)$ . We will prove that  $\rho(x) < \omega_{\varepsilon+1}$  by well-founded induction. If  $x$  is  $R_\gamma$ -minimal, then  $\rho(x) = 0 < \omega_{\varepsilon+1}$ . Suppose  $\rho(y) < \omega_{\varepsilon+1}$  for every  $y \in x\gamma^{-1}$ . Then  $\rho(x) = \sup\{\rho_\delta(y) + 1 : (y, x) \in R_\gamma\}$  by Lemma 5.8, and the result follows.  $\square$

**Lemma 5.10.** *Let  $|X| = \aleph_\varepsilon$ . Then for every nonzero ordinal  $\mu < \omega_{\varepsilon+1}$ , there is  $\gamma \in P(X)$  of type *cho* with root  $x_0$  such that  $\rho(x_0) = \mu$ .*

*Proof.* Let  $0 < \mu < \omega_{\varepsilon+1}$ . We proceed by transfinite induction. The result is clearly true if  $\mu = 1$ . Let  $\mu > 1$  and suppose that the result is true for every ordinal  $\lambda$  such that  $0 < \lambda < \mu$ .

Fix  $x_0 \in X$ , let  $X_0 = X \setminus \{x_0\}$ , and note that  $|X_0| = \aleph_\varepsilon$ . Since  $\mu < \omega_{\varepsilon+1}$ , we have  $|\mu| \leq \aleph_\varepsilon$ . Thus, since  $\aleph_\varepsilon \cdot \aleph_\varepsilon = \aleph_\varepsilon$  and  $\mu = \{\lambda : \lambda \text{ is an ordinal such that } \lambda < \mu\}$ , there is a collection  $\{X_\lambda\}_{0 < \lambda < \mu}$  of pairwise disjoint subsets of  $X_0$  such that  $|X_\lambda| = \aleph_\varepsilon$  for every  $\lambda$ .

Let  $0 < \lambda < \mu$ . By the inductive hypothesis, there is  $\gamma_\lambda \in P(X_\lambda)$  of type *cho* with root  $x_\lambda$  such that  $\rho(x_\lambda) = \lambda$ . We define  $\gamma \in P(X)$  as follows. Set  $\text{dom}(\gamma) = \bigcup_{0 < \lambda < \mu} \text{span}(\gamma_\lambda)$ . For every  $x \in \text{dom}(\gamma)$ , define

$$x\gamma = \begin{cases} x\gamma_\lambda & \text{if } x \in \text{dom}(\gamma_\lambda), \\ x_0 & \text{if } x = x_\lambda. \end{cases}$$

Then  $\gamma$  is of type *cho*,  $x_0$  is the root of  $\gamma$ , and  $x_0\gamma^{-1} = \{x_\lambda : 0 < \lambda < \mu\}$ . Let  $\nu = \sup\{\rho(y) : y \in x_0\gamma^{-1}\}$ . Then

$$\nu = \sup\{\rho(x_\lambda) : 0 < \lambda < \mu\} = \sup\{\lambda : 0 < \lambda < \mu\},$$

where the last equality is true since  $\rho(x_\lambda) = \lambda$  for every nonzero  $\lambda < \mu$ . Hence, either  $\mu = \nu$  (if  $\mu$  is a limit ordinal) or  $\mu = \nu + 1$  (if  $\nu = \lambda$  for some nonzero  $\lambda < \mu$ ). It follows by (4.5) that  $\rho(x_0) = \mu$ .  $\square$

**Lemma 5.11.** *Let  $|X| = \aleph_\varepsilon$  and let  $\langle a_n \rangle$  be an increasing sequence of ordinals  $a_n < \omega_{\varepsilon+1}$  such that  $a_0 = 0$ . Then there is  $\gamma \in T(X)$  of type *rro* with a maximal right ray  $\eta$  such that  $\langle \eta_n^\gamma \rangle = \langle a_n \rangle$ .*

*Proof.* Since  $|X| = \aleph_\varepsilon$ , there is a collection  $\{X_n\}_{n \geq 0}$  of pairwise disjoint subsets of  $X$  such that  $X_0 = \{x_0\}$  and  $|X_n| = \aleph_\varepsilon$  for every  $n \geq 1$ . Let  $n \geq 1$ . By Lemma 5.10, there is  $\gamma_n \in P(X_n)$  of type *cho* with root  $x_n$  such that  $\rho_{\gamma_n}(x_n) = a_n$ . Define  $\gamma \in T(X)$  by

$$x\gamma = \begin{cases} x\gamma_n & \text{if } x \in \text{dom}(\gamma_n), \\ x_{n+1} & \text{if } x = x_n, \\ x_1 & \text{for any other } x. \end{cases}$$

(See Figure 5.4.) Then  $\gamma$  is of type *rro* (since every  $\gamma_n$  is of type *cho*). By the definition of  $\gamma$ , we have that  $\rho_\gamma(x_0) = 0$  and  $\eta = [x_0 x_1 x_2 \dots]$  is a maximal right ray in  $\gamma$ . We have already noticed

that  $\rho_\gamma(x_0) = 0 = a_0$ . We will prove by induction on  $n$  that  $\rho_\gamma(x_n) = a_n$  for every  $n \geq 1$ . Let  $n = 1$ . Then, since  $a_1 \geq 1$ ,

$$\rho_\gamma(x_1) = \max\{1, \rho_{\gamma_1}(x_1)\} = \max\{1, a_1\} = a_1.$$

Let  $n \geq 1$  and suppose  $\rho_\gamma(x_n) = a_n$ . Then

$$\rho_\gamma(x_{n+1}) = \max\{\rho_\gamma(x_n) + 1, \rho_{\gamma_{n+1}}(x_{n+1})\} = \max\{a_n + 1, a_{n+1}\} = a_{n+1},$$

where the last equality is true since  $\langle a_n \rangle$  is increasing, and so  $a_{n+1} > a_n$ . This concludes the inductive argument. Thus  $\eta_n^\gamma = \rho_\gamma(x_n) = a_n$  for every  $n \geq 0$ , which completes the proof.  $\square$




Figure 5.4: The digraph of  $\gamma$  from Lemma 5.11.

**Lemma 5.12.** Let  $\aleph_{\varepsilon+1}$  be a successor cardinal. Then there is a collection  $\{\langle a_n^\mu \rangle\}_{\mu < \omega_{\varepsilon+1}}$  of increasing sequences  $\langle a_n^\mu \rangle$  of ordinals  $a_n^\mu < \omega_{\varepsilon+1}$  such that for all ordinals  $\mu, \lambda < \omega_{\varepsilon+1}$ ,  $a_0^\mu = 0$  and if  $\lambda < \mu$  then  $a_m^\lambda < a_n^\mu$  for all  $m, n \geq 1$ .

*Proof.* We construct such a collection by transfinite recursion. We define  $\langle a_n^0 \rangle = \langle 0, 1, 2, 3, \dots \rangle$ . Let  $\mu$  be an ordinal such that  $0 < \mu < \omega_{\varepsilon+1}$  and suppose  $\langle a_n^\lambda \rangle$  satisfying the hypotheses has been defined for every ordinal  $\lambda < \mu$ . Let  $A = \{a_n^\lambda : \lambda < \mu \text{ and } n \geq 0\}$  and  $\tau = \sup A$ . Then  $|A| = |\mu| \cdot \aleph_0 < \aleph_{\varepsilon+1}$ , and so  $\tau < \omega_{\varepsilon+1}$  by Lemma 5.8. Define  $\langle a_n^\mu \rangle = \langle 0, \tau + 1, \tau + 2, \tau + 3, \dots \rangle$  and note that  $\langle a_n^\mu \rangle$  is an increasing sequence of ordinals  $a_n^\mu < \omega_{\varepsilon+1}$  with  $a_0^\mu = 0$ . The construction has been completed. It is clear from the construction that  $a_m^\lambda < a_n^\mu$  for all  $\lambda, \mu < \omega_{\varepsilon+1}$  with  $\lambda < \mu$  and all  $m, n \geq 1$ .  $\square$

**Remark 5.13.** Let  $\{\langle a_n^\mu \rangle\}_{\mu < \omega_{\varepsilon+1}}$  be a collection from Lemma 5.12. Then it is clear that for all ordinals  $\lambda, \mu < \omega_{\varepsilon+1}$ , if  $\lambda < \mu$  then  $\langle a_n^\lambda \rangle$  does not dominate  $\langle a_n^\mu \rangle$ .

**Definition 5.14.** Let  $\aleph_{\varepsilon+1}$  be a successor cardinal. Denote by  $\text{IS}_{\omega_{\varepsilon+1}}$  the set of all increasing sequences  $\langle a_n \rangle$  of ordinals  $a_n < \omega_{\varepsilon+1}$  such that  $a_0 = 0$ . Define a relation  $\approx$  on  $\text{IS}_{\omega_{\varepsilon+1}}$  by

$$\langle a_n \rangle \approx \langle b_n \rangle \text{ if } \langle b_n \rangle \text{ dominates } \langle a_n \rangle \text{ and } \langle a_n \rangle \text{ dominates } \langle b_n \rangle.$$

It is straightforward to show that  $\approx$  is an equivalence relation on  $\text{IS}_{\omega_{\varepsilon+1}}$ . We denote by  $[\langle a_n \rangle]_\approx$  the equivalence class of  $\langle a_n \rangle$ , and by  $\text{IS}_{\omega_{\varepsilon+1}}^\approx$  the set of all equivalence classes of  $\approx$ .

**Lemma 5.15.** *For any successor cardinal  $\aleph_{\varepsilon+1}$ ,  $|\text{IS}_{\omega_{\varepsilon+1}}| = \aleph_{\varepsilon+1}^{\aleph_0}$  and  $\aleph_{\varepsilon+1} \leq |\text{IS}_{\omega_{\varepsilon+1}}^{\approx}| \leq \aleph_{\varepsilon+1}^{\aleph_0}$ .*

*Proof.* Denote by  $S_{\omega_{\varepsilon+1}}$  the set of all sequences  $\langle s_n \rangle$  of ordinals  $s_n < \omega_{\varepsilon+1}$ . Then  $S_{\omega_{\varepsilon+1}}$  is the set of all functions from  $\mathbb{N}$  to  $\omega_{\varepsilon+1}$ , and so  $|S_{\omega_{\varepsilon+1}}| = |\omega_{\varepsilon+1}|^{|\mathbb{N}|} = \aleph_{\varepsilon+1}^{\aleph_0}$ . Since  $\text{IS}_{\omega_{\varepsilon+1}}$  is a subset of  $S_{\omega_{\varepsilon+1}}$ , we have  $|\text{IS}_{\omega_{\varepsilon+1}}| \leq \aleph_{\varepsilon+1}^{\aleph_0}$ . Let  $S_{\omega_{\varepsilon+1}}^0$  be the subset of  $S_{\omega_{\varepsilon+1}}$  consisting of all sequences  $\langle s_n \rangle$  such that  $s_n > 0$  for all  $n \geq 0$ . Then  $|S_{\omega_{\varepsilon+1}}^0| = |S_{\omega_{\varepsilon+1}}| = \aleph_{\varepsilon+1}^{\aleph_0}$ . Define a function  $f : S_{\omega_{\varepsilon+1}}^0 \rightarrow \text{IS}_{\omega_{\varepsilon+1}}$  by  $\langle s_n \rangle f = \langle a_n \rangle$ , where

$$a_0 = 0 \text{ and } a_{n+1} = a_n + s_n \text{ for all } n \geq 0.$$

Then  $f$  is injective (since for all ordinals  $\mu, \lambda_1, \lambda_2$ , if  $\mu + \lambda_1 = \mu + \lambda_2$  then  $\lambda_1 = \lambda_2$  [20, page 120, Lemma 5.4]), and so  $|\text{IS}_{\omega_{\varepsilon+1}}| \geq |S_{\omega_{\varepsilon+1}}^0| = \aleph_{\varepsilon+1}^{\aleph_0}$ . We have proved that  $|\text{IS}_{\omega_{\varepsilon+1}}| = \aleph_{\varepsilon+1}^{\aleph_0}$ .

We have  $|\text{IS}_{\omega_{\varepsilon+1}}^{\approx}| \leq |\text{IS}_{\omega_{\varepsilon+1}}| = \aleph_{\varepsilon+1}^{\aleph_0}$ . Let  $\{\langle a_n^\mu \rangle\}_{\mu < \omega_{\varepsilon+1}}$  be a collection of sequences constructed as in Lemma 5.12. Then for all ordinals  $\lambda, \mu < \omega_{\varepsilon+1}$ ,  $\langle a_n^\mu \rangle \in \text{IS}_{\omega_{\varepsilon+1}}$  and if  $\lambda < \mu$  then  $\langle a_n^\lambda \rangle$  does not dominate  $\langle a_n^\mu \rangle$  (see Remark 5.13). It follows that any two different sequences from the collection  $\{\langle a_n^\mu \rangle\}_{\mu < \omega_{\varepsilon+1}}$  are in different equivalence classes of  $\approx$ . Since there are  $\aleph_{\varepsilon+1}$  sequences in the collection, it follows that  $|\text{IS}_{\omega_{\varepsilon+1}}^{\approx}| \geq \aleph_{\varepsilon+1}$ . This concludes the proof.  $\square$

We can now prove the counting theorem. For a set  $A$ , we denote by  $\mathcal{P}(A)$  the power set of  $A$ .

**Theorem 5.16.** *Let  $X$  be an infinite set with  $|X| = \aleph_\varepsilon$ . Then in  $P(X)$  there are:*

- (1)  $\max\{2^{\aleph_0}, \aleph_{\varepsilon+1}\}$  conjugacy classes containing a representative with a cycle, of which  $\aleph_0$  have a connected representative;
- (2)  $2^{\aleph_\varepsilon}$  conjugacy classes containing a representative with a connected component of type *rro*, but no cycles, of which at least  $\aleph_{\varepsilon+1}$  and at most  $\aleph_{\varepsilon+1}^{\aleph_0}$  have a connected representative.
- (3)  $\aleph_{\varepsilon+1}$  conjugacy classes containing a representative with a connected component of type *cho*, but no cycles or connected components of type *rro*, of which  $\aleph_{\varepsilon+1}$  have a connected representative.

*In total, there are  $2^{\aleph_\varepsilon}$  conjugacy classes in  $P(X)$ , of which at least  $\aleph_{\varepsilon+1}$  and at most  $\aleph_{\varepsilon+1}^{\aleph_0}$  have a connected representative.*

*Proof.* For  $\alpha \in P(X)$ , we define  $i_\alpha, j_\alpha \in \{0, 1\}$  by  $i_\alpha = 1$  if  $\rho_\gamma(x_0) = s(\alpha)$  for some connected component of  $\alpha$  of type *cho* with root  $x_0$  (and  $i_\alpha = 0$  otherwise); and  $j_\alpha \in \{0, 1\}$  by  $j_\alpha = 1$  if  $\alpha$  has a double ray (and  $j_\alpha = 0$  otherwise).

To prove (1), let  $A = \{[\alpha]_c : \alpha \in P(X) \text{ has a cycle}\}$ . Let

$$A' = \{[\alpha]_c \in A : \alpha \text{ has no maximal left rays}\} \text{ and } A'' = \{[\alpha]_c \in A : \alpha \text{ has a maximal left ray}\}.$$

By Theorem 5.3(4),  $\{A', A''\}$  is a partition of  $A$ . Define  $f' : A' \rightarrow \mathcal{P}(\mathbb{Z}_+) \times (\omega_{\varepsilon+1} + 1) \times \{0, 1\}$  by  $([\alpha]_c)f' = (\text{cs}(\alpha), s(\alpha), i_\alpha)$ . Then  $f'$  is well defined and injective by Theorem 5.3 and Lemma 5.9. (See Definition 5.5 and the discussion following the definition to see why  $i_\alpha$  is needed.) Similarly, the mapping  $f'' : A'' \rightarrow \mathcal{P}(\mathbb{Z}_+) \times (\omega_{\varepsilon+1} + 1) \times \{0, 1\}$  defined by  $([\alpha]_c)f'' = (\text{cs}(\alpha), s(\alpha), i_\alpha)$  is well defined and injective. Thus

$$|A'| \leq |\mathcal{P}(\mathbb{Z}_+)| \cdot |\omega_{\varepsilon+1} + 1| \cdot 2 = 2^{\aleph_0} \cdot \aleph_{\varepsilon+1} \cdot 2 = \max\{2^{\aleph_0}, \aleph_{\varepsilon+1}\},$$

and the same holds for  $|A''|$ . Hence

$$|A| = |A'| + |A''| \leq 2|A'| \leq 2 \max\{2^{\aleph_0}, \aleph_{\varepsilon+1}\} = \max\{2^{\aleph_0}, \aleph_{\varepsilon+1}\}.$$

Let  $P$  be the set of prime positive integers. For any nonempty subset  $Q \subseteq P$ , let  $\{\theta_q\}_{q \in Q}$  be a collection of completely disjoint cycles  $\theta_q$  such that  $\theta_q$  has length  $q$  for every  $q \in Q$ . (Such



a collection exists since  $X$  is infinite.) Define  $\beta_Q \in P(X)$  by  $\beta_Q = \bigsqcup_{q \in Q} \theta_q$ . For all nonempty subsets  $Q_1, Q_2 \subseteq P$  with  $Q_1 \neq Q_2$ , we have  $(\beta_{Q_1}, \beta_{Q_2}) \notin \sim_c$  by Theorem 5.3(1). It follows that  $|A| \geq \mathcal{P}(P) = 2^{\aleph_0}$ . By Lemma 5.10, for every nonzero ordinal  $\mu < \omega_{\varepsilon+1}$ , there is  $\gamma_\mu \in P(X)$  of type *cho* with root  $x_0$  such that  $\rho(x_0) = \mu$ . For all nonzero ordinals  $\lambda, \mu < \omega_{\varepsilon+1}$  with  $\lambda \neq \mu$ , we have  $(\gamma_\lambda, \gamma_\mu) \notin \sim_c$  by Theorem 5.3(5). It follows that  $|A| \geq |\omega_{\varepsilon+1}| = \aleph_{\varepsilon+1}$ . Hence  $|A| \geq \max\{2^{\aleph_0}, \aleph_{\varepsilon+1}\}$ , and so  $|A| = \max\{2^{\aleph_0}, \aleph_{\varepsilon+1}\}$ .

Let  $A_1 = \{[\gamma]_c : \gamma \in P(X) \text{ has a cycle and } \gamma \text{ is connected}\}$ . Fix a subset  $X_0 = \{x_0, x_1, \dots\}$  of  $X$ , and for every integer  $n \geq 0$ , define a cycle  $\gamma_n = (x_0 x_1 \dots x_{n-1}) \in P(X)$ . Then, by Proposition 4.8 and Theorem 5.3,  $A_1 = \{[\gamma_0]_c, [\gamma_1]_c, [\gamma_2]_c, \dots\}$ , and so  $|A_1| = \aleph_0$ . We have proved (1).

To prove (2), let

$$B = \{[\alpha]_c : \alpha \in P(X) \text{ has a connected component of type } rro, \text{ but no cycles}\},$$

and let  $B_1$  be the subset of  $B$  consisting of all conjugacy classes  $[\gamma]_c \in B$  such that  $\gamma$  is connected. Fix a double ray  $\omega = \langle \dots x_{-1} x_0 x_1 \dots \rangle \in P(X)$  and note that

$$B_1 = \{[\gamma]_c : \gamma \in P(X) \text{ is of type } rro\} \cup \{[\omega]_c\}.$$

Let  $B'_1 = \{[\gamma]_c : \gamma \in P(X) \text{ is of type } rro\}$ . For every  $\gamma \in P(X)$  of type *rro*, we fix a maximal right ray  $\eta^\gamma$  in  $\gamma$ . Define a function  $g : B'_1 \rightarrow \text{IS}_{\omega_{\varepsilon+1}}^\approx$  by  $([\gamma]_c)g = [\langle \eta_n^\gamma \rangle]_\approx$ . Note that  $\langle \eta_n^\gamma \rangle \in \text{IS}_{\omega_{\varepsilon+1}}$  by Lemma 5.9. Suppose  $[\gamma_1]_c, [\gamma_2]_c \in B'_1$  with  $[\gamma_1]_c = [\gamma_2]_c$ . Then, by Theorem 5.3(3) and Lemma 4.30, the sequences  $\langle \eta_n^{\gamma_1} \rangle$  and  $\langle \eta_n^{\gamma_2} \rangle$  dominate each other, and so  $[\langle \eta_n^{\gamma_1} \rangle]_\approx = [\langle \eta_n^{\gamma_2} \rangle]_\approx$ . We have proved that  $g$  is well defined. The function  $g$  is also injective (by Theorem 5.3(3)) and surjective (by Lemma 5.11). Thus  $|B'_1| = |\text{IS}_{\omega_{\varepsilon+1}}^\approx|$ , and so, by Lemma 5.15,  $\aleph_{\varepsilon+1} \leq |B'_1| \leq \aleph_{\varepsilon+1}^{\aleph_0}$ .

Then  $\aleph_{\varepsilon+1} \leq |B_1| \leq \aleph_{\varepsilon+1}^{\aleph_0}$  since  $|B_1| = |B'_1| + 1$ .

As to the cardinality of  $B$ , clearly  $|B| \leq |P(X)| = (\aleph_\varepsilon + 1)^{\aleph_\varepsilon} = 2^{\aleph_\varepsilon}$ . Let

$$B' = \{[\alpha]_c \in B : \alpha \text{ has no maximal left rays or double rays}\},$$

$$B'' = \{[\alpha]_c \in B : \alpha \text{ has a maximal left ray but no double rays}\}.$$

By Theorem 5.3(3)(4),  $\{B', B'', \{[\omega]_c\}\}$  is a partition of  $B$ .

We will now prove that  $|B'| \geq 2^{\aleph_\varepsilon}$ . Since  $|B'_1| \geq \aleph_{\varepsilon+1}$ , there is a collection  $\{\gamma_\mu\}_{\mu < \omega_{\varepsilon+1}}$  of transformations  $\gamma_\mu \in P(X)$  of type *rro* such that  $(\gamma_\mu, \gamma_\lambda) \notin \sim_c$  if  $\mu \neq \lambda$ . Since  $|\omega_\varepsilon| = \aleph_\varepsilon$  and  $\aleph_\varepsilon \cdot \aleph_\varepsilon = \aleph_\varepsilon$ , there is a partition  $\{X_\mu\}_{\mu < \omega_\varepsilon}$  of  $X$  such that  $|X_\mu| = |X| = \aleph_\varepsilon$  for every  $\mu < \omega_\varepsilon$ . Let  $\mu < \omega_\varepsilon$ . Since  $|X_\mu| = |X|$ , there is a bijection  $h_\mu : X_\mu \rightarrow X$ . We can use  $h_\mu$  to obtain a “copy” of  $\gamma_\mu$  in  $P(X_\mu)$ : define  $\gamma'_\mu \in P(X_\mu)$  by

$$x\gamma'_\mu = y \Leftrightarrow (xh_\mu)\gamma_\mu = yh_\mu \text{ (for all } x, y \in X_\mu\text{)}.$$

Let  $\mu, \lambda < \omega_\varepsilon$  with  $\mu \neq \lambda$ . Then  $(\gamma_\mu, \gamma_\lambda) \notin \sim_c$ , and so, by Theorem 5.3(3) and Lemma 4.30,  $(\langle \eta_n \rangle, \langle \xi_n \rangle) \notin \approx$  for every maximal right ray  $\eta$  in  $\gamma_\mu$  and every maximal right ray  $\xi$  in  $\gamma_\lambda$ . It follows that

$$(\langle \eta'_n \rangle, \langle \xi'_n \rangle) \notin \approx \tag{5.1}$$

for every maximal right ray  $\eta'$  in  $\gamma'_\mu$  and every maximal right ray  $\xi'$  in  $\gamma'_\lambda$ . Let  $K$  be a nonempty subset of  $\omega_\varepsilon$ . Select  $\nu = \nu_K \in K$  and a maximal right ray  $[x_0 x_1 x_2 \dots]$  in  $\gamma'_\nu$ . Define  $\alpha_K \in P(X)$  by  $\alpha_K = \bigsqcup_{\mu \in K} \gamma'_\mu$ , and note that  $\alpha_K$  does not have a cycle or a double ray. Let  $K, L$  be nonempty subsets of  $\omega_\varepsilon$  such that  $K \neq L$ . We may assume that there is  $\mu \in K$  such that  $\mu \notin L$ . Consider  $\gamma'_\mu$ , which is a connected component of  $\alpha_K$ . Let  $\gamma'_\lambda$  be any connected component of  $\alpha_L$ . Then, by (5.1),  $(\langle \eta'_n \rangle, \langle \xi'_n \rangle) \notin \approx$  for every maximal right ray  $\eta'$  in  $\gamma'_\mu$  and every maximal right ray  $\xi'$  in  $\gamma'_\lambda$ . (Note that, by the definition of  $\alpha_K$ , this is also true when  $\mu = \nu_K$  or  $\lambda = \nu_L$ .) Thus  $(\alpha_K, \alpha_L) \notin \sim_c$  by Theorem 6.1(3). Hence any two different transformations from the collection  $\{\alpha_K\}_{\emptyset \neq K \subseteq \omega_\varepsilon}$  are in different equivalence classes of  $\sim_c$ . Since there are  $2^{\aleph_\varepsilon}$  transformations in the collection,

it follows that  $|B'| \geq 2^{\aleph_\varepsilon}$ . Hence  $|B| = |B'| + |B''| + |\{[\omega]_c\}| \geq |B'| \geq 2^{\aleph_\varepsilon}$ , and so  $|B| = 2^{\aleph_\varepsilon}$ . We have proved (2).

To prove (3), let  $C$  be the set of all  $[\alpha]_c$  such that  $\alpha \in P(X)$  has a connected component of type *cho*, but no cycles or connected components of type *rro*. Let  $C' = \{[\alpha]_c \in C : \alpha \text{ has no maximal left rays}\}$  and  $C'' = \{[\alpha]_c \in C : \alpha \text{ has a maximal left ray}\}$ . By Theorem 5.3(4),  $\{C', C''\}$  is a partition of  $C$ . Fix a maximal left ray  $\lambda = \langle \dots x_2 x_1 x_0 \rangle \in P(X)$  and note that  $C'' = \{[\lambda]_c\}$ . Define  $h : C' \rightarrow (\omega_{\varepsilon+1} + 1) \times \{0, 1\} \times \{0, 1\}$  by  $([\alpha]_c)h = (s(\alpha), i_\alpha, j_\alpha)$ . Then  $h$  is well defined and injective by Theorem 5.3 and Lemma 5.9, and so  $|C'| \leq \aleph_{\varepsilon+1} \cdot 2 \cdot 2 = \aleph_{\varepsilon+1}$ . Thus  $|C| = |C'| + |C''| = |C'| + 1 \leq \aleph_{\varepsilon+1} + 1 = \aleph_{\varepsilon+1}$ .

Let  $C_1$  be the subset of  $C$  consisting of all  $[\gamma]_c \in C$  such that  $\gamma$  is connected. Note that  $C_1 = \{[\gamma]_c : \gamma \in P(X) \text{ is of type } cho\} \cup \{[\lambda]_c\}$ . As in the proof of (1), we can construct a collection  $\{\gamma_\mu\}_{0 < \mu < \omega_{\varepsilon+1}}$  of connected elements of  $P(X)$  of type *cho* such that  $(\gamma_\lambda, \gamma_\mu) \notin \sim_c$  if  $\lambda \neq \mu$ . Thus  $|C_1| \geq \aleph_{\varepsilon+1}$ , and so  $\aleph_{\varepsilon+1} \leq |C_1| \leq |C| \leq \aleph_{\varepsilon+1}$ . Hence  $|C| = |C_1| = \aleph_{\varepsilon+1}$ , which concludes the proof of (3).

The conjugacy classes considered in (1)–(3) cover all conjugacy classes in  $P(X)$ . Thus, there are at most  $\max\{2^{\aleph_0}, \aleph_{\varepsilon+1}\} + 2^{\aleph_\varepsilon} + \aleph_{\varepsilon+1} = 2^{\aleph_\varepsilon}$  conjugacy classes in  $P(X)$  (which also follows from the fact that  $|P(X)| = 2^{\aleph_\varepsilon}$ ). By (2), there are at least  $2^{\aleph_\varepsilon}$  conjugacy classes, so the number of conjugacy classes in  $P(X)$  is  $2^{\aleph_\varepsilon}$ . By (1)–(3), at least  $\aleph_{\varepsilon+1}$  and at most  $\aleph_0 + \aleph_{\varepsilon+1} + \aleph_{\varepsilon+1} = \aleph_{\varepsilon+1}^{\aleph_0}$  of these conjugacy classes have a connected representative. (We point out that if a conjugacy class has a connected representative, it does not imply that all representatives of this class are connected.)  $\square$

## 6 Conjugacy in $T(X)$

A characterization of the conjugacy  $\sim_c$  in the monoid  $T(X)$  of full transformations on  $X$  is simpler than that of the conjugacy in  $P(X)$  (see Section 5). The reason is that a connected component of  $\alpha \in T(X)$  cannot have a maximal left ray or a maximal chain. Suppose  $\alpha, \beta \in T(X)$  and  $\alpha \sim_c \beta$  in  $P(X)$ . Then  $\alpha\phi = \phi\beta$  and  $\beta\psi = \psi\alpha$  for some rp-homomorphisms  $\phi$  and  $\psi$ . By Lemma 3.5,  $X = \text{span}(\alpha) \subseteq \text{dom}(\phi)$  and  $X = \text{span}(\beta) \subseteq \text{dom}(\psi)$ . Therefore,  $\phi, \psi \in T(X)$ , and so  $\alpha \sim_c \beta$  in  $T(X)$ . In other words,  $\sim_c$  in  $T(X)$  is the restriction of  $\sim_c$  in  $P(X)$  to  $T(X) \times T(X)$ .

These observations and Theorem 5.3 give a characterization of  $\sim_c$  in  $T(X)$

**Theorem 6.1.** *Let  $\alpha, \beta \in T(X)$ . Then  $\alpha \sim_c \beta$  in  $T(X)$  if and only if exactly one of the following conditions is satisfied:*

- (1) *both  $\alpha$  and  $\beta$  have a cycle and  $\text{cs}(\alpha) = \text{cs}(\beta)$ ;*
- (2) *both  $\alpha$  and  $\beta$  have a double ray but no cycles;*
- (3) *all connected components of both  $\alpha$  and  $\beta$  have type *rro* and:*
  - (a) *for every connected component  $\gamma$  of  $\alpha$ , there is a connected component  $\delta$  of  $\beta$  such that  $\langle \xi_n^\delta \rangle$  dominates  $\langle \eta_n^\gamma \rangle$  for some maximal right ray  $\eta$  in  $\gamma$  and some maximal right ray  $\xi$  in  $\delta$ , and*
  - (b) *for every connected component  $\delta$  of  $\beta$ , there is a connected component  $\gamma$  of  $\alpha$  such that  $\langle \eta_n^\gamma \rangle$  dominates  $\langle \xi_n^\delta \rangle$  for some maximal right ray  $\xi$  in  $\delta$  and some maximal right ray  $\eta$  in  $\gamma$ .*

**Example 6.2.** Let  $X = \{x_0, x_1, x_2, \dots, y_1, y_2, y_3, \dots\}$  and consider

$$\begin{aligned} \alpha &= [x_0 y_0 x_1 y_1 x_2 y_2 \dots], \\ \beta &= [x_0 y_0 x_1 y_1 x_2 y_2 \dots] \sqcup [y_1 y_2 x_1] \sqcup [y_3 y_4 y_5 y_6 x_2] \sqcup [y_7 y_8 y_9 y_{10} y_{11} y_{12} x_3] \sqcup \dots \end{aligned}$$

in  $T(X)$  (see Figure 6.1). We will argue that  $\alpha$  and  $\beta$  are not conjugate. Both  $\alpha$  and  $\beta$  are connected of type  $rr\alpha$ . The only maximal right ray in  $\alpha$  is  $\eta = [x_0 y_0 x_1 y_1 x_2 y_2 \dots]$  with  $\langle \eta_n^\gamma \rangle = \langle n \rangle$  (where  $\gamma = \alpha$ ). If  $\alpha$  and  $\beta$  were conjugate, then  $\langle \eta_n \rangle$  would dominate  $\langle \xi_n^\delta \rangle$  (where  $\delta = \beta$ ) for some maximal right ray  $\xi$  in  $\beta$ , and so for all maximal right rays  $\xi$  in  $\beta$  (see Lemma 4.30 below). The right chain  $\xi = [x_0 y_0 x_1 y_1 x_2 y_2 \dots]$  is a maximal right chain in  $\beta$  with  $\langle \xi_n^\delta \rangle = \langle 2n \rangle$ . It is clear that the sequence  $\langle n \rangle$  does not dominate the sequence  $\langle 2n \rangle$ . Hence, by Theorem 6.1,  $\alpha$  and  $\beta$  are not conjugate.




Figure 6.1: The digraphs of  $\alpha$  and  $\beta$  from Example 6.2.

If  $X$  is a finite set, then every  $\alpha \in T(X)$  has a cycle. Hence, Theorem 6.1 gives us the following corollary.

**Corollary 6.3.** *Let  $X$  be finite, and let  $\alpha, \beta \in T(X)$ . Then  $\alpha \sim_c \beta$  if and only if  $\text{cs}(\alpha) = \text{cs}(\beta)$ .*

Modifying the proof of Theorem 5.16, we can count the number of conjugacy classes in an infinite  $T(X)$ .

**Theorem 6.4.** *Let  $X$  be an infinite set with  $|X| = \aleph_\varepsilon$ . Then in  $T(X)$  there are:*

- (1)  $2^{\aleph_0}$  conjugacy classes consisting of transformations with a cycle, of which  $\aleph_0$  have a connected representative;
- (2) one conjugacy class consisting of transformations with a double ray but not a cycle;
- (3)  $2^{\aleph_\varepsilon}$  conjugacy classes consisting of transformations without a cycle or a double ray, of which at least  $\aleph_{\varepsilon+1}$  and at most  $\aleph_{\varepsilon+1}^{\aleph_0}$  have a connected representative.

*In total, there are  $2^{\aleph_\varepsilon}$  conjugacy classes in  $T(X)$ , of which at least  $\aleph_{\varepsilon+1}$  and at most  $\aleph_{\varepsilon+1}^{\aleph_0}$  have a connected representative.*

The reason for (1) is that  $\alpha \in T(X)$  does not have any maximal left rays or components of type  $cho$ . Thus, the set  $A = A' \cup A''$  from the proof of (1) of Theorem 5.16 reduces to  $A'$ , and the function  $f' : A' \rightarrow \mathcal{P}(\mathbb{Z}_+) \times (\omega_{\varepsilon+1} + 1) \times \{0, 1\}$  reduces to  $f' : A' \rightarrow \mathcal{P}(\mathbb{Z}_+)$ . The reason for (2) is that if  $\alpha \in T(X)$  has a double ray but not a cycle, then each component of  $\alpha$  either has a double ray or is of type  $rr\alpha$ . Any two such transformations are then conjugate by Lemma 4.15 and Proposition 4.18.

## 7 Conjugacy in $\Gamma(X)$

By  $\Gamma(X)$  we mean the subsemigroup of  $T(X)$  consisting of injective transformations. If  $X$  is finite, then  $\Gamma(X) = \text{Sym}(X)$  but this is not the case for an infinite  $X$ . The semigroup  $\Gamma(X)$  is universal for right cancellative semigroups with no idempotents (except possibly the identity): that is, any such semigroup can be embedded in  $\Gamma(X)$  for some  $X$  [11, Lemma 1.0]. The semigroup  $\Gamma(X)$  has been studied mainly in the context of: ideals and congruences [31, 37];  $\mathcal{G}(X)$ -normal semigroups [29, 30, 35]; Baer-Levi semigroups [32, 33];  $BQ$ -semigroups [23, 36], and centralizers [24, 25]. In this section, we characterize the conjugacy  $\sim_c$  in  $\Gamma(X)$  for an arbitrary set  $X$ .

We note that every connected transformation in  $P(X)$  that is also injective is a cycle, a ray, or a chain. Since transformations in  $\Gamma(X)$  are full,  $\alpha \in \Gamma(X)$  cannot contain a maximal left ray or a maximal chain. These observations give the following lemma.

**Lemma 7.1.** *Let  $\alpha \in \Gamma(X)$ . Then every connected component of  $\alpha$  is a right ray, a double ray, or a cycle.*

The following proposition follows from Lemma 7.1 and Proposition 4.5.

**Proposition 7.2.** *Let  $\alpha \in \Gamma(X)$ . Then there exist unique sets:  $A$  of right rays,  $B$  of double rays, and  $C$  of cycles such that the transformations in  $A \cup B \cup C$  are pairwise completely disjoint and*

$$\alpha = \left( \bigsqcup_{\eta \in A} \eta \right) \sqcup \left( \bigsqcup_{\omega \in B} \omega \right) \sqcup \left( \bigsqcup_{\theta \in C} \theta \right).$$

Let  $\alpha \in \Gamma(X)$ . We will denote the unique sets  $A$ ,  $B$ , and  $C$  from Proposition 7.2 by  $A_\alpha$ ,  $B_\alpha$ , and  $C_\alpha$ , respectively. For  $n \geq 1$ , we will denote by  $C_\alpha^n$  the subset of  $C_\alpha$  consisting of cycles of length  $n$ . Note that:

- $A_\alpha$  = the set of maximal right rays contained in  $\alpha$ ,
- $B_\alpha$  = the set of double rays contained in  $\alpha$ ,
- $C_\alpha$  = the set of cycles contained in  $\alpha$ .

For  $\eta = [x_0 x_1 x_2 \dots]$ ,  $\omega = \langle \dots x_{-1} x_0 x_1 \dots \rangle$ ,  $\theta = (x_0 x_1 \dots x_{k-1})$ , and any  $\phi$  in  $\Gamma(X)$ , we define:

$$\eta\phi^* = [x_0\phi \ x_1\phi \ x_2\phi \ \dots], \ \omega\phi^* = \langle \dots x_{-1}\phi \ x_0\phi \ x_1\phi \ \dots \rangle, \ \theta\phi^* = (x_0\phi \ x_1\phi \ \dots x_{k-1}\phi).$$

**Proposition 7.3.** *Let  $\alpha, \beta, \phi \in \Gamma(X)$ . Then  $\phi$  is a homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$  if and only if for all  $\eta \in A_\alpha$ ,  $\omega \in B_\alpha$ , and  $\theta \in C_\alpha$ :*

- (1) *either there is a unique  $\eta_1 \in A_\beta$  such that  $\eta\phi^* \sqsubset \eta_1$  or there is a unique  $\omega_1 \in B_\beta$  such that  $\eta\phi^* \sqsubset \omega_1$ ;*
- (2)  *$\omega\phi^* \in B_\beta$  and  $\theta\phi^* \in C_\beta$ .*

*Proof.* Suppose  $\phi$  is a homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . Let  $\eta = [x_0 x_1 x_2 \dots] \in A_\alpha$ . Then, since  $\phi$  is an injective homomorphism,  $\eta\phi^* = [x_0\phi \ x_1\phi \ x_2\phi \ \dots]$  is a right ray in  $\Gamma(\beta)$ . By the proof of Proposition 4.18,  $\phi|_{\text{span}(\eta)}$  is a homomorphism from  $\Gamma(\eta)$  to  $\Gamma(\gamma)$  for some connected component  $\gamma$  of  $\beta$ . By Lemma 7.1, either  $\gamma = \eta_1 = [y_0 y_1 y_2 \dots]$  is a right ray in  $\beta$  or  $\gamma = \omega_1 = \langle \dots y_{-1} y_0 y_1 \dots \rangle$  is a double ray in  $\beta$  ( $\gamma$  cannot be a cycle since  $\phi$  is injective). In the former case,  $\eta\phi^* \sqsubset \eta_1$ , and in the latter case,  $\eta\phi^* \sqsubset \omega_1$ . The uniqueness of  $\eta_1$  and  $\omega_1$  follows from the fact that the elements of  $A_\beta \cup B_\beta$  are pairwise completely disjoint. We have proved (1). The proof of (2) is similar.

Conversely, suppose that  $\phi$  satisfies (1) and (2). Then it follows immediately that for all  $x, y \in X$ ,  $x \xrightarrow{\alpha} y$  implies  $x\phi \xrightarrow{\beta} y\phi$ , and so  $\phi$  is a homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ .  $\square$

**Definition 7.4.** Let  $\alpha, \beta \in \Gamma(X)$ . For a homomorphism  $\phi \in \Gamma(X)$  from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ , we define a mapping  $h_\phi : A_\alpha \cup B_\alpha \cup C_\alpha \rightarrow A_\beta \cup B_\beta \cup C_\beta$  by:

$$\delta h_\phi = \begin{cases} \eta & \text{if } \delta \in A_\alpha \text{ and } \delta\phi^* \sqsubset \eta \text{ for some } \eta \in A_\beta, \\ \omega & \text{if } \delta \in A_\alpha \text{ and } \delta\phi^* \sqsubset \omega \text{ for some } \omega \in B_\beta, \\ \delta\phi^* & \text{if } \delta \in B_\alpha \cup C_\alpha. \end{cases}$$

Note that  $h_\phi$  is well defined (by Proposition 7.3) and injective (since  $\phi$  is injective).

We will need the following lemma from set theory (whose proof is straightforward).

**Lemma 7.5.** Let  $A_1, B_1, A_2$ , and  $B_2$  be sets such that  $A_1 \cap B_1 = \emptyset$ ,  $A_2 \cap B_2 = \emptyset$ ,  $|A_1| + |B_1| \leq |A_2| + |B_2|$ , and  $|B_1| \leq |B_2|$ . Then there is an injective mapping  $f : A_1 \cup B_1 \rightarrow A_2 \cup B_2$  such that  $xf \in B_2$  for every  $x \in B_1$ .

We can now characterize the conjugacy  $\sim_c$  in  $\Gamma(X)$ .

**Theorem 7.6.** Let  $\alpha, \beta \in \Gamma(X)$ . Then  $\alpha \sim_c \beta$  in  $\Gamma(X)$  if and only if  $|A_\alpha| + |B_\alpha| = |A_\beta| + |B_\beta|$ ,  $|B_\alpha| = |B_\beta|$ , and  $|C_\alpha^n| = |C_\beta^n|$  for every  $n \geq 1$ .

*Proof.* Suppose  $\alpha \sim_c \beta$  in  $\Gamma(X)$ . Then, by Corollary 3.9, there is  $\phi \in \Gamma(X)$  such that  $\phi$  is a homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . Define  $f : A_\alpha \cup B_\alpha \rightarrow A_\beta \cup B_\beta$  by  $\delta f = \delta h_\phi$ . (By the definitions of  $h_\phi$  and  $\phi^*$ ,  $\delta f$  is indeed in  $A_\beta \cup B_\beta$  if  $\delta \in A_\alpha \cup B_\alpha$ .) The mapping  $f$  is injective (since  $h_\phi$  is injective),  $A_\alpha \cap B_\alpha = \emptyset$ , and  $A_\beta \cap B_\beta = \emptyset$ . Thus

$$|A_\alpha| + |B_\alpha| = |A_\alpha \cup B_\alpha| \leq |A_\beta \cup B_\beta| = |A_\beta| + |B_\beta|.$$

Similarly,  $|B_\alpha| \leq |B_\beta|$  since  $g : B_\alpha \rightarrow B_\beta$  defined by  $\omega g = \omega h_\phi$  is well defined and injective. Let  $n \geq 1$ . Define  $h : C_\alpha^n \rightarrow C_\beta^n$  by  $\theta h = \theta h_\phi$ . (If  $\theta = (x_0 \dots x_{n-1}) \in C_\alpha^n$ , then  $\theta h_\phi = \theta\phi^* = (x_0\phi^* \dots x_{n-1}\phi^*) \in C_\beta^n$ .) The mapping  $h$  is injective, and so  $|C_\alpha^n| \leq |C_\beta^n|$ . By symmetry,  $|A_\beta| + |B_\beta| \leq |A_\alpha| + |B_\alpha|$ ,  $|B_\beta| \leq |B_\alpha|$ , and  $|C_\beta^n| \leq |C_\alpha^n|$ . Hence the stated equalities hold.

Conversely, suppose  $|A_\alpha| + |B_\alpha| = |A_\beta| + |B_\beta|$ ,  $|B_\alpha| = |B_\beta|$ , and  $|C_\alpha^n| = |C_\beta^n|$  for every  $n \geq 1$ . We will define an injective homomorphism  $\phi$  from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . By Lemma 7.5, there is an injective mapping  $f : A_\alpha \cup B_\alpha \rightarrow A_\beta \cup B_\beta$  such that  $\omega f \in B_\beta$  for every  $\omega \in B_\alpha$ . For every  $n \geq 1$ , fix a bijection  $g_n : C_\alpha^n \rightarrow C_\beta^n$ . Let  $n \geq 1$ . For all  $\eta \in A_\alpha$ ,  $\omega \in B_\alpha$ , and  $\theta \in C_\alpha^n$ , we define  $\phi$  on  $\text{dom}(\eta) \cup \text{dom}(\omega) \cup \text{dom}(\theta)$  in such a way that  $\eta\phi^* \sqsubset \eta f$ ,  $\omega\phi^* = \omega f$ , and  $\theta\phi^* = \theta g_n$ . Note that this defines  $\phi$  for every  $x \in X$ . By the definition of  $\phi$  and Proposition 7.3,  $\phi \in \Gamma(X)$  and  $\phi$  is a homomorphism from  $\Gamma(\alpha)$  to  $\Gamma(\beta)$ . By symmetry, there is an injective homomorphism  $\psi$  from  $\Gamma(\beta)$  to  $\Gamma(\alpha)$ . Hence  $\alpha \sim_c \beta$  by Corollary 3.9.  $\square$

**Example 7.7.** Let  $\alpha$  and  $\beta$  be partial transformations on an infinite set whose digraphs are presented in Figures 7.1 and 7.2, respectively. Then  $|A_\alpha| = 1$ ,  $|B_\alpha| = \aleph_0$ ,  $|A_\alpha| + |B_\alpha| = \aleph_0$ , and  $|C_\alpha^n| = 0$  for every  $n \geq 1$ . Also,  $|A_\beta| = 2$ ,  $|B_\beta| = \aleph_0$ ,  $|A_\beta| + |B_\beta| = \aleph_0$ , and  $|C_\beta^n| = 0$  for every  $n \geq 1$ . Thus  $\alpha \sim_c \beta$  by Theorem 7.6.

Using Theorem 7.6, we can count the conjugacy classes in  $\Gamma(X)$ . First, we need the following lemma.

**Lemma 7.8.** Let  $X$  be an infinite set with  $|X| = \aleph_\varepsilon$ , let  $\alpha \in \Gamma(X)$ . Then  $|A_\alpha| \leq \aleph_\varepsilon$ ,  $|B_\alpha| \leq \aleph_\varepsilon$ , and  $|C_\alpha^n| \leq \aleph_\varepsilon$  for every  $n \geq 1$ .

*Proof.* Let  $Y = \bigcup_{\eta \in A_\alpha} \text{dom}(\eta) \subseteq X$ . Since the elements of  $A_\alpha$  are pairwise completely disjoint and  $|\text{dom}(\eta)| = \aleph_0$  for every  $\eta \in A_\alpha$ , we have

$$\aleph_\varepsilon = |X| \geq |Y| = \left| \bigcup_{\eta \in A_\alpha} \text{dom}(\eta) \right| = |A_\alpha| \cdot \aleph_0 \geq |A_\alpha|.$$

Thus  $|A_\alpha| \leq \aleph_\varepsilon$ . The proofs for  $B_\alpha$  and  $C_\alpha^n$  ( $n \geq 1$ ) are similar.  $\square$




Figure 7.1: The digraph of  $\alpha$  from Example 7.7.




Figure 7.2: The digraph of  $\beta$  from Example 7.7.

For sets  $A$  and  $B$ , we denote by  $A^B$  the set of all functions from  $B$  to  $A$ .

**Theorem 7.9.** *Let  $X$  be an infinite set with  $|X| = \aleph_\varepsilon$ . Let  $\kappa = \aleph_0 + |\varepsilon|$ . Then there are  $\kappa^{\aleph_0}$  conjugacy classes in  $\Gamma(X)$ , of which two have a connected representative if  $\aleph_\varepsilon = \aleph_0$ , and none has a connected representative if  $\aleph_\varepsilon > \aleph_0$ .*

*Proof.* Let  $K$  be the set of all cardinals  $\tau$  such that  $\tau \leq \aleph_\varepsilon$ . Then  $K$  contains  $\aleph_0$  finite cardinals and  $|\varepsilon| + 1$  infinite cardinals, hence  $|K| = \aleph_0 + |\varepsilon| + 1 = \aleph_0 + |\varepsilon| = \kappa$ . Let  $\Gamma(X)/\sim_c$  be the set of conjugacy classes of  $\Gamma(X)$ . Define a function  $f : \Gamma(X)/\sim_c \rightarrow K^{\mathbb{N}}$ , where  $\mathbb{N} = \{0, 1, 2, \dots\}$ , by

$$([\alpha]_c)f = (|A_\alpha| + |B_\alpha|, |B_\alpha|, |C_\alpha^1|, |C_\alpha^2|, |C_\alpha^3|, \dots).$$

By Theorem 7.6,  $f$  is well defined and injective. Thus  $|\Gamma(X)/\sim_c| \leq |K^{\mathbb{N}}| = |K|^{|\mathbb{N}|} = \kappa^{\aleph_0}$ .

We next define an injective mapping  $g : K^{\mathbb{N}} \rightarrow \Gamma(X)/\sim_c$ . Let

$$\xi = (\tau_2, \tau_3, \tau_4, \dots) \in K^{\mathbb{N}}.$$

(It will be clear from the definition of  $g$  why we begin the indexing with  $n = 2$ .) Let  $\tau = \sum_{n=2}^{\infty} n\tau_n$  (see [20, Chapter 9]). For every  $n \geq 2$ ,  $n\tau_n \leq \aleph_\varepsilon$  (since  $\tau_n \leq \aleph_\varepsilon$  and  $\aleph_\varepsilon$  is infinite). Thus

$$\tau = \sum_{n=2}^{\infty} n\tau_n \leq \aleph_0 \cdot \aleph_\varepsilon = \aleph_\varepsilon,$$

and so  $\aleph_\varepsilon + \tau = \aleph_\varepsilon$ . Hence, there is a collection  $\{X_n\}_{n \geq 1}$  of pairwise disjoint subsets of  $X$  such that  $\bigcup_{n=1}^{\infty} X_n = X$ ,  $|X_1| = \aleph_\varepsilon$ , and  $|X_n| = n\tau_n$  for every  $n \geq 2$ . Let  $n \geq 2$ . Since  $|X_n| = n\tau_n$ ,

there is a collection  $C_n$  of  $n$ -cycles in  $\Gamma(X)$  such that  $|C_n| = \tau_n$  and  $\text{dom}(\bigsqcup_{\theta \in C_n} \theta) = X_n$ . Let  $\alpha_n = \bigsqcup_{\theta \in C_n} \theta$ . Define a transformation  $\alpha_\xi$  on  $X$  by

$$\alpha_\xi = \bigsqcup_{n \geq 2} \alpha_n \sqcup \bigsqcup_{x \in X_1} (x).$$

Then  $\alpha \in \Gamma(X)$ ,  $A_\alpha = B_\alpha = \emptyset$ , and  $C_\alpha^n = C_n$  for all  $n \geq 2$ . Thus

$$(|C_\alpha^1|, |C_\alpha^2|, |C_\alpha^3|, |C_\alpha^4|, \dots) = (\aleph_\varepsilon, \tau_2, \tau_3, \tau_4, \dots),$$

and it follows from Theorem 7.6 that the mapping  $g : K^\mathbb{N} \rightarrow \Gamma(X)/\sim_c$  defined by  $\xi g = \alpha_\xi$  is injective. Hence  $|\Gamma(X)/\sim_c| \geq |K^\mathbb{N}| = |K|^{|\mathbb{N}|} = \kappa^{\aleph_0}$ .

Suppose  $|X| = \aleph_0$ , say  $X = \{x_1, x_2, x_3, \dots\}$ . Then, by Theorem 7.6 and Lemma 7.1, the only conjugacy classes in  $\Gamma(X)$  with a connected representative are  $[(x_1 x_2 x_3 \dots)]$  and  $[\langle \dots x_6 x_4 x_2 x_1 x_3 x_5 \dots \rangle]$ . (There is no single cycle in  $\theta$  in  $\Gamma(X)$  since  $\text{dom}(\theta)$  is finite.)

If  $|X| > \aleph_0$ , then no element  $\alpha \in \Gamma(X)$  is connected since  $\text{dom}(\alpha) = X$  and the domain of any right ray, double ray, or cycle has cardinality at most  $\aleph_0$ . The result follows.  $\square$

## 8 Problems

The results of this paper prompt a number of problems in combinatorics, semigroups, matrix theory, and set theory. The first problem asks for the number of conjugacy classes in some important finite semigroups.

**Problem 8.1.** Let  $X$  be a finite set. Is it possible to find a closed formula that gives the number of conjugacy classes in  $T(X)$ ,  $P(X)$  or  $\mathcal{I}(X)$  (where  $\mathcal{I}(X)$  denotes the symmetric inverse semigroup on  $X$ )?

The second problem might attract the attention of experts in set theory.

**Problem 8.2.** Let  $X$  be an infinite set with  $|X| = \aleph_\varepsilon$ . According to Theorem 6.4, the number of conjugacy classes in  $T(X)$  that have a connected representative is in the interval  $[\aleph_{\varepsilon+1}, \aleph_{\varepsilon+1}^{\aleph_0}]$ . Is it possible to be more precise and reduce the length of this interval?

In this paper we characterized the conjugate elements in some well-known transformation semigroups, but there are many other transformation semigroups, or endomorphism monoids of some relational algebras that may be considered.

**Problem 8.3.** Characterize  $\sim_c$ , and calculate the number of conjugacy classes, in other transformation semigroups such as, for example, those appearing in the problem list of [7, Section 6] or those appearing in the large list of transformation semigroups included in [12]. Especially interesting would be a characterization of the conjugacy classes in the centralizers of idempotents [5, 6].

The theorems and problems in this paper have natural linear counter-parts.

**Problem 8.4.** Characterize  $\sim_c$  in the endomorphism monoid of a (finite or infinite dimensional) vector space.

Whenever some result holds for both sets and vector spaces the natural step forward is to prove those results for independence algebras.

**Problem 8.5.** Characterize  $\sim_c$  in the endomorphism monoid of a (finite or infinite dimensional) independence algebra. (For historical notes on the importance of these algebras, see [3, 4]; for definitions and basic results, see [1, 2, 8, 9, 10, 13, 14, 16]).

**Problem 8.6.** The notion of conjugation  $\sim_p$  defined in (1.2) is very important in symbolic dynamics in connection with the Williams Conjecture [38]. Characterize  $\sim_p$  in  $T(X)$ ,  $P(X)$  and  $\mathcal{I}(X)$  for an infinite set  $X$ . (Kudryavtseva and Mazorchuk [26] have characterized  $\sim_p^*$  (the transitive closure of  $\sim_p$ ) in  $T(X)$ ,  $P(X)$  and  $\mathcal{I}(X)$  for a finite  $X$ , and in  $\mathcal{I}(X)$  for a countably infinite  $X$ .)

## 9 Acknowledgments

The authors would like to thank the referee for the very careful reading of our paper and excellent suggestions, especially the one regarding Definition 4.20.

## References

- [1] J. Araújo, Normal semigroups of endomorphisms of proper independence algebras are idempotent generated, *Proc. Edinburgh Math. Soc. (2)* **45** (2002), 205–217.
- [2] J. Araújo, Idempotent generated endomorphisms of an independence algebra, *Semigroup Forum* **67** (2003), 464–467.
- [3] J. Araújo, M. Edmundo, and S. Givant,  $v^*$ -Algebras, independence algebras and logic, *Internat. J. Algebra Comput.* **21** (2011), 1237–1257.
- [4] J. Araújo and J. Fountain, The origins of independence algebras, *Semigroups and languages*, 54–67, World Sci. Publ., River Edge, NJ, 2004.
- [5] J. Araújo and J. Konieczny, Automorphisms groups of centralizers of idempotents, *J. Algebra* **269** (2003), 227–239.
- [6] J. Araújo and J. Konieczny, Semigroups of transformations preserving an equivalence relation and a cross-section, *Comm. Algebra* **32** (2004), 1917–1935.
- [7] J. Araújo and J. Konieczny, Centralizers in the full transformation semigroup, *Semigroup Forum* **86** (2013), 1–31.
- [8] J. Araújo and J. M. Mitchell, Relative ranks in the monoid of endomorphisms of an independence algebra, *Monatsh. Math.* **151** (2007), 1–10.
- [9] J. Araújo and F. Wehrung, Embedding properties of endomorphism semigroups, *Fund. Math.* **202** (2009), 125–146.
- [10] P. J. Cameron and C. Szabó, Independence algebras, *J. London Math. Soc.* **61** (2000), 321–334.
- [11] A.H. Clifford and G.B. Preston, *The Algebraic Theory of Semigroups*, Mathematical Surveys, No. 7, American Mathematical Society, Providence, Rhode Island, 1964 (Vol. I) and 1967 (Vol. II).
- [12] V.H. Fernandes, Presentations for some monoids of partial transformations on a finite chain: a survey, *Semigroups, algorithms, automata and languages* (Coimbra, 2001), 363–378, World Sci. Publ., River Edge, NJ, 2002.
- [13] J. Fountain and A. Lewin, Products of idempotent endomorphisms of an independence algebra of finite rank, *Proc. Edinburgh Math. Soc.* **35** (1992), 493–500.
- [14] J. Fountain and A. Lewin, Products of idempotent endomorphisms of an independence algebra of infinite rank, *Math. Proc. Cambridge Philos. Soc.* **114** (1993), 303–319.



- [15] O. Ganyushkin and V. Mazorchuk, *Classical Finite Transformation Semigroups: An Introduction*, Algebra and Applications, Volume 9, Springer-Verlag, London, 2010.
- [16] V. Gould, Independence algebras, *Algebra Universalis* **33** (1995), 294–318.
- [17] P. Hell and J. Nešetřil, *Graphs and Homomorphisms*, Oxford University Press, New York, 2004.
- [18] P.M. Higgins, The semigroup of conjugates of a word, *Internat. J. Algebra Comput.* **16** (2006), 1015–1029.
- [19] C. Hollings, From right PP monoids to restriction semigroups: a survey, *European J. Pure Appl. Math.* **2** (2009), 21–57.
- [20] K. Hrbacek and T. Jech, *Introduction to Set Theory*, Third Edition, Taylor & Francis, New York, 1999.
- [21] T. Jech, *Set Theory*, Third Edition, Springer-Verlag, New York, 2006.
- [22] A.S. Kechris, *Classical Descriptive Set Theory*, Springer-Verlag, New York, 1995.
- [23] Y. Kemprasit, Some transformation semigroups whose sets of bi-ideals and quasi-ideals coincide, *Comm. Algebra* **30** (2002), 4499–4506.
- [24] J. Konieczny, Centralizers in the semigroup of injective transformations on an infinite set, *Bull. Austral. Math. Soc.* **82** (2010), 305–321.
- [25] J. Konieczny, Infinite injective transformations whose centralizers have simple structure, *Cent. Eur. J. Math.* **9** (2011), 23–35.
- [26] G. Kudryavtseva and V. Mazorchuk, On conjugation in some transformation and Brauer-type semigroups, *Publ. Math. Debrecen* **70** (2007), 19–43.
- [27] G. Kudryavtseva and V. Mazorchuk, On three approaches to conjugacy in semigroups, *Semigroup Forum* **78** (2009), 14–20.
- [28] G. Lallement, *Semigroups and Combinatorial Applications*, John Wiley & Sons, New York, 1979.
- [29] I. Levi, Green’s relations on  $\mathcal{G}_X$ -normal semigroups, *Math. Japon.* **39** (1994), 19–28.
- [30] I. Levi, Group closures of one-to-one transformations, *Bull. Austral. Math. Soc.* **64** (2001), 177–188.
- [31] I. Levi and B.M. Schein, The semigroup of one-to-one transformations with finite defects, *Glasgow Math. J.* **31** (1989), 243–249.
- [32] I. Levi, B.M. Schein, R.P. Sullivan, and G.R. Wood, Automorphisms of Baer-Levi semigroups, *J. London Math. Soc.* **28** (1983), 492–495.
- [33] D. Lindsey and B. Madison, The lattice of congruences on a Baer-Levi semigroup, *Semigroup Forum* **12** (1976), 63–70.
- [34] F. Otto, Conjugacy in monoids with a special Church-Rosser presentation is decidable, *Semigroup Forum* **29** (1984), 223–240.
- [35] R.P. Sullivan, Automorphisms of injective transformation semigroups, *Studia Sci. Math. Hungar.* **15** (1980), 1–4.
- [36] R.P. Sullivan,  $BQ$ -semigroups of transformations, *Acta Sci. Math. (Szeged)* **75** (2009), 59–74.

- [37] È.G. Šutov, Semigroups of one-to-one transformations, *Dokl. Akad. Nauk SSSR* **140** (1961), 1026–1028 (Russian); translated as *Soviet Math. Dokl.* **2** (1961) 1319–1321.
- [38] R.F. Williams, Classification of subshifts of finite type, *Ann. of Math. (2)* **98** (1973), 120–153.
- [39] L. Zhang, Conjugacy in special monoids, *J. Algebra* **143** (1991), 487–497.
- [40] L. Zhang, On the conjugacy problem for one-relator monoids with elements of finite order, *Internat. J. Algebra Comput.* **2** (1992), 209–220.