

Huygen's principle and iterative methods in inverse obstacle scattering

Olha Ivanyshyn*, Rainer Kress* and Pedro Serranho†

Abstract

The main purpose of this paper is to establish relations between three iterative numerical methods based on Huygen's principle that arose almost simultaneously to solve a particular inverse problem. The inverse problem we consider in this paper is to determine the shape of an obstacle from the knowledge of the far field pattern for scattering of time-harmonic plane waves. In the case of scattering from a sound-soft obstacle, we will interpret Huygen's principle as a system of two integral equations for the unknown boundary of the scatterer and the induced surface flux, i.e., the unknown normal derivative of the total field on the boundary. Reflecting the ill-posedness of the inverse obstacle scattering problem these integral equations are ill-posed. They are linear with respect to the unknown flux and nonlinear with respect to the unknown boundary. In this paper we establish relations and connections between various possibilities for their iterative solution. We also compare them with the usual Newton method applied to the boundary to far-field operator and illustrate that one can get alternatives to this method that imply less computational costs and similar qualitative results. Given a current approximation for the boundary, one of the two equations can be solved for the flux and then the other equation can be linearized with respect to the boundary to obtain an update. Obviously, this possibility offers two options for choosing the linear equation for the flux. A second possibility is to interpret the

*Institut für Numerische und Angewandte Mathematik, Universität Göttingen, 37083 Göttingen, Germany

†CEMAT, Instituto Superior Técnico, Lisbon, Portugal and Polytechnical Institute of Leiria, 2411-901 Leiria, Portugal

equations as a nonlinear system for two unknowns and linearize simultaneously with respect to both the boundary and the flux. We will describe the basic ideas of these approaches and their interrelations.

1 Introduction

The scattering of a time-harmonic acoustic incident wave u^i by a sound-soft obstacle D , that is, a bounded domain $D \subset \mathbb{R}^m$ for $m = 2, 3$ with a connected complement, is modeled by an exterior boundary value problem for the Helmholtz equation

$$\Delta u^s + k^2 u^s = 0 \quad \text{in } \mathbb{R}^m \setminus \bar{D} \quad (1.1)$$

with positive wave number k . The scattered wave u^s has to satisfy the Dirichlet boundary condition

$$u^i + u^s = 0 \quad \text{on } \partial D \quad (1.2)$$

and the Sommerfeld radiation condition

$$\frac{\partial u^s}{\partial r} - ik u^s = o\left(\frac{1}{r^{\frac{m-1}{2}}}\right), \quad r = |x| \rightarrow \infty, \quad (1.3)$$

uniformly for all directions. The total wave u is obtained via superposition $u = u^i + u^s$ and, for most of this presentation we assume the incident wave to be a plane wave, that is,

$$u^i(x, d) = e^{ikx \cdot d}$$

where the unit vector d is the direction of propagation.

Although most of our analysis has extensions to the impedance and/or the Neumann boundary condition, for the sake of exposition and simplicity, we have deliberately chosen to confine our presentation to the case of the Dirichlet boundary condition.

The Sommerfeld radiation condition characterizes outgoing waves and ensures uniqueness for the obstacle scattering problem (see [3]). For brevity, solutions u^s to the Helmholtz equation that satisfy the Sommerfeld radiation condition are called radiating solutions. They can be shown to have an asymptotic behavior of the form

$$u^s(x) = \frac{e^{ik|x|}}{|x|^{\frac{m-1}{2}}} \left\{ u_\infty(\hat{x}) + O\left(\frac{1}{|x|}\right) \right\}, \quad |x| \rightarrow \infty, \quad \hat{x} := \frac{x}{|x|}, \quad (1.4)$$

uniformly with respect to all directions. The function u_∞ is known as the *far field pattern* of the scattered wave and is an analytic function of \hat{x} on the unit sphere $S^{m-1} := \{x \in \mathbb{R}^m : |x| = 1\}$. As one of the most important tools in scattering theory, Rellich's lemma (see Theorem 2.13 in [3]) provides a one-to-one correspondence between a radiating solution u^s to the Helmholtz equation and its far field pattern u_∞ in the sense that $u_\infty = 0$ on S^{m-1} (or on an open subset of S^{m-1}) implies that $u^s = 0$ in its domain of definition.

The *inverse scattering problem* that we are concerned with is to determine the shape and location of the scatterer D from the knowledge of the far field pattern u_∞ for one incident plane wave. We note that this inverse problem is nonlinear in the sense that the scattered wave depends nonlinearly on the scatterer D . More importantly, it is ill-posed since the determination of D does not depend continuously on the far field pattern in any reasonable norm.

The question of uniqueness for the inverse scattering problem for sound-soft obstacles is not yet completely resolved. However, the far field pattern for one incident plane wave is sufficient to uniquely determine the obstacle D if we have the a priori information that the diameter of D is less than $2c_m/k$ where c_m is the smallest positive zero of the Bessel function $J_{m/2}$ of order $m/2$, see [5, 10]. For detailed presentations on inverse obstacle scattering including the issue of uniqueness we refer to the monographs [1, 3, 33] and the surveys [2, 4, 26, 35, 36].

Here, for the foundation of a class of iterative reconstruction methods we start from Green's representation for the scattered wave for a sound-soft obstacle

$$u^s(x) = - \int_{\partial D} \frac{\partial u}{\partial \nu}(y) \Phi(x, y) ds(y), \quad x \in \mathbb{R}^m \setminus \bar{D}, \quad (1.5)$$

in terms of the fundamental solution

$$\Phi(x, y) = \begin{cases} \frac{i}{4} H_0^{(1)}(k|x-y|), & x \neq y, \quad m = 2, \\ \frac{e^{ik|x-y|}}{4\pi|x-y|}, & x \neq y, \quad m = 3, \end{cases}$$

where $H_0^{(1)}$ denotes the Hankel function of the first kind and order zero (see Theorem 3.12 in [3]). The unit normal vector ν to ∂D is assumed to be directed into the exterior of D . The far field pattern of (1.5) is given by

$$u_\infty(\hat{x}) = -\gamma_m \int_{\partial D} \frac{\partial u}{\partial \nu}(y) e^{-ik\hat{x}\cdot y} ds(y), \quad \hat{x} \in S^{m-1}, \quad (1.6)$$

where

$$\gamma_m = \begin{cases} \frac{e^{i\frac{\pi}{4}}}{\sqrt{8\pi k}}, & m = 2, \\ \frac{1}{4\pi}, & m = 3. \end{cases}$$

The representations (1.5) and (1.6) of the scattered field and its far field pattern in terms of its secondary sources on the boundary is known as Huygen's principle.

In view of the sound-soft boundary condition from (1.5) we conclude that

$$u^i(x) = \int_{\partial D} \frac{\partial u}{\partial \nu}(y) \Phi(x, y) ds(y), \quad x \in \partial D, \quad (1.7)$$

Now we can interpret Huygen's principle, that is, (1.6) and (1.7) as a system of two integral equations for the unknown boundary ∂D of the scatterer and the induced surface flux

$$\varphi := -\frac{\partial u}{\partial \nu} \quad \text{on } \partial D.$$

Although, in principle, we are not interested in the surface flux in methods based on the simultaneous solution of (1.6) and (1.7) we cannot avoid solving also for φ . It is convenient to call (1.6) the data equation since it contains the given far field for the inverse problem and (1.7) as the field equation since it represents the boundary condition. Both equations are linear with respect to the flux and nonlinear with respect to the boundary. Equation (1.6) is severely ill-posed whereas (1.7) is only mildly ill-posed.

There exist three options for an iterative solution of (1.6) and (1.7). Given a current approximation for the boundary ∂D we can solve the mildly ill-posed integral equation of the first kind (1.7) for φ . Then, keeping φ fixed we linearize the equation (1.6) with respect to ∂D to update the boundary approximation. This approach has been suggested by Johansson and Sleeman [23] and analyzed further by Ivanyshyn and Johansson [19, 20]. Here, the burden of the ill-posedness and nonlinearity is put on one equation. A more even distribution of the difficulties is obtained by reversing the roles of (1.6) and (1.7), that is, by solving the severely ill-posed equation (1.6) for φ and linearize (1.7) for gaining the boundary update. In a slight modification, this approach may be interpreted also as a decomposition method since to some extent it separates the ill-posedness and the nonlinearity. It combines

the traditional decomposition method with elements of Newton iterations. Therefore it has also been termed as a hybrid method and as such analyzed in a series of papers by Kress and Serranho [27, 29, 30, 37, 38, 39].

Following ideas first developed for the Laplace equation by Kress and Rundell [28], one also can solve the system (1.6) and (1.7) simultaneously for ∂D and φ by Newton iterations, that is, by linearizing both equations with respect to both unknowns. This approach has been intensively studied by Ivanyshyn, Johansson and Kress [15, 16, 17, 18, 21, 22].

2 Parameterized operators and derivatives

In order to define the operators occurring in (1.6) and (1.7) rigorously, for ease of presentation, we restrict ourselves to boundaries ∂D that can be parameterized through mapping them globally onto the unit sphere S^{m-1} , that is,

$$\partial D = \{p(\hat{x}) : \hat{x} \in S^{m-1}\},$$

for some injective C^∞ function $p : S^{m-1} \rightarrow \mathbb{R}^m$. As simple example, the reader should consider the case of starlike domains, where

$$p(\hat{x}) = r(\hat{x})\hat{x}, \quad \hat{x} \in S^{m-1}, \quad (2.1)$$

with a radial distance function $r : S^{m-1} \rightarrow (0, \infty)$. Now, we introduce the parameterized single-layer operator and far field operator

$$A, A_\infty : C^2(S^{m-1}, \mathbb{R}^m) \times L^2(S^{m-1}, \mathbb{C}) \rightarrow L^2(S^{m-1}, \mathbb{C})$$

by

$$A(p, \psi)(\hat{x}) := \int_{S^{m-1}} \Phi(p(\hat{x}), p(\hat{y}))\psi(\hat{y}) ds(\hat{y}), \quad \hat{x} \in S^{m-1}, \quad (2.2)$$

and

$$A_\infty(p, \psi)(\hat{x}) := \gamma_m \int_{S^{m-1}} e^{-ik\hat{x} \cdot p(\hat{y})}\psi(\hat{y}) ds(\hat{y}), \quad \hat{x} \in S^{m-1}. \quad (2.3)$$

Then the equations (1.6) and (1.7) can be written in the operator form

$$A_\infty(p, \psi) = u_\infty \quad (2.4)$$

and

$$A(p, \psi) = -u^i \circ p \quad (2.5)$$

where we have incorporated the surface element into the density function via

$$\psi(\hat{x}) := J(\hat{x}) \varphi(p(\hat{x})) \quad (2.6)$$

with the Jacobian J of the mapping p . The linearization of these equations requires the Fréchet derivatives of the operators A and A_∞ with respect to p . According to [32], these can be obtained by formally differentiating their kernels with respect to p , that is,

$$A'(p, \psi; q)(\hat{x}) = \int_{S^{m-1}} \text{grad}_x \Phi(p(\hat{x}), p(\hat{y})) \cdot [q(\hat{x}) - q(\hat{y})] \psi(\hat{y}) ds(\hat{y}) \quad (2.7)$$

and

$$A'_\infty(p, \psi; q)(\hat{x}) = -ik\gamma_m \int_{S^{m-1}} e^{-ik\hat{x}\cdot p(\hat{y})} \hat{x} \cdot q(\hat{y}) \psi(\hat{y}) ds(\hat{y}) \quad (2.8)$$

for $\hat{x} \in S^{m-1}$. The operators A' and A'_∞ are linear with respect to q and they represent linearizations in the sense that

$$\|A(p+q, \psi) - A(p, \psi) - A'(p, \psi; q)\|_{L^2(S^{m-1})} = o(\|q\|_{C^2(S^{m-1})}^2)$$

and

$$\|A_\infty(p+q, \psi) - A_\infty(p, \psi) - A'_\infty(p, \psi; q)\|_{L^2(S^{m-1})} = o(\|q\|_{C^2(S^{m-1})}^2).$$

3 Iterative solution of the integral equations

3.1 Linearization of the data equation

Johansson and Sleeman [23] suggested the following iterative method for approximately solving the system (2.4) and (2.5). For fixed p , provided k^2 is not a Dirichlet eigenvalue of the negative Laplacian in D , both in a Hölder space setting $A(p, \cdot) : C^{0,\alpha}(S^{m-1}) \rightarrow C^{1,\alpha}(S^{m-1})$ or in a Sobolev space setting $A(p, \cdot) : H^{-1/2}(S^{m-1}) \rightarrow H^{1/2}(S^{m-1})$, the operator $A(p, \cdot)$ is a homeomorphism (see [3]). In this case, given an approximation to the boundary parameterization p we can solve the field equation (2.5) for the density ψ . Then, keeping ψ fixed, we linearize the data equation (2.4) with respect to p to obtain the linear equation

$$A'_\infty(p, \underbrace{[A(p, \cdot)]^{-1}(u^i \circ p)}_{-\psi}; q) = -u_\infty - A_\infty(p, \underbrace{[A(p, \cdot)]^{-1}(u^i \circ p)}_{-\psi}) \quad (3.1)$$

for q to update the parameterization p via $p + q$. This procedure is repeated until some stopping criterium is satisfied.

In principle, the parameterization of the update

$$\partial D_q = \{p(\hat{x}) + q(\hat{x}) : \hat{x} \in S^{m-1}\}$$

is not unique. To cope with the ambiguity, the simplest possibility is to allow only perturbations of the form

$$q = z(\nu \circ p) \tag{3.2}$$

with a scalar function z defined on the unit sphere S^{m-1} and denote the corresponding linear space of normal L^2 vector fields by $L^2_{\text{normal}}(S^{m-1})$.

For fixed p the operator $A'_\infty(p, [A(p, \cdot)]^{-1}(u^i \circ p); \cdot)$ has a smooth kernel and therefore is severely ill-posed. This requires stabilization, for example, via Tikhonov regularization. The following result which was obtained in [17] for the two-dimensional case ensures injectivity and dense range as prerequisites for Tikhonov regularization.

Theorem 3.1 *Assume that k^2 is not a Neumann eigenvalue of the negative Laplacian in D . Then the operator*

$$A'_\infty(p, [A(p, \cdot)]^{-1}(u^i \circ p); \cdot) : L^2_{\text{normal}}(S^{m-1}) \rightarrow L^2(S^{m-1})$$

is injective and has dense range.

Proof. We abbreviate $\psi := -[A(p, \cdot)]^{-1}(u^i \circ p)$ and assume that $q = z(\nu \circ p)$ satisfies $A'_\infty(p, \psi; q) = 0$. Then, from (2.8) we observe that the double-layer potential

$$v(x) = \int_{S^{m-1}} \nu(p(\hat{y})) \cdot \text{grad}_x \Phi(x, p(\hat{y})) \psi(\hat{y}) z(\hat{y}) ds(\hat{y})$$

has vanishing far field pattern. Hence, by Rellich's lemma we can conclude that v vanishes outside of D . By the L^2 jump relations for double-layer potentials together with the Fredholm alternative in dual systems it can be concluded that z is continuous (see the proof of Theorem 5.5 in [3]). This in turn, by the jump relations, implies that v solves the homogeneous Neumann problem in D and therefore vanishes identically in D by our assumption on k . A third application of the jump relations yields $\psi z = 0$ on S^{m-1} . However, by

Holmgren's theorem $\psi = -[A(p, \cdot)]^{-1}(u^i \circ p)$ cannot vanish on open subsets of S^{m-1} , since it represents the normal derivative $\partial u / \partial \nu$ of the total field u for scattering of the plane wave u^i from D . Consequently, we have $z = 0$ on S^{m-1} .

Conversely, let g be a solution to the homogeneous adjoint equation

$$\nu(p(\hat{y})) \cdot \int_{S^{m-1}} \hat{x} e^{ik \hat{x} \cdot p(\hat{y})} \psi(\hat{y}) g(\hat{x}) ds(\hat{x}) = 0, \quad \hat{y} \in S^{m-1}.$$

Then, keeping in mind that ψ cannot vanish on open subsets of S^{m-1} , this implies that the Herglotz wave function

$$w(z) := \int_{S^{m-1}} e^{ik \hat{x} \cdot z} g(\hat{x}) ds(\hat{x}), \quad z \in \mathbb{R}^m,$$

has a vanishing normal derivative on ∂D . By the assumption of the theorem on k , this implies that w vanishes identically in D and by analyticity w vanishes everywhere. Therefore, the kernel g of the Herglotz wave function must be identically zero (see Theorem 3.15 in [3]). \square

We can relate the above approach to traditional Newton iterations for solving the inverse obstacle scattering problem. Denoting by $F : p \rightarrow u_\infty$ the operator that maps the boundary ∂D represented by the parameterization p onto the far field pattern for scattering of the incident wave u^i from D , the inverse problem is equivalent to solving the nonlinear operator equation

$$F(p) = u_\infty. \tag{3.3}$$

With the above notations, in the case when k^2 is not a Dirichlet eigenvalue of the negative Laplacian in D , we can represent

$$F(p) = -A_\infty(p, [A(p, \cdot)]^{-1}(u^i \circ p)). \tag{3.4}$$

By the product and chain rule this implies the Fréchet derivative

$$\begin{aligned} F'(p; q) &= -A'_\infty(p, [A(p, \cdot)]^{-1}(u^i \circ p); q) \\ &\quad + A_\infty(p, [A(p, \cdot)]^{-1} A'(p, [A(p, \cdot)]^{-1}(u^i \circ p); q)) \\ &\quad - A_\infty(p, [A(p, \cdot)]^{-1} ((\text{grad } u^i) \circ p) \cdot q). \end{aligned} \tag{3.5}$$

Hence, the iteration scheme as given through (3.1) can be interpreted as Newton iterations for (3.3) with the derivate of F approximated through the first term in the representation (3.5) only, which corresponds to keeping $\psi = -[A(p, \cdot)]^{-1}(u^i \circ p)$ fixed in the linearization process.

The iteration scheme given through (3.1) has been successfully numerically implemented in two dimensions (see [18, 23]). As to be expected from the close relations to Newton iterations for (3.3) as just pointed out, the quality of the reconstructions can compete with those of Newton iterations at the benefit of reduced computational costs.

The above method of linearizing the data equation has been extended to the case of sound-hard obstacles [20].

3.2 Simultaneous linearization of both equations

Extending a method proposed by Kress and Rundell [28] for an inverse Dirichlet problem for the Laplace equation, a second approach for iteratively solving the system (2.4) and (2.5) consists in simultaneously linearizing both equations with respect to both unknowns. In this case, given approximations p and ψ both for the boundary parameterization and the density we obtain the system of linear equations

$$A'_\infty(p, \psi; q) + A_\infty(p, \chi) = -A_\infty(p, \psi) + u_\infty \quad (3.6)$$

and

$$A'(p, \psi; q) + ((\text{grad } u^i) \circ p) \cdot q + A(p, \chi) = -A(p, \psi) - u^i \circ p. \quad (3.7)$$

This system has to be solved for q and χ in order to obtain updates $p + q$ for the boundary parameterization and $\psi + \chi$ for the density. This procedure is repeated until some stopping criterium is satisfied.

Again we restrict the updates to normal fields of the form (3.2) and note that due to the smoothness of the kernels both equations (3.6) and (3.7) are severely ill-posed and require regularization with respect to both variables. In particular, for the parameterization update it is appropriate to incorporate penalties for Sobolev norms of q to guarantee smoothness of the boundary, whereas for the density L^2 penalty terms on χ are sufficient.

The simultaneous iterations (3.6) and (3.7) exhibit some relations to the Newton iteration

$$F'(p; q) = u_\infty - F(p) \quad (3.8)$$

obtained by linearization of (3.3).

Theorem 3.2 *Assume that k^2 is not a Dirichlet eigenvalue of the negative Laplacian in D and set $\psi := -[A(p, \cdot)]^{-1}(u^i \circ p)$. Provided q satisfies (3.8) then q and*

$$\chi := -[A(p, \cdot)]^{-1}(A'(p, \psi; q) + ((\text{grad } u^i) \circ p) \cdot q)$$

satisfy (3.6) and (3.7). Conversely, if q and χ solve (3.6) and (3.7) then q satisfies (3.8).

Proof. From (3.4) and the definition of ψ we have

$$F'(p; q) = u_\infty - A_\infty(p, \psi).$$

The representation (3.5) of the derivative of F yields

$$F'(p, q) = A'_\infty(p, \psi; q) + A_\infty(p, \chi)$$

and combining this with the previous equation establishes that (3.6) holds. From the definition of χ we observe

$$A(p, \chi) + A'(p, \psi; q) + ((\text{grad } u^i) \circ p) \cdot q = 0$$

Therefore, in view of $A(p, \psi) = -u^i \circ p$ we also have (3.7) satisfied.

Conversely, the second equation (3.7) implies that

$$\chi = -[A(p, \cdot)]^{-1}(A'(p, \psi; q) + ((\text{grad } u^i) \circ p) \cdot q)$$

and inserting this into (3.6) leads to

$$A'_\infty(p, \psi; q) - A_\infty(p, [A(p, \cdot)]^{-1}(A'(p, \psi; q) + ((\text{grad } u^i) \circ p) \cdot q)) = -A_\infty(p, \psi) + u_\infty$$

and via (3.5) this implies (3.8). \square

Based on Theorem 3.2 known results (see [3, 25]) on injectivity and dense range of the derivative F' can be carried over to the system (3.6) and (3.7) (see [16]).

Theorem 3.2 also illustrates the difference between the iteration method based on (3.6) and (3.7) and the Newton iterations for (3.3). In general, when performing (3.6) and (3.7) in the sequence of updates the relation $A(p, \psi) = -(u^i \circ p)$ between the current approximations p and ψ for the parameterization and the density will not be satisfied. This observation also

indicates a possibility to use (3.6) and (3.7) for implementing a Newton scheme for (3.3). We only need to change the update $\psi + \chi$ for the density by

$$-[A(p + q, \cdot)]^{-1}(u^i \circ (p + q)),$$

that is, at the expense of throwing away χ and solving a boundary integral equation for a new density. In the literature, the implementation of Newton iterations for (3.3), in general, is based on the characterization of the Fréchet derivative of F which is given by the far field $F'(p; q) = v_\infty$ of the radiating solution v to the Helmholtz equation in the exterior of D satisfying the Dirichlet condition

$$v \circ q = -(\text{grad } u) \circ q \tag{3.9}$$

in terms of the total field u (see Theorem 5.14 in [3]). For numerical implementations in three dimensions we refer to [8, 11].

The iteration scheme given through (3.6) and (3.7) has been numerically tested in two dimensions (see [21]). The performance of the method based on the linearization of the data equation from the previous subsection and the simultaneous linearization of this subsection, respectively, have been compared in [18]. In general, both methods work equally well with the following slight difference in strategy as observed from a large number of examples. The first method usually performs such that within the first iterations the initial guess is moved towards the center of the obstacle without much change in the shape and then first the illuminated part and then the shadow region of the obstacle is found. On the other hand, the second method searches right away for location and shape simultaneously. In some sense, this is to be expected since the far field pattern describes the asymptotic behavior of the scattered field and therefore is more sensible to the location of the scatterer than to its shape. Consequently, as the field equation is solved first and then (for a fixed flux density) the data equation is linearized, the first method first reconstructs the location and only later the shape. Since in the second method both the data and the field equation are linearized simultaneously, the search for location and shape also starts simultaneously. However, so far no deeper theoretical explanation for this behavior could be given.

Although there are a few results available on the convergence of regularized Newton iterations (see [13, 14, 34]) for inverse obstacle scattering problems, i.e., for the solution of (3.3) via (3.8), this issue is not satisfactorily resolved. Despite the progress made by Hohage [13, 14] with this respect, so far it has not been clarified whether the general results on the

solution of ill-posed nonlinear equations in a Hilbert space setting (among others see [6, 7]) are applicable to inverse obstacle scattering or, in general, to inverse boundary value problems in the frame work of solving the operator equation (3.3). The more problem oriented approach of Potthast [34] for a convergence analysis suffers from the restrictive assumption of a non-vanishing normal derivative of the total field on the boundary ∂D in the case of exact data. Furthermore, in the analysis for noisy data, convergence for the noise level tending to zero, as usual, requires a stopping rule and with this particular rule the method has not yet been numerically implemented. These difficulties with establishing convergence results are also present in the iterations of this section, that is, currently the convergence issue remains an open problem.

The above method of simultaneous linearization has been extended to the case of sound-soft [22] and sound-hard [31] cracks in two dimensions. Here we note that as opposed to the hybrid method of the following section for crack reconstructions no additional penalty term on the length of the crack is required. In addition, it has been also applied to reconstructions of sound-soft or sound-hard scatterers from the modulus of the far field pattern [15]. In order to avoid the exceptional values occurring, for example, in Theorem 3.1 and 3.2 modifications using combined single- and double-layer potentials are described in [19].

4 Decomposition methods revisited

To evenly distribute the burden of the ill-posedness and the nonlinearity of the inverse obstacle scattering problem, instead of solving the field equation (2.5) for the density and then linearizing the data equation one can also think of solving the severely ill-posed data equation (2.5) for the density and linearize the only mildly ill-posed field equation to update the boundary. In this case, given an approximation for the boundary parameterization p we first solve the data equation (2.4) for the density ψ . Then, keeping ψ fixed, we linearize the field equation (2.5), that is, we obtain the linear equation

$$A'(p, \psi; q) + ((\text{grad } u^i) \circ p) \cdot q = -A(p, \psi) - u^i \circ p \quad (4.1)$$

for q to update the parameterization p via $p + q$. Again, this procedure of alternatingly solving (2.4) and (4.1) must be repeated until some stopping criterium is satisfied. To some extent this procedure mimics a decomposition

method in the sense that it decomposes the inverse problem into a severely ill-posed linear problem and an at most mildly ill-posed nonlinear problem.

To our knowledge, this approach has not been implemented numerically. However, the method that has been suggested and investigated by Kress and Serranho can be considered as a slight modification of the above procedure. In this method, given an approximation p for the parameterization of the boundary, the data equation, that is,

$$A_\infty(p, \psi) = u_\infty \quad (4.2)$$

is solved for the density ψ via regularization. Injectivity and dense range of the operator $A_\infty(p, \cdot) : L^2(S^{m-1}) \rightarrow L^2(S^{m-1})$ are guaranteed by Theorem 5.17 in [3] provided k^2 is not a Dirichlet eigenvalue for the negative Laplacian in D . Then we define the single-layer potential

$$u^s(x) = \int_{S^{m-1}} \Phi(x, p(\hat{y})) \psi(\hat{y}) ds(\hat{y})$$

and evaluate the boundary values of $u := u^i + u^s$ and its derivatives on the surface represented by p via the jump relations. Finally we find an update $p + q$ for the boundary by linearizing the boundary condition $u \circ (p + q) = 0$, that is, by solving the linear equation

$$u \circ p + ((\text{grad } u) \circ p) \cdot q = 0 \quad (4.3)$$

for q . In an obvious way, the two steps (4.2) and (4.3) are iterated. Again for uniqueness of the update representation we allow only perturbations of the form (3.2). Then, by Holmgren's theorem, injectivity for the linear equation (4.3) can be established at the exact boundary.

After introducing the operator

$$\tilde{A}(p, \psi; q)(\hat{x}) = \int_{S^{m-1}} \text{grad}_x \Phi(p(\hat{x}), p(\hat{y})) \cdot q(\hat{x}) \psi(\hat{y}) ds(\hat{y}) - \frac{1}{2} \frac{\psi(\hat{x}) [\nu(p(\hat{x})) \cdot q(\hat{x})]}{J(\hat{x})}$$

and observing the jump relations for the single-layer potential and (2.6) we can rewrite (4.3) into the form

$$\tilde{A}(p, \psi; q) + ((\text{grad } u^i) \circ p) \cdot q = -A(p, \psi) - u^i \circ p. \quad (4.4)$$

Comparing this with (4.1) we observe that the Fréchet derivative of A with respect to p is replaced by the operator \tilde{A} in the sense that we linearize only

with respect to the evaluation surface for the single-layer potential but not with respect to the integration surface.

Since this approach combines ideas of the decomposition method due to Kirsch and Kress [24] with aspects of Newton iterations, it was termed a hybrid method. Convergence for exact data in the spirit of Potthast [34] and a stopping rule for noisy data were investigated in [38].

Without giving any details on the computations, in Fig. 1–4 we present some examples for reconstructions by the above hybrid method. The implementation was based on solving (4.3) and setting up the linear system via the jump relations rather than the equivalent form (4.4). The numerical quadratures were based on Wienert’s method [40] as described in section 3.6 of [3] (see also [9]) and the radial distance functions in the sense of (2.1) were approximated by linear combinations of spherical harmonics up to order eight. In each example the figure on the left hand side gives the exact boundary shape and the figure on the right the reconstruction with one incident wave in direction of the arrow. The reconstructions are obtained with 2% random noise added to the synthetic far field pattern. The wave number is $k = 1$.

This approach has been successfully extended to the case of sound-hard [30] and impedance obstacles [37]. For the impedance boundary condition both the shape and the impedance function can be reconstructed. In addition, the method has also been used for the reconstruction of sound-soft cracks in two dimensions [29]. In this case, it was necessary to impose a penalty term on the length of the crack. Otherwise, the algorithm would reconstruct cracks with length shrinking to zero.

Although the linearization of the homogeneous boundary Neumann condition (see [30]) is slightly more technical than the linearization of the homogeneous Dirichlet condition as described above, it is considerably less involved than the derivation of the boundary condition for the corresponding boundary to far field map F for the sound hard case (see [12]).

5 Conclusion

We have described and established connections between three closely related methods for solving the inverse obstacle scattering problem via an iterative solution of nonlinear integral equations and have also illuminated their connections to Newton iterations for the boundary to far field map. We have also illustrated that these methods are a good alternative to Newton iter-

ations in the sense that one gets the same quality of reconstruction with fewer effort. These methods have been shown to be applicable for a range of boundary conditions. Their practical feasibility has been established in the literature through a variety of numerical examples. More work is required for the theoretical foundation through an appropriate convergence analysis and for further implementations in three dimensions.

Acknowledgment. The research of O.I. was partially supported by the German Research Foundation DFG through the collaborative research center SFB 755. The research of P.S. was partially supported by the Calouste Gulbenkian Foundation.

References

- [1] Cakoni, F. and Colton, D.: *Qualitative Methods in Inverse Scattering Theory: An Introduction*. Springer, New York 2005.
- [2] Colton, D., Coyle, J., and Monk, P.: Recent developments in inverse acoustic scattering theory. *SIAM Review* **42**, 369–414 (2000).
- [3] Colton, D. and Kress, R.: *Inverse Acoustic and Electromagnetic Scattering Theory*. 2nd. ed. Springer, Berlin 1998.
- [4] Colton, D. and Kress, R.: Using fundamental solutions in inverse scattering. *Inverse Problems* **22**, R49–R66
- [5] Colton, D. and Sleeman, B.D.: Uniqueness theorems for the inverse problem of acoustic scattering. *IMA J. Appl. Math.* **31**, 253–259 (1983).
- [6] Deuffhard, P., Engl, H.W., and Scherzer, O.: A convergence analysis of iterative methods for the solution of nonlinear ill-posed problems under affinity invariant conditions. *Inverse Problems* **14**, 1081–1106 (1998).
- [7] Engl, H.W., Hanke, M., and Neubauer, A.: *Regularization of inverse problems*. Kluwer Academic Publisher, Dordrecht 1996.
- [8] Farhat, C., Tezaur, R., and Djellouli, R. : On the solution of three-dimensional inverse obstacle acoustic scattering problems by a regularized Newton method. *Inverse Problems* **18**, 1229–1246 (2002).
- [9] Ganesh, M. and Graham, I.: A high-order algorithm for obstacle scattering in three dimensions. *J. Comp. Phys.*, **198**, 211–242 (2004).

- [10] Gintides, D.: Local uniqueness for the inverse scattering problem in acoustics via the Faber–Krahn inequality. *Inverse Problems* **21**, 1195–1205 (2005).
- [11] Harbrecht, H. and Hohage, T.: Fast methods for three-dimensional inverse obstacle scattering problems. *Jour. Integral Equations and Appl.* **19**, 237–260 (2007).
- [12] Hettlich, F.: Frechet derivatives in inverse obstacle scattering. *Inverse Problems* **14**, 209–210 (1998).
- [13] Hohage, T.: Logarithmic convergence rates of the iteratively regularized Gauss-Newton method for an inverse potential and an inverse scattering problem. *Inverse Problems* **13**, 1279–1299 (1997).
- [14] Hohage, T.: *Iterative Methods in Inverse Obstacle Scattering: Regularization Theory of Linear and Nonlinear Exponentially Ill-Posed Problems*. Dissertation, Linz 1999.
- [15] Ivanyshyn, O.: Shape reconstruction of acoustic obstacles from the modulus of the far field pattern. *Inverse Problems and Imaging* **1**, 609–622 (2007).
- [16] Ivanyshyn, O.: *Nonlinear Boundary Integral Equations in Inverse Scattering*. Dissertation Göttingen 2007.
- [17] Ivanyshyn, O. and Johansson, T.: Boundary integral equations for acoustical inverse sound-soft scattering. *J. Inverse Ill-Posed Probl.* **15**, 1–14 (2007).
- [18] Ivanyshyn, O. and Johansson, T.: Nonlinear integral equation methods for the reconstruction of an acoustically sound-soft obstacle. *J. Integral Equations Appl.* **19**, 289–308 (2007).
- [19] Ivanyshyn, O. and Johansson, T.: A coupled boundary integral equation method for inverse sound-soft scattering. In: *Proceedings of waves 2007. The 8th international conference on mathematical and numerical aspects of waves*, pp. 153–155 University of Reading 2007.
- [20] Ivanyshyn, O. and Johansson, T.: Reconstruction of acoustically sound-hard obstacles from the far field using a boundary integral equation method. In: *Mathematical Methods in Scattering Theory and Biomedical Engineering* (Fotiatis, Massalas, eds.), World Scientific, Singapore, (2008). (to appear)
- [21] Ivanyshyn, O. and Kress, R.: Nonlinear integral equations in inverse obstacle scattering. In: *Mathematical Methods in Scattering Theory and Biomedical Engineering* (Fotiatis, Massalas, eds.), pp 39–50 World Scientific, Singapore, (2006).

- [22] Ivanyshyn, O. and Kress, R.: Inverse scattering for planar cracks via nonlinear integral equations. *Math. Meth. Appl. Sciences* (2007).
- [23] Johansson, T. and Sleeman, B.: Reconstruction of an acoustically sound-soft obstacle from one incident field and the far field pattern. *IMA J. Appl. Math.* **72**, 96–112 (2007).
- [24] Kirsch, A. and Kress, R.: An optimization method in inverse acoustic scattering. In: *Boundary elements IX, Vol 3. Fluid Flow and Potential Applications*, (Brebbia et al, eds.), pp 3–18, Springer-Verlag, Berlin 1987.
- [25] Kress, R.: Integral equation methods in inverse acoustic and electromagnetic scattering. In: *Boundary Integral Formulations for Inverse Analysis* (Ingham and Wrobel, eds.), pp 67–92, Computational Mechanics Publications, Southampton 1997.
- [26] Kress, R.: Acoustic scattering: Scattering by obstacles. In: *Scattering*, (Pike, Sabatier, eds.) pp. 52–73, Academic Press, London 2001.
- [27] Kress, R.: Newton’s Method for inverse obstacle scattering meets the method of least squares. *Inverse Problems* **19**, 91–104 (2003).
- [28] Kress, R. and Rundell, W.: Nonlinear integral equations and the iterative solution for an inverse boundary value problem. *Inverse Problems* **21**, 1207–1223 (2005).
- [29] Kress R. and Serranho P.: A hybrid method for two-dimensional crack reconstruction, *Inverse Problems* **21**, 773–784 (2005).
- [30] Kress R. and Serranho P.: A hybrid method for sound-hard obstacle reconstruction. *J. Comput. Appl. Math.* **204**, 418–427 (2007).
- [31] Lee, Kuo-Ming: Inverse scattering via nonlinear integral equations for a Neumann crack *Inverse Problems* **22**, 1989–2000 (2006).
- [32] Potthast, R.: Fréchet differentiability of boundary integral operators in inverse acoustic scattering. *Inverse Problems* **10**, 431–447 (1994).
- [33] Potthast, R.: *Point-Sources and Multipoles in Inverse Scattering Theory*. Chapman & Hall, London 2001.
- [34] Potthast, R.: On the convergence of a new Newton-type method in inverse scattering. *Inverse Problems* **17**, 1419–1434 (2001).

- [35] Potthast, R.: Sampling and probe methods - an algorithmical view. *Computing* **75**, 215–235 (2005).
- [36] Potthast, R.: A survey on sampling and probe methods for inverse problems. *Inverse Problems* (to appear).
- [37] Serranho, P.: A hybrid method for inverse scattering for shape and impedance. *Inverse Problems* **22**, 663–680 (2006).
- [38] Serranho, P.: A hybrid method for sound-soft obstacles in 3D. *Inverse Problems and Imaging* **1**, 691–712 (2007).
- [39] Serranho, P.: *A Hybrid Method for Inverse Obstacle Scattering Problems*. Dissertation Göttingen 2007.
- [40] Wienert, L.: *Die numerische Approximation von Randintegraloperatoren für die Helmholtzgleichung im \mathbb{R}^3* . Dissertation Göttingen 1990.




Figure 1: Reconstruction of an acorn shaped domain.




Figure 2: Reconstruction of a pinched acorn shaped domain.




Figure 3: Reconstruction of a star shaped domain.




Figure 4: Reconstruction of a cushion shaped domain.