

High Positive End-expiratory Pressure (PEEP) with Recruitment Maneuvers *versus* Low PEEP during General Anesthesia for Surgery: A Bayesian Individual Patient Data Meta-analysis of Three Randomized Clinical Trials

Guido Mazzinari, Ph.D., Fernando G. Zampieri, M.D., Ph.D., Lorenzo Ball, M.D., Ph.D., Niklas S. Campos, M.D., Thomas Bluth, M.D., Sabine N. T. Hemmes, M.D., Ph.D., Carlos Ferrando, M.D., Ph.D., Julian Librero, M.D., Marina Soro, M.D., Ph.D., Paolo Pelosi, M.D., Marcelo Gama de Abreu, M.D., Ph.D., Marcus J. Schultz, M.D., Ph.D., Ary Serpa Neto, M.D., M.Sc., Ph.D.; for REPEAT on behalf of the PROVHILO, iPROVE, and PROBESE investigators and the PROVE Network investigators*

ANESTHESIOLOGY 2025; 142:72–97



EDITOR'S PERSPECTIVE

What We Already Know about This Topic

- Postoperative pulmonary complications frequently occur
- High positive end-expiratory pressure with recruitment maneuvers can protect the lungs against repetitive tidal recruitment but can potentially cause lung overdistension
- A recent individual patient data meta-analysis pooling the information from three large randomized clinical trials found that subjects ventilated with high positive end-expiratory pressure with

ABSTRACT

Background: The influence of high positive end-expiratory pressure (PEEP) with recruitment maneuvers on the occurrence of postoperative pulmonary complications after surgery is still not definitively established. Bayesian analysis can help to gain further insights from the available data and provide a probabilistic framework that is easier to interpret. The objective was to estimate the posterior probability that the use of high PEEP with recruitment maneuvers is associated with reduced postoperative pulmonary complications in patients with intermediate-to-high risk under neutral, pessimistic, and optimistic expectations regarding the treatment effect.

Methods: Multilevel Bayesian logistic regression analysis was performed on individual patient data from three randomized clinical trials carried out on surgical patients at intermediate to high risk for postoperative pulmonary complications. The main outcome was the occurrence of postoperative pulmonary complications in the early postoperative period. This study examined the effect of high PEEP with recruitment maneuvers *versus* low PEEP ventilation. Priors were chosen to reflect neutral, pessimistic, and optimistic expectations of the treatment effect.

Results: Using a neutral, pessimistic, or optimistic prior, the posterior mean odds ratio for high PEEP with recruitment maneuvers compared to low PEEP was 0.85 (95% credible interval, 0.71 to 1.02), 0.87 (0.72 to 1.04), and 0.86 (0.71 to 1.02), respectively. Regardless of prior beliefs, the posterior probability of experiencing a beneficial effect exceeded 90%. Subgroup analysis indicated a more pronounced effect in patients who underwent laparoscopy (odds ratio, 0.67 [0.50 to 0.87]) and those at high risk for postoperative pulmonary complications (odds ratio, 0.80 [0.53 to 1.13]). Sensitivity analysis, considering severe postoperative pulmonary complications only or applying a different heterogeneity prior, yielded consistent results.

Conclusions: High PEEP with recruitment maneuvers demonstrated a moderate reduction in the probability of postoperative pulmonary complication occurrence, with a high posterior probability of benefit observed consistently across various prior beliefs, particularly among patients who underwent laparoscopy.

(ANESTHESIOLOGY 2025; 142:72–97)

recruitment maneuvers were less likely to develop postoperative pulmonary complications; however, this result was not significant at the standard *P* value threshold

What This Article Tells Us That Is New

- This *post hoc* Bayesian reanalysis of individual patient data from three large randomized clinical trials showed that using high positive end-expiratory pressure and recruitment maneuvers led to a high likelihood of a slight reduction in the probability of complications occurrence

This article is featured in "This Month in ANESTHESIOLOGY," page A1. This article is accompanied by an editorial on p. 6. Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are available in both the HTML and PDF versions of this article. Links to the digital files are provided in the HTML text of this article on the Journal's Web site (www.anesthesiology.org). This article has an audio podcast. This article has a visual abstract available in the online version.

Submitted for publication November 22, 2023. Accepted for publication July 9, 2024. Published online first on July 22, 2024.

Guido Mazzinari, Ph.D.: Department of Anesthesiology and Pain Medicine, La Fe Research Institute, Valencia, Spain; Perioperative Medicine Research Group, Valencia, Spain; and Department of Statistics and Operational Research, Universidad de Valencia, Valencia, Spain.

Copyright © 2024 American Society of Anesthesiologists. All Rights Reserved. ANESTHESIOLOGY 2025; 142:72–97. DOI: 10.1097/ALN.00000000000005170

Postoperative pulmonary complications frequently occur, increasing mortality, hospital length of stay, and healthcare costs.¹ Although the protective role of low tidal volume has been established,² the effect of high positive end-expiratory pressure (PEEP) with recruitment maneuvers during intraoperative ventilation is less certain, because it can protect the lungs against repetitive tidal recruitment but can potentially cause lung overdistension.³ A recent individual patient data meta-analysis pooled the information from three large randomized clinical trials to assess the effect of high PEEP and recruitment maneuvers *versus* low PEEP on postoperative pulmonary complications occurrence.⁴ Subjects ventilated with high PEEP with recruitment maneuvers were less likely to develop postoperative pulmonary complications. However, this result was not significant at the standard *P* value threshold.

Several recent papers used a Bayesian approach to reanalyze data from previous anesthesiology and critical care randomized clinical trials with indeterminate frequentist results to gain further insights.^{5–10} We aimed to reanalyze the individual patient data set from the previous REPEAT frequentist meta-analysis⁴ with a Bayesian approach according to previously published recommendations.⁷ We hypothesized that intraoperative high PEEP with recruitment maneuvers reduced the posterior probability of developing postoperative pulmonary complications.

Materials and Methods

Study Design and Context

The study is an individual patient data meta-analysis using a database pooling individual patients from three recent

randomized clinical studies of intraoperative ventilation during general anesthesia for surgery.⁴ This analysis's protocol and statistical analysis plan were previously published.¹¹ The data harmonization procedure is described in detail elsewhere.¹² Briefly, the three studies shared a similar intraoperative management and outcome definition and compared high PEEP (10 to 12 cm H₂O) with recruitment maneuvers to low PEEP (0 to 5 cm H₂O) without Rm. Of note, two of the studies were two-arm parallel trials, whereas one was a four-arm parallel trial with additional interventions in the early postoperative period. The primary analysis focused on the intraoperative intervention, *i.e.* the high PEEP with recruitment maneuvers strategy. The protocols of the three studies were approved by the institutional review boards of the participating hospitals and published previously.^{13–15} The detailed characteristics of the studies are shown in supplemental table S1 (<https://links.lww.com/ALN/D637>).

Patients

The studies included patients with an intermediate risk of developing postoperative pulmonary complication as assessed by the Assess Respiratory Risk in Surgical Patients in Catalonia (ARISCAT) score¹⁶ who underwent major abdominal surgery. No additional exclusion criteria were added for this analysis.

Data Collected

Baseline characteristics, as well as intraoperative data and postoperative pulmonary complications occurrence, were collected. Of note, for postoperative pulmonary complications, the follow-up was available up to 5 postoperative days for every patient and up to 7 days in one study.¹⁴

Fernando G. Zampieri, M.D., Ph.D.: Department of Critical Care Medicine, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada; and PROVE Network, Alberta Health Services, Edmonton, Alberta, Canada.

Lorenzo Ball, M.D., Ph.D.: IRCCS San Martino Policlinico Hospital, Genoa, Italy; University of Genoa, Genoa, Italy; Department of Surgical Sciences and Integrated Diagnostics, Genoa, Italy; and Hospital Israelita Albert Einstein, São Paulo, Brazil.

Niklas S. Campos, M.D.: Department of Critical Care Medicine, Av Hospital Israelita Albert Einstein, São Paulo, Brazil; and Cardio-Pulmonary Department, Pulmonary Division, Heart Institute, Hospital das Clínicas HCFMUSP, Faculty of Medicine, University of Sao Paulo, Sao Paulo, Brazil.

Thomas Bluth, M.D.: Pulmonary Engineering Group, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, Technical University Dresden, Dresden, Germany.

Sabrina N. T. Hemmes, M.D., Ph.D.: Departments of Intensive Care and of Anesthesiology, Amsterdam University Medical Centers, Amsterdam, The Netherlands.

Carlos Ferrando, M.D., Ph.D.: Department of Anesthesiology and Critical Care, Hospital Clinic de Barcelona, Research Institute August Pi i Sunyer, Barcelona, Spain; and Center of Biomedical Research in Respiratory Diseases, Health Institute Carlos III, Madrid, Spain.

Julian Librero, M.D.: Navarrabiomed-Fundación Miguel Servet, Red de Investigación en Servicios de Salud en Enfermedades Crónicas (REDISSEC), Pamplona, Spain.

Marina Soro, M.D., Ph.D.: INCLIVA Clinical Research Institute, Clinical Hospital, University of Valencia, Valencia, Spain.

Paolo Pelosi, M.D.: IRCCS San Martino Policlinico Hospital, Genoa, Italy; and Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy.

Marcelo Gama de Abreu, M.D., Ph.D.: Departments of Intensive Care and Resuscitation, of Cardiothoracic Anesthesia, and of Outcomes Research, Institute of Anesthesiology, Cleveland Clinic, Cleveland, Ohio.

Marcus J. Schultz M.D., Ph.D.: Department of Intensive Care and Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand; Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom; and Department of Anesthesia, General Intensive Care and Pain Management, Division of Cardiothoracic and Vascular Anesthesia and Critical Care Medicine, Medical University Vienna, Wien, Austria.

Ary Serpa Neto M.D., M.Sc., Ph.D.: Department of Critical Care Medicine and Cardio-Pulmonary Department, Pulmonary Division, São Paulo, Brasil; Department of Intensive Care, Australian and New Zealand Intensive Care Research Centre (ANZIC-RC), Monash University, Melbourne, Australia; and Department of Critical Care, Melbourne Medical School, University of Melbourne, Austin Hospital, Melbourne, Australia.

*Investigators with REPEAT, PROVHILO, iPROVE, PROBESE, and PROVE are listed in the appendix.

Definitions

Postoperative pulmonary complications are defined as a collapsed composite of mild and severe respiratory failure, acute respiratory distress syndrome, pulmonary infection, pleural effusion, atelectasis, pneumothorax, and bronchospasm. The definition of the individual components of the composite is shown in supplemental table S2 (<https://links.lww.com/ALN/D637>).

Outcomes

The outcome of this analysis was the proportion of patients developing one or more postoperative pulmonary complication.

Statistical Methods and Analysis

We used all of the available data without any *a priori* power calculations and carried out a one-stage approach meta-analysis by fitting a multilevel Bayesian logistic model using the *brms* R package,¹⁷ including the effect of the PEEP strategy as population (fixed) effect and the study and site as a random effect modeling heterogeneity

of effect across different studies. The site was introduced as a nested effect for the study (supplemental fig. S1, <https://links.lww.com/ALN/D637>).

Prior Distributions

Priors were set for the fixed effect, *i.e.*, the effect of PEEP strategy, standard deviations of the random effects, and the correlation structure of the random effects. We defined a neutral, a pessimistic, and an optimistic prior probability distribution with moderate belief strength to cover the full range of possible beliefs for the effect of the PEEP strategy as previously recommended and shown in detail in table 1.⁷ We used normal distributions for the effects on the log odds scale (supplemental fig. S2, <https://links.lww.com/ALN/D637>). This allows the null effect to be centered at zero and a symmetrical distribution, which can be transformed back to odds ratios by exponentiating the log odds values. We assigned some strength to our prior beliefs to be weakly informative; *i.e.*, we set them so that we cannot rule out an eventual benefit but can mostly rule out large effect sizes for the intervention and acknowledge a non-negligible chance of the intervention being harmful.⁴

Table 1. Defining Priors to Explain What We Expect the Intervention Will Achieve and How Strong These Assumptions Are

Strength	Belief		
	Neutral	Optimistic	Pessimistic
Weak	We do not know much about the intervention effect. We will use a normal distribution with a mean of 0 and a large SD. The extreme case will be a “flat” prior with an infinite SD. Example: $N \sim (0, 5)$	We believe the intervention is beneficial, but we have no reason to exclude that it can be harmful. We will use a normal distribution with a mean at the expected effect and an SD to retain at least a 30% probability of harm (<i>i.e.</i> , $OR > 1$, log odds > 0).	We believe the intervention is harmful, but we rule out a benefit. We will use a normal distribution with a mean at the expected effect and an SD to retain at least a 30% probability of benefit (<i>i.e.</i> , $OR < 1$, log odds < 0).
Moderate	We do not know whether the intervention may be beneficial or harmful, but we exclude that the effect is large. We will use a normal prior with a mean of 0 and an SD to retain only 2.5% of probability of large effect of more than $OR > 2$ or $OR < 0.5$. Example: $N \sim (0, 0.355)$. <i>We used this prior for the analysis.</i>	We believe the intervention is beneficial, but we cannot exclude that there is a certain chance that it can be harmful. We will use a normal distribution with a mean at the expected effect and an SD to retain at least a 15% probability of harm ($OR > 1$, log odds > 0). <i>We used this prior in the analysis using averaged effect size estimate from the three original studies ($OR = 0.51$, log odds = -0.67).</i>	We believe the intervention is harmful, but we cannot exclude that there is a certain chance that it can be beneficial. We will use a normal distribution with a mean at the expected effect and an SD to retain at least a 15% probability of benefit (<i>i.e.</i> , $OR < 1$, log odds < 0). <i>We used this prior in the analysis using averaged effect size estimate from the three original studies ($OR = 1/0.51$, log odds = 0.67).</i>
Strong	We strongly believe that the intervention does not have an effect. We will use a normal prior with a mean of 0 and an SD to retain only 2.5% of probability of large effect of more than $OR > 1.5$ or $OR < 0.66$. Example: $N \sim (0, 0.205)$	We believe the intervention is beneficial, but there is a very low chance for it to be harmful. We will use a normal distribution with a mean at the expected effect and an SD to retain at least a 5% probability of harm ($OR > 1$, log odds > 0).	We believe the intervention is harmful, but there is a very low chance for it to be beneficial. We will use a normal distribution with a mean at the expected effect and an SD to retain at least a 5% probability of benefit (<i>i.e.</i> , $OR < 1$, log odds < 0).

This table is based on an example to study the intervention effect on a harmful binary outcome, meaning that positive ratios represent harm. In this context, we usually express our initial guess as an effect ratio; for instance, the odds ratio (OR) using the log of this ratio that allows to use normal distribution to model it. Such distribution will be defined by the mean (location) and the standard deviation (SD; scale) and parameterized according to our set of beliefs. For instance, a log odds of 0 represents no effect, *i.e.*, an OR of 1, and a log odds greater than 0 represents higher odds for the event. The normal distributions are reported as $N \sim (\text{mean}, \text{SD})$. The *weakly pessimistic* prior was initially chosen to retain a 0.30 probability of benefit [$Pr(OR < 1)$]; however, because this returned an almost flat prior due to the great variance needed to retain that probability density for benefit with an average of OR of 1.98 that mirrors the effect from the optimistic prior, we opted for a symmetric prior retaining 0.15 probability of benefit, as previously used in a recent study.⁸

The prior distribution for the standard deviations of the random effects, *i.e.* the intervention heterogeneity across studies, was defined as a half-normal distribution with a mean of 0 and a SD = 0.5; this yields prior probabilities of 52% in the reasonable category, 27% in the fairly high category, and 5% in the extreme category (supplemental fig. S3, <https://links.lww.com/ALN/D637>).¹⁸ Furthermore, the prior distribution for the correlation matrix of the random effects was based on the Lewandowski–Kurowicka–Joe distribution with an η parameter of 2 (supplemental fig. S4, <https://links.lww.com/ALN/D637>).¹⁹

We also looked at how different beliefs about the intervention affected the main results as postulated in previously published recommendations.⁷ We did this by treating the estimates from different priors' models as if they were separate studies and fitting an aggregate data Bayesian meta-analysis. This helped us see how much these beliefs influenced the results because the heterogeneity of this meta-analysis is the influence of the priors, because there is no other sampling difference. We used a weakly informative neutral prior similar to the one we used in the main analysis for the effect size and a DuMouchel prior for heterogeneity. We chose such heterogeneity prior following the recommendations on Bayesian reanalysis;⁷ the DuMouchel prior has been shown to provide good coverage for this kind of analysis, not deviating much from uniform priors while providing some regularization.²⁰

We chose the region of practical equivalence as the interval between odds ratios of 0.9 and 1.1. The region of practical equivalence is usually defined based on the context of the study and specific use; it is set based on previous knowledge, expert opinions, or thinking about how parameter values practically affect the outcome; parameters within region of practical equivalence are seen as having similar effects, and any differences between them might not be seen as important in a clinical context.²¹ The extent of the 95% posterior highest density interval overlap with region of practical equivalence helps to understand the chances of an intervention being helpful or harmful. In addition, we defined a threshold for severe harm at an odds ratio of 1.25.

We used the threshold suggested by previously published recommendations in a critical care setting⁷ because we considered that our clinical scenario bears sufficient similarities. We drew 7,000 samples from the posterior distribution after fitting the models with *brms* Hamiltonian Markov No U-turn sampler using each of the previously defined priors and determined how much of the posterior probability density laid in the region of practical equivalence interval or exceeded the threshold for severe harm and determined the posterior probability density of treatment effect using estimated marginal means with the *emmeans* package for R (version 1.8.5).²²

We reported the effect of the intervention both as a marginal effect and as a conditional effect. For the conditional effect, we conditioned on a new random effect simulated by

drawing posterior samples from the random effect mixture distribution of trials and sites. In other words, this conditional effect draws random values from the joint posterior distribution to reflect the observed uncertainty within and across all subjects for the random effects included in the model, thus allowing assessment of the effect of the intervention for hypothetical or unobserved scenarios as an additional assessment of the heterogeneity. Furthermore, we compared the interval of the region of practical equivalence with the 95% posterior highest density interval, as previously recommended, to see whether the highest density interval posterior probability density falls outside the region of practical equivalence.

We performed the following preplanned subgroup analyses by refitting the multilevel logistic regression model by adding an interaction term between treatment and a subgroup of interest and reporting the effect estimated by pairwise contrasts in the groups of interest. We carried out these analyses for the effect of high PEEP and recruitment maneuver *versus* low PEEP in (1) type of surgery, *i.e.*, laparoscopic *versus* open surgery; (2) risk for postoperative pulmonary complications, *i.e.*, ARISCAT score less than 45 *versus* greater than or equal to 45; and (3) body mass index less than 30 *versus* greater than 30. Next, we evaluated potential heterogeneity due to the different design of one of the studies.¹⁴ Therefore, we assessed the effect of (4) type of PEEP selection, *i.e.*, fixed *versus* titrated, in the high PEEP and recruitment maneuver cohort and the (5) use of a postoperative element as part of the tested intervention, *i.e.*, applied postoperative continuous positive airway pressure (CPAP) *versus* standard oxygen support.

We performed sensitivity analyses by (1) refitting the model and using only severe postoperative pulmonary complications as the primary outcome; (2) varying the heterogeneity prior to a half-Cauchy prior¹⁸; and (3) assessing prior–data conflicts by examining the perturbation on posterior distributions of parameters by power-scaling priors with the R *priorsense* package.²³ We could not perform the previously planned analysis including only patients with lung collapse before the intervention as previously specified,¹¹ because we could not identify such a group with the available data. All analyses used R software (version 4.2.3, Core Team, Vienna, Austria).

Results

The pooled database contained 3,836 patients: 1,913 high PEEP and recruitment maneuver patients and 1,923 low PEEP patients. Demographic, perioperative, and care-delivery characteristics were well balanced at baseline between high PEEP and recruitment maneuver and low PEEP patients (supplemental table S3, <https://links.lww.com/ALN/D637>). Intraoperative and ventilatory variables and differences between groups by every outcome

component are shown in supplemental tables S4 and S5 (<https://links.lww.com/ALN/D637>).

The posterior mean effect of high PEEP and recruitment maneuver on the occurrence of postoperative pulmonary complications was 0.85 (95% credible interval, 0.71 to 1.02), 0.87 (0.72 to 1.04), and 0.86 (0.71 to 1.02) for the models fitted with neutral, pessimistic, and optimistic priors, respectively. The posterior probability of a beneficial effect was more than 90% regardless of the prior (table 2; figs. 1 and 2). The conditional effects estimated by considering the random effect of trial and site have similar posterior means but a wider 95% credible interval and some posterior probability of harm, *i.e.*, more than 10% regardless of prior (supplemental table S6 and figs. S5 and S6, <https://links.lww.com/ALN/D637>). The posterior probability density comprised in the region of practical equivalence was always less than 30%. The prior choice had little influence on the results; the estimated heterogeneity I^2 from the aggregate estimate meta-analysis was 0.107. Therefore, approximately 10% of the variance was caused by the priors (supplemental fig. S7, <https://links.lww.com/ALN/D637>).

The effect of high PEEP and recruitment maneuver *versus* low PEEP had a higher posterior probability of benefit in the laparoscopic surgery group compared to the open surgery group. High PEEP and recruitment maneuver was slightly more likely to be beneficial in the high-risk ARISCAT cohort. We did not find any differences between the other subgroups (table 3). As for the main analysis, the conditional effects reported a similar posterior mean with wider 95% credible intervals and some posterior probability of harm (supplemental table S6 and figs. S8 to S12, <https://links.lww.com/ALN/D637>). We found that the posterior distributions changed little when we used only severe postoperative pulmonary complications as the outcome or when we used a different prior for heterogeneity (table 3; fig. 3). In addition, the *post hoc* prior sensitivity did not report any conflicts for the intervention estimates.

Discussion

We applied a Bayesian analysis framework on a large, harmonized data set to estimate the effect of high PEEP and recruitment maneuver ventilation on the occurrence of postoperative pulmonary complications compared to a strategy based on low PEEP, an approach that allows for flexible hypothesis testing, the intuitive use of probabilities.^{24–28} Our results showed that (1) using high PEEP and recruitment maneuver led to a slight reduction in the posterior probability of complications occurrence; (2) this benefit was highly likely in general, although (3) it can be less likely to benefit depending on the variability of patients. Moreover, (4) the different prior distributions did not influence the results. Subgroups and sensitivity analyses showed that (5) the effect of the high PEEP and recruitment maneuver was more beneficial in the laparoscopic and high ARISCAT risk score patients' groups, and no effect was associated with the use of postoperative CPAP or an individualized high PEEP instead of a fixed high PEEP.

The physiologic basis of mechanical ventilation studies in the operating room is that general anesthesia with muscle paralysis causes airway and lung collapse. This can lead to injury due to the excessive pressure or volume applied to the shrunk parenchyma or to the shear force occurring in cyclically opening regions.²⁹ The use of high PEEP with recruitment maneuvers is intended to counteract these deleterious effects. However, apart from a recent meta-analysis of mainly small studies in which the authors acknowledged some publication bias,³⁰ several large randomized clinical trials yielded indeterminate results on the benefit of such a ventilatory strategy. Our results based on a different statistical framework provide a reasonable belief that a high PEEP and recruitment maneuver strategy can be beneficial. The magnitude of this benefit, although not negligible based on the prespecified region of practical equivalence criteria, is probably small. We applied a region of practical equivalence criterion suitable for similar clinical contexts

Table 2. Average Effect Estimates from Main Model for Primary Outcome

Belief	Prior Subject Matter Justification	Odds Ratio (95% Credible Interval)*	Absolute Risk Reduction* (95% Credible Interval)	Probability Mass in the Region of Practical Equivalence	Probability of Benefit Odds Ratio < 1 (Odds Ratio < 0.8; > 1.25)
Neutral	There is no previous firm evidence to believe the intervention is good or bad, but large effect sizes can be reasonably ruled out.	0.85 (0.71 to 1.02)	–3% (–6 to 0%)	25%	96% (23%; 0%)
Pessimistic	The intervention is believed to be harmful, but there are few data, and an eventual benefit cannot be ruled out.	0.87 (0.72 to 1.04)	–3% (–5 to 0%)	30%	93% (18%; 0%)
Optimistic	The intervention is believed to be good, but a non-negligible chance of harm must be acknowledged.	0.86 (0.71 to 1.02)	–3% (–6 to 0%)	25%	96% (25%; 0%)

The region of practical equivalence was defined by an odds ratio interval of 0.9 and 1.1.

*This risk reduction is a simple estimation derived from the posterior draws of the predictor transformed by the inverse-link function with only the intervention as previously described.⁹




Fig. 1. Prior and posterior distributions for the high positive end-expiratory pressure and recruitment maneuver intervention are reported in the odds ratio (OR) scale. *Blue lines*, prior distribution; *red lines*, posterior distribution of the effect.

and discovered that approximately 25% of the posterior probability density lies within this interval. A similar density is observed for a significant benefit (odds ratio less than 0.8), whereas no density is found above an odds ratio of 1.25. Ultimately, the essence of Bayesian analysis lies in the full posterior distribution, akin to a map illustrating probabilities across various parameter values. Making decisions

based on different parameters or combinations thereof using regions of practical equivalence occurs after evaluating this primary information. In addition, we should try to avoid making strict yes or no decisions, because we might ignore how important or uncertain a parameter value really is. This type of thinking can lead to misunderstandings and confusion.^{31,32}




Fig. 2. Posterior distributions of the effects of high positive end-expiratory pressure and recruitment maneuver on the probability of postoperative pulmonary complication occurrence with a highlighted probability of benefit area. *Blue area*, probability density for the benefit threshold (odds ratio, less than 1); *orange area*, probability density for the harm threshold (odds ratio, greater than 1); *dark orange area*, probability density for the harm threshold (odds ratio, greater than 1.25); *black vertical lines*, region of practical equivalence boundaries (odds ratio, 0.9 to 1.1). PPC, postoperative pulmonary complications; ROPE, region of practical equivalence.

Table 3. Subgroups and Sensitivity Analyses

Pairwise Comparison	Subgroup	Odds Ratio (95% Credible Interval)*	Region of Practical Equivalence Probability, %	Probability of Benefit or Harm, %		
				Odds Ratio < 1	Odds Ratio < 0.8	Odds Ratio > 1.25
Subgroup analyses						
High PEEP and RM versus low PEEP	Laparoscopic surgery	0.67 (0.50 to 0.87)	0	99	90	0
		0.67 (0.50 to 0.87)	0	99	90	0
	Open surgery	0.67 (0.50 to 0.85)	0	99	91	0
		1.01 (0.79 to 1.23)	66	48	2	3
		1.01 (0.79 to 1.27)	62	46	3	4
		0.98 (0.76 to 1.22)	62	56	4	2
High ARISCAT risk		0.80 (0.53 to 1.13)	22	88	50	1
		0.80 (0.53 to 1.13)	23	88	49	1
		0.79 (0.54 to 1.12)	21	89	52	1
Intermediate ARISCAT risk		0.90 (0.73 to 1.07)	46	87	13	1
		0.90 (0.74 to 1.10)	50	85	11	1
		0.88 (0.71 to 1.06)	37	91	18	1
Body mass index < 30 kg m ⁻¹		0.88 (0.67 to 1.13)	40	83	24	1
		0.89 (0.66 to 1.16)	42	79	21	1
		0.84 (0.62 to 1.08)	29	89	35	1
		0.84 (0.66 to 1.07)	27	91	34	1
Body mass index > 30 kg m ⁻¹		0.85 (0.65 to 1.06)	28	91	33	1
		0.84 (0.64 to 1.06)	26	91	38	1
		1.19 (0.73 to 1.78)	28	21	3	42
		1.20 (0.71 to 1.79)	27	21	3	42
Postoperative CPAP versus no postoperative CPAP	High PEEP and RM	1.22 (0.74 to 1.85)	25	18	3	46
		1.14 (0.79 to 1.57)	34	21	2	31
		1.15 (0.77 to 1.57)	34	22	2	32
		1.14 (0.79 to 1.58)	34	22	2	30
Individualized PEEP versus fixed high PEEP†	High PEEP and RM	1.14 (0.71 to 1.70)	32	27	4	34
		1.14 (0.72 to 1.71)	32	27	4	34
		1.14 (0.70 to 1.68)	32	26	2	34
Sensitivity analyses						
Severe postoperative pulmonary complications		0.84 (0.69 to 1.01)	21	96	30	0
		0.87 (0.70 to 1.06)	34	91	20	0
		0.86 (0.69 to 1.04)	28	93	26	0
Different heterogeneity prior		0.86 (0.71 to 1.01)	25	96	23	0
		0.87 (0.72 to 1.03)	33	93	18	0
		0.86 (0.70 to 1.01)	26	95	24	0

The region of practical equivalence is the probability distribution between 0.9 and 1.1 odds ratio.

*Results for the posterior means and posterior probabilities are reported according to the prior distribution from top to bottom: neutral, pessimistic, and optimistic. †This comparison is carried out in the cohort receiving high PEEP and RM because no patient in the low PEEP cohort received it.

ARISCAT, Assess Respiratory Risk in Surgical Patients in Catalonia; CPAP, continuous positive airway pressure; PEEP, positive end-expiratory pressure; RM, recruitment maneuver.

Furthermore, the effects of the intervention estimated conditional on a new random effect reflecting the uncertainty within and across subjects show how the estimate distributions widen even to contain some posterior probability of harm. Hence, high PEEP and recruitment maneuver should not be a one-size-fits-all solution, but when and how to apply them should be carefully chosen. A partial explanation of these results comes from subgroup analysis. For instance, the effect of high PEEP and recruitment maneuver is more beneficial in specific groups such as laparoscopic surgery and high-risk ARISCAT patients, whereas in obesity, there is only a small interaction effect.

In addition, lung injury is ultimately related to the amount of stress and strain applied to the parenchyma during the mechanical ventilation process,³³ and ventilation must be

tailored to minimize it. Thus, applying high PEEP, recruitment maneuvers, or both can be beneficial as long as they decrease lung strain. This could not always be the case because, on the one hand, they can reduce the strain dynamic component by applying the same tidal volume to a greater number of alveoli thanks to recruitment and the pressure to keep them open, but on the other, they can increase the static strain by administering more force to the lung scaffold.³⁴ Other parameters, such as driving pressure, have been proposed to guide ventilation during general anesthesia, although its physiologic rationale is still being debated and laid out,³⁵ and specific randomized clinical trials to evaluate its effect are ongoing.³⁶ Also, other more comprehensive measurements of the total force, *i.e.*, the mechanical power applied to the lung during ventilation, have been described. Results from *post hoc* analysis show that this




Fig. 3. Sensitivity analysis results for the posterior probability distributions for the effect of high positive end-expiratory pressure and recruitment maneuvers on the probability of postoperative pulmonary complications occurrence. Reported are the effect estimates conditionally on a new random effect for a new hypothetical scenario. *Blue lines*, different heterogeneity prior; *red lines*, only severe postoperative complications as the outcome; *orange lines*, main model.

can be a promising measurement,³⁷ but we are not aware of any study specifically designed to test whether this parameter can reduce postoperative pulmonary complications.

Several limitations of this study must be acknowledged. First, although our database was previously harmonized, a

non-negligible heterogeneity between studies cannot be ruled out. Although we introduced a random effect in the model to precisely account for potential between studies differences, the effect of high PEEP and recruitment maneuver may be different in particular clinical scenarios or patients with specific

preoperative medications that cannot be ascertained by this study. Second, we chose to focus on individual patient data from large randomized clinical trial, leaving out some small studies. We could have switched to aggregate data meta-analysis and potentially had some difference in the point estimates, although with the trade-off of losing data granularity. As in our previous frequentist analysis, we decided to concentrate on the binary composite main outcome. However, it is important to note that this type of outcome has limitations, such as increased heterogeneity. Moreover, because the studied intervention combined the use of recruitment maneuvers and high PEEP, we cannot isolate the individual role of each component, and the synergistic or antagonistic effect in specific clinical situations could not be investigated. In addition, data on dynamic and static strain, such as functional residual capacity estimation or stress index, were not collected in the original studies thus, we cannot elucidate when the PEEP effect changes from beneficial to injurious by adding only overdistension. Moreover, although we did not observe any effect from other types of intervention, such as postoperative CPAP or the individualized setting of PEEP according to the pulmonary compliance after the recruitment maneuver, these interventions were administered in only one of the three studies; thus, our results should be interpreted with caution. Finally, we could not carry out the prespecified subgroup analysis on patients who presented atelectasis before the intervention administration because no data could identify such a group reliably. This must be investigated in future studies because the application of PEEP, and recruitment maneuvers in particular, seem sensible if a significant amount of collapsed parenchyma is present.

In conclusion, intraoperative mechanical ventilation with high PEEP and recruitment maneuvers slightly reduces the risk of occurrence and adverse postoperative pulmonary events as compared to low PEEP. This effect is likely not generalizable and is more pronounced in patients undergoing laparoscopy or at high risk of complications.

Research Support

Support was provided solely from institutional and/or departmental sources.

Competing Interests

Dr. Zampieri received consulting fees from Baxter (Deerfield, Illinois). Dr. Gama de Abreu received consulting fees from Zoll (Chelmsford, Massachusetts), and Dräger Medical (Lubeck, Germany), and Ambu Inc. (Copenhagen, Denmark), and honoraria for lectures from Merck Sharpe Dohme (Readington, New Jersey). Dr. Neto received honorary from Dräger Medical for lectures and own stocks in Endpoint Health (Palo Alto, California).

Correspondence

Address correspondence to Dr. Mazzinari: Hospital Universitario y Politécnico la Fe, Avenida Fernando Abril Martorell 106, Valencia, Spain. gmazzinari@gmail.com

Supplemental Digital Content

Additional tables, figures, and R code, <https://links.lww.com/ALN/D637>

References

- Schultz MJ, Hemmes SNT, Neto AS, et al.: Epidemiology, practice of ventilation and outcome for patients at increased risk of postoperative pulmonary complications: LASVEGAS—An observational study in 29 countries. *Eur J Anaesthesiol* 2017; 34:492–507
- Futier E, Constantin JM, Paugam-Burtz C, et al.; IMPROVE Study Group: A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013; 369:428–37
- Lagier D, Zeng C, Fernandez-Bustamante A, Vidal Melo MF: Perioperative pulmonary atelectasis: Part II. Clinical implications. *ANESTHESIOLOGY* 2022; 136:206–36
- Campos NS, Bluth T, Hemmes SNT, et al.; REPEAT: Intraoperative positive end-expiratory pressure and postoperative pulmonary complications: A patient-level meta-analysis of three randomised clinical trials. *Br J Anaesth* 2022; 128:1040–51
- Wasserstein RL, Lazar N: The ASA's statement on p-values: Context, process, and purpose. *Am Stat* 2016; 70:129–33
- Yarnell CJ, Abrams D, Baldwin MR, et al.: Clinical trials in critical care: Can a Bayesian approach enhance clinical and scientific decision making? *Lancet Respir Med* 2021; 9:207–16
- Zampieri FG, Casey JD, Shankar-Hari M, Harrell FE, Harhay MO: Using Bayesian methods to augment the interpretation of critical care trials: An overview of theory and example reanalysis of the alveolar recruitment for acute respiratory distress syndrome trial. *Am J Respir Crit Care Med* 2021; 203:543–52
- Zampieri FG, Damiani LP, Bakker J, et al.: Effects of a resuscitation strategy targeting peripheral perfusion status versus serum lactate levels among patients with septic shock: A Bayesian reanalysis of the ANDROMEDA-SHOCK trial. *Am J Respir Crit Care Med* 2020; 201:423–9
- Goligher EC, Tomlinson G, Hajage D, et al.: Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome and posterior probability of mortality benefit in a post hoc Bayesian analysis of a randomized clinical trial. *JAMA* 2018; 320:2251–9
- Komorowski M, Lemyze M: Informing future intensive care trials with machine learning. *Br J Anaesth* 2019; 123:14–6
- Mazzinari G, Zampieri FG, Ball L, et al.; PROVHILO investigators: Effect of intraoperative PEEP with recruitment maneuvers on the occurrence of postoperative pulmonary complications during general

- anesthesia—Protocol for Bayesian analysis of three randomized clinical trials of intraoperative ventilation. *F1000Res* 2022; 11:1090
12. Campos NS, Bluth T, Hemmes SNT, et al.: Re-evaluation of the effects of high PEEP with recruitment manoeuvres versus low PEEP without recruitment manoeuvres during general anaesthesia for surgery—Protocol and statistical analysis plan for an individual patient data meta-analysis of PROVHILO, iPROVE and PROBESE. *Rev Esp Anesthesiol Reanim (Engl Ed)* 2020; 67:76–89
 13. Schultz MJ, Hemmes SNT, de Abreu MG, et al.: High versus low positive end-expiratory pressure during general anaesthesia for open abdominal surgery (PROVHILO trial): A multicentre randomised controlled trial. *Lancet* 2014; 384:495–503
 14. Ferrando C, Soro M, Unzueta C, et al.: Individualized Perioperative Open-lung VEntilation (iPROVE) Network: Individualised perioperative open-lung approach versus standard protective ventilation in abdominal surgery (iPROVE): A randomised controlled trial. *Lancet Respir Med* 2018; 6:193–203
 15. Bluth T, Serpa Neto A, Schultz MJ, et al.: Effect of intraoperative high positive end-expiratory pressure (PEEP) with recruitment maneuvers vs. low PEEP on postoperative pulmonary complications in obese patients: A randomized clinical trial. *JAMA* 2019; 321:2292–305
 16. Canet J, Gallart L, Gomar C, et al.; ARISCAT Group: Prediction of postoperative pulmonary complications in a population-based surgical cohort. *ANESTHESIOLOGY* 2010; 113:1338–50
 17. Bürkner PC: brms: An R package for Bayesian multi-level models using Stan. *J Stat Softw* 2017; 80:1–28
 18. Röver C, Bender R, Dias S, et al.: On weakly informative prior distributions for the heterogeneity parameter in Bayesian random-effects meta-analysis. *Res Synth Methods* 2021; 12:448–74
 19. McElreath R: *Statistical Rethinking: A Bayesian Course with Examples in R and Stan*, 2nd edition. Boca Raton, Chapman & Hall/CRC Texts, 2020.
 20. Daniels MJ: A prior for the variance in hierarchical models. *Can J Stat* 1999; 27:567–78
 21. Makowski D, Ben-Shachar MS, Lüdtke D: bayestest R: Describing effects and their uncertainty, existence and significance within the bayesian framework. *J Open Source Softw* 2019; 4:1541
 22. Lenth R. emmeans: Estimated marginal means, aka least-squares means. Last R package version 1.10.0. 2024. Available at: <https://CRAN.R-project.org/package=emmeans>. Accessed June 5, 2024.
 23. Kallioinen N, Paananen T, Burkner P-C, Vehtari A: Detecting and diagnosing prior and likelihood sensitivity with power-scaling. *Stat Comput* 2024; 34:57
 24. Infranger D, Schmidt-Truckass A: P value functions: An underused method to present research results and to promote quantitative reasoning. *Stat Med* 2019; 20:4189–97
 25. Greenland S: Valid p-values behave exactly as they should: Some misleading criticisms of p-values and their resolution with s-values. *Am Stat* 2019; 73:106–14
 26. Friede T, Röver C, Wandel S, Neuenschwander B: Meta-analysis of two studies in the presence of heterogeneity with applications in rare diseases. *Biom J* 2017; 59:658–71
 27. Rhodes KM, Turner RM, White IR, Jackson D, Spiegelhalter DJ, Higgins JPT: Implementing informative priors for heterogeneity in meta-analysis using meta-regression and pseudo data. *Stat Med* 2016; 35:5495–511
 28. Cornell JE, Mulrow CD, Localio R, et al.: Random-effects meta-analysis of inconsistent effects: A time for change. *Ann Intern Med* 2014; 160:267–70
 29. Bigatello L, Pesenti A: Respiratory physiology for the anesthesiologist. *ANESTHESIOLOGY* 2019; 130:1064–77
 30. Zorrilla-Vaca A, Grant MC, Urman RD, Frenzl G: Individualised positive end-expiratory pressure in abdominal surgery: A systematic review and meta-analysis. *Br J Anaesth* 2022; 129:815–25
 31. Kruschke JK: Bayesian analysis reporting guidelines. *Nat Hum Behav* 2021; 5:1282–91
 32. Kruschke JK, Liddell TM: The Bayesian new statistics: Hypothesis testing, estimation, meta-analysis, and power analysis from a Bayesian perspective. *Psychon Bull Rev* 2018; 25:178–206
 33. Chiumello D, Carlesso E, Cadringer P, et al.: Lung stress and strain during mechanical ventilation for acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2008; 178:346–55
 34. Protti A, Andreis DT, Monti M, et al.: Lung stress and strain during mechanical ventilation: Any difference between statics and dynamics? *Crit Care Med* 2013; 41:1046–55
 35. Pelosi P, Ball L: Should we titrate ventilation based on driving pressure? Maybe not in the way we would expect. *Ann Transl Med* 2018; 6:389–389
 36. Hol L, Nijbroek SGLH, Neto AS, et al.: Driving Pressure during General Anesthesia for Open Abdominal Surgery (DESIGNATION): Study protocol of a randomized clinical trial. *Trials* 2020; 21:198
 37. Karalapillai D, Weinberg L, Neto AS, et al.: Intraoperative ventilator mechanical power as a predictor of postoperative pulmonary complications in surgical patients: A secondary analysis of a randomised clinical trial. *Eur J Anaesthesiol* 2022; 39:67–74

Appendix: Complete List of Collaborators

All of the collaborators listed below collected data and provided care for study patients.

Reevaluation of PEEP in Anesthesia Meta-analysis (REPEAT)

Writing Committee.

Niklas S. Campos: Department of Critical Care Medicine, Hospital Israelita Albert Einstein, São Paulo, Brazil; and Cardio-Pulmonary Department, Pulmonary Division, Instituto do Coração, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil.

Thomas Bluth: Pulmonary Engineering Group, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Sabrine N.T. Hemmes: Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology and the Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Carlos Ferrando: Department of Anesthesiology and Critical Care, Hospital Clinic de Barcelona, Barcelona, Spain; and CIBER of Respiratory Disease, Instituto de Salud Carlos III, Madrid, Spain.

Paolo Pelosi: IRCCS San Martino Policlinico Hospital, Genoa, Italy; and Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy.

Marcelo Gama de Abreu: Cardio-Pulmonary Department, Pulmonary Division, Instituto do Coração, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil.

Marcus J. Schultz: Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands; and Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand.

Ary Serpa Neto: Department of Critical Care Medicine, Hospital Israelita Albert Einstein, São Paulo, Brazil; Cardio-Pulmonary Department, Pulmonary Division, Instituto do Coração, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil; and Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Steering Committee.

Niklas S. Campos: Department of Critical Care Medicine, Hospital Israelita Albert Einstein, São Paulo, Brazil; and Cardio-Pulmonary Department, Pulmonary Division, Instituto do Coração, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil.

Thomas Bluth: Pulmonary Engineering Group, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Sabrine N.T. Hemmes: Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology and Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Julian Librero: Navarrabiomed-Fundación Miguel Servet, Red de Investigación en Servicios de Salud en Enfermedades Crónicas (REDISSEC), Pamplona, Spain.

Natividad Pozo: INCLIVA, Biomedical Research Institute, Valencia, Spain.

Carlos Ferrando: Department of Anesthesiology and Critical Care, Hospital Clinic de Barcelona, Barcelona, Spain; and CIBER of Respiratory Disease, Instituto de Salud Carlos III, Madrid, Spain.

Lorenzo Ball: IRCCS San Martino Policlinico Hospital, Genoa, Italy; and Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy.

Guido Mazzinari: Department of Anesthesiology and Pain Medicine, Hospital de Manises, Valencia, Spain; and Research Group in Perioperative Medicine, Hospital Universitario y Politécnico la Fe, Valencia, Spain.

Paolo Pelosi: IRCCS San Martino Policlinico Hospital, Genoa, Italy; and Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy.

Marcelo Gama de Abreu: Pulmonary Engineering Group, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Marcus J. Schultz: Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands; and Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand.

Ary Serpa Neto: Department of Critical Care Medicine, Hospital Israelita Albert Einstein, São Paulo, Brazil; Cardio-Pulmonary Department, Pulmonary Division, Instituto do Coração, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil; and Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Protective Ventilation using High versus Low Positive End-expiratory Pressure Trial (PROVHIL0) Investigators

Steering and Executive Committees.

Sabrine N.T. Hemmes: Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Marcelo Gama de Abreu: Pulmonary Engineering Group, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Paolo Severgnini: Department of Biotechnologies and Sciences of Life, University of Insubria, Varese, Italy.

Markus W. Hollmann: Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Jan M. Binnekade: Department of Intensive Care, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Hermann Wrigge: Department of Anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany.

Jaume Canet: Department of Anesthesiology, Hospital Universitari Germans Trias i Pujol, Barcelona, Spain.

Michael Hiesmayr: Division of Cardiac, Thoracic, and Vascular Anesthesia and Intensive Care, Medical University Vienna, Vienna, Austria.

Werner Schmid: Division of Cardiac, Thoracic, and Vascular Anesthesia and Intensive Care, Medical University Vienna, Vienna, Austria.

Edda Tschernko: Division of Cardiac, Thoracic, and Vascular Anesthesia and Intensive Care, Medical University Vienna, Vienna, Austria.

Samir Jaber: Department of Critical Care Medicine and Anesthesiology (SAR B), Saint Eloi University Hospital, Montpellier, France.

Göran Hedenstierna: Department of Medical Sciences, Clinical Physiology, Uppsala University, Uppsala, Sweden.

Christian Putensen: Department of Anesthesiology and Intensive Care Medicine, University Hospital Bonn, Bonn, Germany.

Paolo Pelosi: IRCCS San Martino Policlinico Hospital, Genoa, Italy; and Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy.

Marcus J. Schultz: Department of Intensive Care and Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands; and Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand.

Investigators. Asterisks (*) indicate local principal investigators. The names are listed in alphabetical order.

Agnes Marti: Hospital Universitari Germans Trias i Pujol, Barcelona, Spain.

Alessandro Bacuzzi: University of Insubria, Azienda Ospedaliera Fondazione Macchi, Ospedale di Circolo, Varese, Italy.

Alexander Brodhun: Johannes Gutenberg, Universität Mainz, Mainz, Germany.

Alexandre Molin*: Università degli Studi di Genova, IRCCS San Martino IST, Genoa, Italy.

Alfred Merten: Hospital Sant Pau, Barcelona, Spain.

Ana Parera: Hospital Sant Pau, Barcelona, Spain.

Andrea Brunelli*: Hospital Universitari Germans Trias i Pujol, Barcelona, Spain.

Andrea Cortegiani: Università degli Studi di Palermo, Palermo, Italy.

Andreas Güldner: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.

Andreas W. Reske: University of Leipzig, Leipzig, Germany.

Angelo Gratarola: Università degli Studi di Genova, IRCCS San Martino IST, Genoa, Italy.

Antonino Giarratano*: Università degli Studi di Palermo, Palermo, Italy.

Bea Bastin: Düsseldorf University Hospital, Heinrich-Heine University Düsseldorf, Düsseldorf, Germany.

Bjorn Heyse: Ghent University Hospital, Ghent, Belgium.

Branka Mazul-Sunko*: University Hospital Sveti Duh, Zagreb, Croatia.

Bruno Amantea: University “Magna Graecia” of Catanzaro, Catanzaro, Italy.

Bruno Barberis: Azienda Sanitaria Locale TO3, Ospedale di Rivoli, Rivoli, Italy.

Christian Putensen*: University Hospital of Bonn Medical School, Bonn, Germany.

Christopher Uhlig: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.

Conrado Minguez Marín: Consorcio Hospital General Universitario Valencia, Valencia, Spain.

Cristian Celentano: Azienda Sanitaria Locale TO3, Ospedale di Rivoli, Rivoli, Italy.

Daniela La Bella: University of Foggia, Foggia, Italy.

David D’Antini: University of Foggia, Foggia, Italy.

David Velghe*: ZNA Middelheim, Antwerp, Belgium.

Demet Sulemanji: Massachusetts General Hospital, Boston, Massachusetts.

Edoardo De Robertis*: University of Napoli Federico II, Napoli, Italy.

Eric Hartmann: Johannes Gutenberg, Universität Mainz, Mainz, Germany.

Francesca Montalto: Università degli Studi di Palermo, Palermo, Italy.

Francesco Tropea: University “Magna Graecia” of Catanzaro, Catanzaro, Italy.

Gary H. Mills*: Sheffield Teaching Hospitals, Sheffield, United Kingdom.

Gilda Cinnella*: University of Foggia, Foggia, Italy.

Giorgio Della Rocca*: Università degli Studi di Udine, Udine, Italy.

Girolamo Caggianelli: University of Foggia, Foggia, Italy.

Giulia Pellerano: Università degli Studi di Genova, IRCCS San Martino IST, Genoa, Italy.

- Giuseppina Mollica: University of Foggia, Foggia, Italy.
 Guillermo Buggedo*: Hospital Clínico de la Pontificia Universidad Católica de Chile, Santiago, Chile.
 Hermann Wrigge*: University of Leipzig, Leipzig, Germany.
 Jan-Paul Mulier*: AZ St Jan, Brugge, Belgium.
 Jeroen Vandenbrande: Virga Jesse Ziekenhuis, Hasselt, Belgium.
 Johann Geib: Düsseldorf University Hospital, Heinrich-Heine University Düsseldorf, Düsseldorf, Germany.
 Jonathan Yaqub: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.
 Jorge Florez: Hospital Clínico de la Pontificia Universidad Católica de Chile, Santiago, Chile.
 Juan F. Mayoral: Fundación Puigvert, Barcelona, Spain.
 Juraj Sprung*: Mayo Clinic, Rochester, Minnesota.
 Jurgen Van Limmen: Ghent University Hospital, Ghent, Belgium.
 Lieuwe D.J. Bos: Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.
 Luc de Baerdemaeker: Ghent University Hospital, Ghent, Belgium.
 Luc Jamaer*: Virga Jesse Ziekenhuis, Hasselt, Belgium.
 Luigi Spagnolo*: Azienda Sanitaria Locale TO3, Ospedale di Rivoli, Rivoli, Italy.
 Lydia Stry: Johannes Gutenberg, Universität Mainz, Mainz, Germany.
 Manuel Granell Gil*: Consorcio Hospital General Universitario Valencia, Valencia, Spain.
 Marcelo Gama de Abreu*: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.
 Marcos F. Vidal Melo*: Massachusetts General Hospital, Boston, Massachusetts.
 Marcus J. Schultz*: Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.
 Maria Carmen Unzueta*: Hospital Sant Pau, Barcelona, Spain.
 Maria Victoria Moral: Hospital Sant Pau, Barcelona, Spain.
 Marion Ferner: Johannes Gutenberg, Universität Mainz, Mainz, Germany.
 Markus W. Hollmann: Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.
 Martin Weiss: Düsseldorf University Hospital, Heinrich-Heine University Düsseldorf, Düsseldorf, Germany.
 Massimo Vanoni: University of Insubria, Azienda Ospedaliera Fondazione Macchi, Ospedale di Circolo, Varese, Italy.
 Maximilian S. Schaefer: Düsseldorf University Hospital, Heinrich-Heine University Düsseldorf, Düsseldorf, Germany.
 Mercè Prieto: Fundación Puigvert, Barcelona, Spain.
 Michele Griò: Azienda Sanitaria Locale TO3, Ospedale di Rivoli, Rivoli, Italy.
 Paolo Severgnini*: University of Insubria, Azienda Ospedaliera Fondazione Macchi, Ospedale di Circolo, Varese, Italy.
 Peter Markus Spieth: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.
 Philipp Simon: University of Leipzig, Leipzig, Germany.
 Phoebe Bodger*: Barts Health National Health Service Trust, London, United Kingdom.
 Pilar Sierra: Fundación Puigvert, Barcelona, Spain.
 Rita Laufenberg-Feldmann*: Johannes Gutenberg, Universität Mainz, Mainz, Germany.
 Roberta Rusca: Università degli Studi di Genova, IRCCS San Martino IST, Genoa, Italy.
 Rodolfo Proietti*: Università degli Studi di Roma Cattolica, Rome, Italy.
 Sabrine N.T. Hemmes: Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.
 Santi Maurizio Raineri: Università degli Studi di Palermo, Palermo, Italy.
 Santo Caroleo*: University “Magna Graecia” of Catanzaro, Catanzaro, Italy.
 Sergi Sabaté*: Fundación Puigvert, Barcelona, Spain.
 Stefan De Hert*: Ghent University Hospital, Ghent, Belgium.
 Stefano Pezzato: Università degli Studi di Genova, IRCCS San Martino IST, Genoa, Italy.
 Tanja A. Treschan*: Düsseldorf University Hospital, Heinrich-Heine University Düsseldorf, Düsseldorf, Germany.
 Tatjana Goranovic: University Hospital Sveti Duh, Zagreb, Croatia.
 Thea Koch: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.
 Thomas Bluth: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.
 Thomas Kiss: University Hospital Dresden, Technische Universität Dresden, Dresden, Germany.
 Valter Perilli: Università degli Studi di Roma Cattolica, Rome, Italy.
 Virginia Cegarra: Hospital Sant Pau, Barcelona, Spain.
 Werner Schmid*: Medical University Vienna, Vienna.
- Information and Funding.** The PROVHILO trial is a collaboration of the Protective Ventilation Network (PROVENet). The trial was funded by the European Society of Anesthesiology and the Academic Medical Center (Amsterdam, The Netherlands).
- Individualized Perioperative Open-lung Ventilation Trial (iPROVE) Investigators**
- Steering and Executive Committees.**
 Carlos Ferrando: Department of Anesthesiology and Critical Care, Hospital Clinic de Barcelona, Barcelona, Spain; and CIBER of Respiratory Disease, Instituto de Salud Carlos III, Madrid, Spain.
 Javier Belda: Department of Surgery, Medicine Faculty, Universidad de Valencia, Valencia, Spain.

Marina Soro: Department of Anesthesiology and Critical Care, Hospital Clínico Universitario de Valencia, Valencia, Spain.

Jaume Canet: Department of Anesthesiology and Critical Care, Hospital Universitario Germans Trias i Pujol, Barcelona, Spain.

Carmen Unzueta: Department of Anesthesiology and Critical Care, Hospital Universitario San Pau, Barcelona, Spain.

Fernando Suarez-Sipmann: Department of Anesthesiology and Critical Care, Hospital Universitario La Princesa, Madrid, Spain.

Julián Librero: Navarrabiomed-Fundación Miguel Servet, Red de Investigación en Servicios de Salud en Enfermedades Crónicas (REDISSEC), Pamplona, Spain.

Alicia Llombart: Department of Pharmacology, Corchan Hospital, Barcelona, Spain.

Lucas Rovira: Department of Anesthesiology and Critical Care, Hospital General Universitario de Valencia, Valencia, Spain.

Manuel Granell: Department of Anesthesiology and Critical Care, Hospital General Universitario de Valencia, Valencia, Spain.

César Aldecoa: Department of Anesthesiology and Critical Care, Hospital Universitario Río Hortega, Valladolid, Spain.

Oscar Diaz-Cambronero: Department of Anesthesiology and Critical Care, Hospital Universitario y Politécnico La Fe, Valencia, Spain.

Jaume Balust: Department of Anesthesiology and Critical Care, Hospital Clínic i Provincial Universitario de Barcelona, Barcelona, Spain.

Ignacio Garutti: Department of Anesthesiology and Critical Care, Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Rafael Gonzalez: Department of Anesthesiology and Critical Care, Hospital de León, León, Spain.

Lucía Gallego: Department of Anesthesiology and Critical Care, Hospital Universitario Miguel Servet de Zaragoza, Zaragoza, Spain.

Santiago Garcia del Valle: Department of Anesthesiology and Critical Care, Hospital Fundación de Alcorcón, Alcorcón, Spain.

Javier Redondo: Department of Anesthesiology and Critical Care, Hospital General Universitario de Ciudad Real, Ciudad Real, Spain.

David Pestaña: Department of Anesthesiology and Critical Care, Hospital Universitario Ramón y Cajal, Madrid, Spain.

Aurelio Rodríguez: Department of Anesthesiology and Critical Care, Hospital de Gran Canaria, Dr. Negrín, Gran Canaria, Spain.

Javier García: Department of Anesthesiology and Critical Care, Hospital Universitario Puerta de Hierro de Majadahonda, Madrid, Spain.

Manuel de la Matta: Department of Anesthesiology and Critical Care, Hospital Universitario Virgen del Rocío de Sevilla, Sevilla, Spain.

Maite Ibáñez: Department of Anesthesiology, Hospital de la Marina Baixa de la Vila Joiosa, Alicante, Spain.

Francisco Barrios: Department of Anesthesiology and Critical Care, Hospital Principe de Asturias, Madrid, Spain.

Samuel Hernández: Department of Anesthesiology, Hospital NS de Candelaria, Santa Cruz de Tenerife, Spain.

Vicente Torres: Department of Anesthesiology and Critical Care, Hospital Son Espases de Mallorca, Mallorca, Spain.

Salvador Peiró: IISLAFE Clinical Research Institute, Hospital Universitario y Politécnico La Fe, Valencia, Spain.

Natividad Pozo: INCLIVA, Biomedical Research Institute, Valencia, Spain.

Investigators. The names are listed in alphabetical order.

Abigail Villena: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Albert Carramiñana: Hospital Universitario Germans Trias i Pujol, Barcelona, Spain.

Alberto Gallego-Casilda: Hospital Universitario Miguel Servet de Zaragoza, Zaragoza, Spain.

Alejandro Duca: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Amalia Alcón: Hospital Clínic i Provincial Universitario de Barcelona, Barcelona, Spain.

Amanda Miñana: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Ana Asensio: Hospital Universitario Miguel Servet de Zaragoza, Zaragoza, Spain.

Ana Colás: Hospital Universitario Miguel Servet de Zaragoza, Zaragoza, Spain.

Ana Isabel Galve: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Ana Izquierdo: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Ana Jurado: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Ana María Pérez: Hospital de León, León, Spain.

Ana Mugarra: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Ana Parera: Hospital Universitario San Pau, Barcelona, Spain.

Andrea Brunelli: Hospital Universitario Germans Trias i Pujol, Barcelona, Spain.

Andrea Gutierrez: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Ángeles De Miguel: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Angels Lozano: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Antonio Katime: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Antonio Romero: Hospital Universitario Puerta de Hierro de Majadahonda, Madrid, Spain.

Beatriz Garrigues: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Begoña Ayas: Hospital Universitario y Politécnico La Fe, Valencia, Spain.

Blanca Arocas: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Carlos Delgado: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Carmen Fernández: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Carolina Romero: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Clara Gallego: Hospital Universitario Ramón y Cajal, Madrid, Spain.

Cristina Garcés: Hospital Universitario Miguel Servet de Zaragoza, Zaragoza, Spain.

Cristina Lisbona: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Cristina Parrilla: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Daniel López-Herrera: Hospital Universitario Virgen del Rocío de Sevilla, Sevilla, Spain.

Domingo González: Hospital Universitario Virgen del Rocío de Sevilla, Sevilla, Spain.

Eduardo Llamazares: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Elena Del Rio: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Elena Lozano: Hospital de la Marina Baja de la Vila Joiosa, Alicante, Spain.

Ernesto Pastor: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Estefanía Chamorro: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Estefanía Gracia: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Ester Sánchez: Hospital de la Marina Baja de la Vila Joiosa, Alicante, Spain.

Esther Romero: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Fernando Díez: Hospital de León, León, Spain.

Ferran Serralta: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Francisco Daviu: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Francisco Sandín: Hospital Universitario Miguel Servet de Zaragoza, Zaragoza, Spain.

Gerardo Aguilar: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Gerardo Tusman: Hospital Privado de Comunidad, Mar del Plata, Buenos Aires, Argentina.

Gonzalo Azparren: Hospital Universitario San Pau, Barcelona, Spain.

Graciela Martínez-Pallí: Hospital Clínic i Provincial Universitario de Barcelona, Barcelona, Spain.

Guido Mazzinari: Hospital de Manises, Valencia, Spain.

Inmaculada Benítez: Hospital Universitario Virgen del Rocío de Sevilla, Sevilla, Spain.

Inmaculada Hernández: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Inmaculada India: Hospital Universitario San Pau, Barcelona, Spain.

Irene León: Hospital Clínic i Provincial Universitario de Barcelona, Barcelona, Spain.

Isabel Fuentes: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Isabel Ruiz: Hospital Universitario Ramón y Cajal, Madrid, Spain.

Jaume Puig: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Javie Ignacio Román: Hospital Son Espases de Mallorca, Mallorca, Spain.

Jesús Acosta: Hospital Universitario Virgen del Rocío de Sevilla, Sevilla, Spain.

Jesús Rico-Feijoo: Hospital Universitario Río Hortega, Valladolid, Spain.

Jonathan Olmedo: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Jose A. Carbonell: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Jose M. Alonso: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Jose María Pérez: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Jose Miguel Marcos: Hospital de León, León, Spain.

Jose Navarro: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Jose Valdivia: Hospital de la Marina Baja de la Vila Joiosa, Alicante, Spain.

Juan Carrizo: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Laura Piqueras: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Laura Soriano: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Laura Vaquero: Hospital Universitario Río Hortega, Valladolid, Spain.

Lisset Miguel: Hospital Universitario Ramón y Cajal, Madrid, Spain.

Lorena Muñoz: Hospital General Universitario de Valencia, Valencia, Spain.

Lucía Valencia: Hospital de Gran Canaria, Dr. Negrín, Gran Canaria, Spain.

Luis Olmedilla: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Ma Justina Etulain: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Manuel Tisner: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

María Barrio: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

María Dolores Alonso: Hospital General Universitario de Valencia, Valencia, Spain.

María García: Hospital Universitario Río Hortega, Valladolid, Spain.

María J. Hernández: Hospital General Universitario de Valencia, Valencia, Spain.

María José Alberola: Hospital Universitario y Politécnico La Fe, Valencia, Spain.

María Parra: Hospital Clínico Universitario de Valencia, Valencia, Spain.

María Pilar Argente: Hospital Universitario y Politécnico La Fe, Valencia, Spain.

María Vila: Hospital Universitario y Politécnico La Fe, Valencia, Spain.

Mario De Fez: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Marta Agilaga: Hospital Universitario San Pau, Barcelona, Spain.

Marta Gine: Hospital Universitario San Pau, Barcelona, Spain.

Mercedes Ayuso: Hospital Príncipe de Asturias, Madrid, Spain.

Mercedes García: Hospital Universitario San Pau, Barcelona, Spain.

Natalia Bejarano: Hospital General Universitario de Ciudad Real, Ciudad Real, Spain.

Natalia Peña: Hospital Universitario Miguel Servet de Zaragoza, Zaragoza, Spain.

Nazario Ojeda: Hospital de Gran Canaria, Dr. Negrín, Gran Canaria, Spain.

Nilda Martínez: Hospital Universitario Ramón y Cajal, Madrid, Spain.

Nuria García: Hospital Universitario y Politécnico La Fe, Valencia, Spain.

Oto Padrón: Hospital de Gran Canaria, Dr. Negrín, Gran Canaria, Spain.

Pablo García: Hospital Fundación de Alcorcón, Alcorcón, Spain.

Paola Valls: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Patricia Cruz: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Patricia Piñeiro: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Pedro Charco: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Rafael Anaya: Hospital Universitario San Pau, Barcelona, Spain.

Ramiro López: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Rayco Rodríguez: Hospital de Gran Canaria, Dr. Negrín, Gran Canaria, Spain.

Rocío Martínez: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Roger Pujol: Hospital Clínic i Provincial Universitario de Barcelona, Barcelona, Spain.

Rosa Dosdá: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Rosa Lardies: Hospital Clínic i Provincial Universitario de Barcelona, Barcelona, Spain.

Ruben Díaz: Hospital Universitario San Pau, Barcelona, Spain.

Rubén Villazala: Hospital General Universitario de Ciudad Real, Ciudad Real, Spain.

Sara Zapatero: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Sergio Cabrera: Hospital de Gran Canaria, Dr. Negrín, Gran Canaria, Spain.

Sergio Sánchez: Hospital General Universitario de Ciudad Real, Ciudad Real, Spain.

Silvia Martín: Hospital Universitario Río Hortega, Valladolid, Spain.

Suzana Diaz: Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Tania Franco: Hospital Universitario Ramón y Cajal, Madrid, Spain.

Tania Moreno: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Tania Socorro: Hospital Clínico Universitario de Valencia, Valencia, Spain.

Vicente Gilabert: Hospital de la Marina Baixa de la Vila Joiosa, Alicante, Spain.

Victor Baladrón: Hospital General Universitario de Ciudad Real, Ciudad Real, Spain.

Victoria Moral: Hospital Universitario San Pau, Barcelona, Spain.

Virgina Cegarra: Hospital Universitario San Pau, Barcelona, Spain.

Viviana Varón: Hospital Fundación de Alcorcón, Alcorcón, Spain.

Information and Funding. The iPROVE trial wants to thank all staff who participated, including researchers, nurses, emergency medical technicians, and anyone who has made possible the successful completion of the project. The trial investigators also thank the technical support provided by the Biomedical Research Foundation Institute. This trial was funded by the Instituto de Salud Carlos III of the Spanish Ministry of Economy and Competitiveness (grant No. PI14/00829, co-financed by the European Regional Development Fund) and received the support of the grants program of the European Society of Anesthesiology.

Protective Intraoperative Ventilation with Higher versus Lower Levels of Positive End-Expiratory Pressure in OBESE Patients' Trial (PROBESE) Investigators

Writing, Steering, and Executive Committees.

Thomas Bluth: Pulmonary Engineering Group, Department of Anesthesiology and Intensive Care Medicine, University

Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Ary Serpa Neto: Department of Critical Care Medicine, Hospital Israelita Albert Einstein, São Paulo, Brazil; Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands; and Cardio-Pulmonary Department, Pulmonary Division, Instituto do Coração, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil.

Ilona Bobek: Aneszteziológiai és Intenzív Terápiás Klinika, Semmelweis Egyetem, Budapest, Hungary.

Jaume Canet: Department of Anesthesiology, Hospital Universitari Germans Trias i Pujol, Barcelona, Spain.

Gilda Cinnella: Department of Anesthesiology and Intensive Care Medicine, University of Foggia, Foggia, Italy.

Luc de Baerdemaeker: Department of Anesthesiology and Perioperative Medicine, Ghent University Hospital, Ghent, Belgium.

Cesare Gregoretti: Department of Anesthesiology, Città della Salute e della Scienza, Turin, Italy.

Göran Hedenstierna: Department of Medical Sciences, Clinical Physiology, Uppsala University, Uppsala, Sweden.

Sabrina N.T. Hemmes: Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Michael Hiesmayr: Division of Cardiac, Thoracic, and Vascular Anesthesia and Intensive Care, Medical University Vienna, Vienna, Austria.

Markus W. Hollmann: Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Samir Jaber: Department of Critical Care Medicine and Anesthesiology (SAR B), Saint Eloi University Hospital, Montpellier, France.

John Laffey: Department of Anesthesiology, St. Michael's Hospital, University of Toronto, Toronto, Ontario, Canada.

Marc-Joseph Licker: Division of Anesthesiology, University Hospitals of Geneva, Geneva, Switzerland.

Klaus Markstaller: Department of Anesthesia, Critical Care and Pain Medicine, Medical University Vienna, Vienna, Austria.

Idit Matot: Department of Anesthesia, Pain and Critical Care, Tel-Aviv Medical Center, Sackler Medical School, Tel Aviv, Israel.

Gary H. Mills: Operating Services, Critical Care and Anesthesia, Sheffield Teaching Hospitals, University of Sheffield, Sheffield, United Kingdom.

Jan Paul Mulier: Department of Anesthesiology, AZ Sint Jan Brugge-Oostende AV, Brugge, Belgium.

Christian Putensen: Department of Anesthesiology and Intensive Care Medicine, University Hospital Bonn, Bonn, Germany.

Rolf Rossaint: Department of Anesthesiology, Medical Faculty, Rheinisch-Westfälische Technische Hochschule Aachen University, Aachen, Germany.

Jochen Schmitt: Center for Evidence-based Healthcare, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Mert Senturk: Department of Anesthesiology and Intensive Care, Istanbul University, Istanbul Faculty of Medicine, Istanbul, Turkey.

Paolo Severgnini: Dipartimento di Anestesia, Università dell'Insubria, Azienda Ospedaliera ASST Sette Laghi Ospedale di Circolo e Fondazione Macchi, Varese, Italy.

Juraj Sprung: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota.

Marcos F.Vidal Melo: Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, Massachusetts.

Hermann Wrigge: Department of Anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany; and Department of anesthesiology, Intensive Care and Emergency Medicine, Pain Therapy, Bergmannstrost Hospital Halle, Halle, Germany.

Marcus J. Schultz: Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands; and Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand.

Paolo Pelosi: IRCCS San Martino Policlinico Hospital, Genoa, Italy; and Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy.

Marcelo Gama de Abreu: Pulmonary Engineering Group, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Protective Ventilation (PROVE) Investigators

Asterisks (*) indicate local principal investigators. The names are listed in alphabetical order.

Abelha, Fernando*: Department of Anesthesiology, Centro Hospitalar de São João, Porto, Portugal; and Department of Surgery and Physiology, Cardiovascular Research Center, Faculty of Medicine of the University of Porto, Porto, Portugal.

Abitağaoğlu, Sühayla: Fatih Sultan Mehmet Educational and Research Hospital, İstanbul, Turkey.

Achilles, Marc*: Marienhospital Wesel, Wesel, Germany.
Adebesin, Afeez: Imperial College Healthcare National Health Service Trust, London, United Kingdom.

Adriaenssens, Ine: University Hospital Antwerp, Antwerp, Belgium.

Ahene, Charles*: Cleveland Clinic Abu Dhabi, Al Maryah Island, Abu Dhabi, United Arab Emirates.

Akbar, Fatima: Imperial College Healthcare National Health Service Trust, London, United Kingdom.

AlHarbi, Mohammed: Ministry of National Guard Health Affairs, King Abdulaziz Medical City, Riyadh,

Saudi Arabia; and Anesthesia Department, King Saud Bin Abdulaziz University for Health Science, Riyadh, Saudi Arabia.

Al Khoury al Kallab, Rita: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Albanel, Xavier: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Aldenkortt, Florence: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Alfouzan, Rawan Abdullah Saleh: Anesthesia Department, Ministry of National Guard Health Affairs, King Abdulaziz Medical City, Riyadh, Saudi Arabia.

Alruqaie, Reef: Anesthesia Department, Ministry of National Guard Health Affairs, King Abdulaziz Medical City, Riyadh, Saudi Arabia.

Altermatt, Fernando*: Department of Anesthesiology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile.

Araujo, Bruno Luís de Castro*: Department of Anesthesiology, Hospital do Câncer II, National Cancer Institute of Brazil, Rio de Janeiro, Brazil.

Arbesú, Genaro: University Hospital Germans Trias i Pujol, Institut Catalán de la Salut, Universidad Autónoma de Barcelona, Barcelona, Spain.

Artsi, Hanna: Division of Anesthesia, Pain and Critical Care, Tel-Aviv Medical Center affiliated with Sackler Medical School, Tel Aviv University, Tel Aviv, Israel.

Aurilio, Caterina*: Department of Women, Child and General and Specialized Surgery, University of Campania "L. Vanvitelli," Naples, Italy.

Ayanoglu, Omer Hilmi: Marmara University Pendik Training and Research Hospital, Istanbul, Turkey.

Bacuzzi, Alessandro: Università dell'Insubria, Dipartimento di Anestesia, Azienda Ospedaliera Asst Settlaghi Ospedale di Circolo e Fondazione Macchi, Varese, Italy.

Baig, Harris*: University of Mississippi Medical Center, Jackson, Mississippi.

Baird, Yolanda: St. Richard's Hospital, Chichester, United Kingdom.

Balonov, Konstantin*: Department of Anesthesiology and Perioperative Medicine, Tufts Medical Center, Boston, Massachusetts.

Balust, Jaume: Department of Anesthesiology, Hospital Clínic, Barcelona, Spain.

Banks, Samantha: Royal Cornwall Hospital National Health Service Trust, Truro, United Kingdom.

Bao, Xiaodong: Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, Massachusetts.

Baumgartner, Mélanie: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Belda Tortosa, Isabel*: Hospital Sagrat Cor, Barcelona, Spain.

Bergamaschi, Alice: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Bergmann, Lars*: Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Universitätsklinikum Knappschaftskrankenhaus Bochum, Bochum, Germany.

Bigatello, Luca*: St. Elizabeth's Medical Center, Boston, Massachusetts.

Biosca Pérez, Elena: Consorcio Hospital General Universitario of Valencia, Spain.

Birr, Katja: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Bluth, Thomas: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Bojaxhi, Elird: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Jacksonville, Florida.

Bonenti, Chiara: Università dell'Insubria, Dipartimento di Anestesia, Azienda Ospedaliera Asst Settlaghi Ospedale di Circolo e Fondazione Macchi, Varese, Italy.

Bonney, Iwona: Department of Anesthesiology and Perioperative Medicine, Tufts Medical Center, Boston, Massachusetts.

Bos, Elke M.E.: Amsterdam University Medical Center, University of Amsterdam, Department of Anesthesiology, Amsterdam, The Netherlands.

Bowman, Sara: Homerton University Hospitals National Health Service Foundation Trust, London, United Kingdom.

Braz, Leandro Gobbo: Department of Anesthesiology, Universidade Estadual Paulista, Botucatu, Brazil.

Brugnoni, Elisa: Università dell'Insubria, Dipartimento di Anestesia, Azienda Ospedaliera Asst Settlaghi Ospedale di Circolo e Fondazione Macchi, Varese, Italy.

Brull, Sorin J.: Department of Anesthesiology and Perioperative Medicine, College of Medicine, Mayo Clinic, Jacksonville, Florida.

Brunetti, Iole: Anestesia e Terapia Intensiva, IRCCS Policlinico San Martino, Genova, Italy.

Bruni, Andrea: Anesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Graeca University, Catanzaro, Italy.

Buenvenida, Shonie L.: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota.

Busch, Cornelius Johannes*: Department of Anesthesiology, Heidelberg University Hospital, Heidelberg, Germany.

Camerini, Giovanni: Dipartimento di Scienze Chirurgiche e Diagnostiche Integrate, Università degli Studi di Genova, Genova, Italy; and Clinica Chirurgica 1, IRCCS Policlinico San Martino, Genova, Italy.

Canet, Jaume*: University Hospital Germans Trias i Pujol, Institut Català de la Salut, Universidad Autónoma de Barcelona, Barcelona, Spain.

Capatti, Beatrice: Department Morphology, Surgery and Experimental Medicine, Anesthesia and Intensive Care University Section, University of Ferrara, Italy.

Carmona, Javiera: Department of Anesthesiology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile.

Carungcong, Jaime: Magill Department of Anesthesia, Chelsea and Westminster National Health Service Foundation Trust, London, United Kingdom.

Carvalho, Marta: Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

Cattan, Anat: Division of Anesthesia, Pain and Critical Care, Tel-Aviv Medical Center affiliated with Sackler Medical School, Tel Aviv University, Tel Aviv, Israel.

Cavaleiro, Carla: Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal and Centro de Investigação Clínica em Anestesiologia, Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

Chiumello, Davide*: Dipartimento di Scienze della Salute, Università degli Studi di Milano, Milan, Italy; and SC Anestesia e Rianimazione, ASST Santi Paolo e Carlo, Centro Ricerca Coordinata Insufficienza Respiratoria, Milan, Italy.

Ciardo, Stefano: Department Morphology, Surgery and Experimental Medicine, Anesthesia and Intensive Care University Section, University of Ferrara, Ferrara, Italy.

Coburn, Mark: Department of Anesthesiology, Medical Faculty, Rheinisch-Westfälische Technische Hochschule Aachen University, Aachen, Germany.

Colella, Umberto: Department of Women, Child and General and Specialized Surgery, University of Campania “L. Vanvitelli,” Naples, Italy.

Contreras, Victor: Department of Anesthesiology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile.

Corman Dincer, Pelin*: Marmara University Pendik Training and Research Hospital, Istanbul, Turkey.

Cotter, Elizabeth: Section of Critical Care Medicine, Department of Anesthesia and Critical Care, University of Chicago, Chicago, Illinois.

Crovetto, Marcia: Department of Anesthesiology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile.

Darrah, William: Department of Anesthesiology, St. Michael's Hospital, University of Toronto, Toronto, Ontario, Canada.

Davies, Simon*: York Teaching Hospitals National Health Service Foundation Trust, York, United Kingdom.

de Baerdemaeker, Luc*: Ghent University Hospital, Ghent, Belgium.

De Hert, Stefan: Ghent University Hospital, Ghent, Belgium.

Del Cojo Peces, Enrique: Área de Salud Don Benito-Villanueva, Don Benito, Spain.

Delphin, Ellise: Montefiore Medical Center, Bronx, New York.

Diaper, John: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

do Nascimento, Jr., Paulo*: Department of Anesthesiology, Universidade Estadual Paulista, Botucatu, Brazil.

Donatiello, Valerio: Department of Women, Child and General and Specialized Surgery, University of Campania “L. Vanvitelli,” Naples, Italy.

Dong, Jing: Department of Anesthesiology, Fudan University Shanghai Cancer Center, Shanghai, China; and Department of Oncology, Shanghai Medical College, Fudan University, Shanghai, China.

Dourado, Maria do Socorro: Associação Hospitalar Beneficente São Vicente de Paulo, Passo Fundo, Brazil.

Dullenkopf, Alexander*: Frauenfeld Cantonal Hospital, Frauenfeld, Switzerland.

Ebner, Felix: Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Universitätsklinikum Knappschafts-Krankenhaus Bochum, Bochum, Germany.

Elgendy, Hamed*: Departments of Anesthesiology, King Abdullah Medical City, Makkah, Saudi Arabia and Assiut University, Assiut, Egypt and Hamad Medical Corporation, Qatar.

Ellenberger, Christoph: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Erdoğan Arı, Dilek*: Fatih Sultan Mehmet Educational and Research Hospital, İstanbul, Turkey.

Erment, Thomas: Department of Anesthesiology, Intensive Care and Pain Medicine, University Hospital Münster, Münster, Germany.

Farah, Fadi: St. Elizabeth's Medical Center, Boston, Massachusetts.

Fernandez-Bustamante, Ana*: University of Colorado School of Medicine, Department of Anesthesiology, Aurora, Colorado.

Ferreira, Cristina: Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

Fiore, Marco: Department of Women, Child and General and Specialized Surgery, University of Campania “L. Vanvitelli,” Naples, Italy.

Fonte, Ana: Department of Anesthesiology from Centro Hospitalar de Entre o Douro e Vouga, São Sebastião Hospital, Santa Maria da Feira, Portugal.

Fortià Palahí, Christina: Hospital Sagrat Cor, Barcelona, Spain.

Galimberti, Andrea: Dipartimento di Scienze della Salute, Università degli Studi di Milano, Milan, Italy; and SC Anestesia e Rianimazione, Asst Santi Paolo e Carlo, Centro Ricerca Coordinata Insufficienza Respiratoria, Milan, Italy.

Gama de Abreu, Marcelo*: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Garofano, Najja: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Giaccari, Luca Gregorio: Department of Women, Child and General and Specialized Surgery, University of Campania "L. Vanvitelli," Naples, Italy.

Gilsanz, Fernando: Department of Anesthesiology and Surgical Critical Care, Hospital Universitario La Paz, Madrid, Spain.

Girrbach, Felix: Department of anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany.

Gobbi, Luca: Department Morphology, Surgery and Experimental Medicine, Anesthesia and Intensive Care University Section, University of Ferrara, Ferrara, Italy.

Godfried, Marc Bernard*: Department of Anesthesiology, Onze Lieve Vrouwe Gasthuis, Amsterdam, The Netherlands.

Goettel, Nicolai*: Department of Anesthesia, Surgical Intensive Care, Prehospital Emergency Medicine and Pain Therapy, University Hospital Basel, University of Basel, Basel, Switzerland.

Goldstein, Peter A.*: Weill Cornell Medicine, New York, New York.

Goren, Or: Division of Anesthesia, Pain and Critical Care, Tel-Aviv Medical Center affiliated with Sackler Medical School, Tel Aviv University, Tel Aviv, Israel.

Gorlin, Andrew: Mayo Clinic Arizona, Phoenix, Arizona.

Granell Gil, Manuel*: Consorcio Hospital General Universitario of Valencia, Valencia, Spain.

Gratarola, Angelo: Anestesia e Rianimazione, IRCCS Policlinico San Martino, Genova, Italy.

Graterol, Juan*: Royal Cornwall Hospital National Health Service Trust, Truro, United Kingdom.

Guyon, Pierre: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Haire, Kevin: Magill Department of anesthesia, Chelsea and Westminster National Health Service Foundation Trust, London, United Kingdom.

Harou, Philippe: Polyclinique Montier la Celle, Saint-André-les-Vergers, France.

Helf, Antonia: Department of anesthesiology, University Hospital Würzburg, Würzburg, Germany.

Hemmes, Sabrina N.T.*: Amsterdam University Medical Center, University of Amsterdam, Department of Anesthesiology, Amsterdam, The Netherlands.

Hempel, Gunther: Department of Anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany.

Hernández Cádiz, María José: Consorcio Hospital General Universitario of Valencia, Spain

Heyse, Björn: Ghent University Hospital, Ghent, Belgium.

Hollmann, Markus W.*: Amsterdam University Medical Center, University of Amsterdam, Department of Anesthesiology, Amsterdam, The Netherlands.

Huercio, Ivan: Department of Anesthesiology and Surgical Critical Care, Hospital Universitario La Paz, Madrid, Spain.

Ilievska, Jasmina: University Clinic of Surgery "Ss. Naum Ohridski," Faculty of Medicine, University "Ss. Cyril and Methodius," Skopje, Macedonia.

Jakus, Lien: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Jeganath, Vijay: Royal Stoke University Hospital, University Hospitals of North Midlands National Health Service Trust, Stoke-on-Trent, Staffordshire, United Kingdom.

Jelting, Yvonne: Department of Anesthesiology, University Hospital Würzburg, Würzburg, Germany.

Jung, Minoa: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Kabon, Barbara*: Department of Anesthesia, Critical Care and Pain Medicine, Medical University Vienna, Vienna, Austria.

Kacha, Aalok*: Section of Critical Care Medicine, Department of Anesthesia and Critical Care, University of Chicago, Chicago, Illinois.

Karaman Ilić, Maja: Clinical Hospital Sveti Duh, J.J. Strossmayer, Faculty of Medicine, Zagreb, Croatia.

Karuppih, Arunthevaraja: St. Elizabeth's Medical Center, Boston, Massachusetts.

Kavas, Ayse Duygu: Marmara University Pendik Training and Research Hospital, Istanbul, Turkey.

Keli Barcelos, Gleicy: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Kellogg, Todd A.: Department of Surgery, Mayo Clinic, Rochester, Minnesota.

Kemper, Johann: Department of Anesthesiology, University Hospital Duesseldorf, Heinrich-Heine University Duesseldorf, Duesseldorf, Germany.

Kerbrat, Romain: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service

d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Khodr, Suraya: Cleveland Clinic Abu Dhabi, Al Maryah Island, Abu Dhabi, United Arab Emirates.

Kienbaum, Peter: Department of Anesthesiology, University Hospital Duesseldorf, Heinrich-Heine University Duesseldorf, Duesseldorf, Germany.

Kir, Bunyamin: Marmara University Pendik Training and Research Hospital, Turkey.

Kiss, Thomas: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Kivrak, Selin: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Klarić, Vlasta*: Clinic of Anesthesiology, Reanimation and Intensive Care Medicine, University Hospital Dubrava, Zagreb, Croatia.

Koch, Thea: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Köksal, Ceren: Fatih Sultan Mehmet Educational and Research Hospital, İstanbul, Turkey.

Kowark, Ana: Department of Anesthesiology, Medical Faculty, Rheinisch-Westfälische Technische Hochschule Aachen University, Aachen, Germany.

Kranke, Peter*: Department of Anesthesiology, University Hospital Würzburg, Würzburg, Germany.

Kuvaki, Bahar*: Dokuz Eylül University Faculty of Medicine, Department of Anesthesiology and Intensive Care, Balçova, Turkey.

Kuzmanovska, Biljana: Department of Anesthesiology, University Clinic for Traumatology, Orthopedics, Anesthesia, Reanimation, Intensive Care and Emergency Center, Skopje, Macedonia.

Laffey, John*: Department of Anesthesiology, St. Michael's Hospital, University of Toronto, Toronto, Ontario, Canada.

Lange, Mirko: Department of Anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany.

Lemos, Marília Freitas de: Department of Anesthesiology, Hospital do Câncer II, National Cancer Institute of Brazil, Rio de Janeiro, Brazil.

Licker, Marc-Joseph*: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie; Faculty of Medicine, Hôpitaux Universitaires de Genève, University of Geneva, Geneva, Switzerland.

López-Baamonde, Manuel: Department of Anesthesiology, Hospital Clínic, Barcelona, Spain.

López-Hernández, Antonio: Department of Anesthesiology, Hospital Clínic, Barcelona, Spain.

Lopez-Martinez, Mercedes: Department of Anesthesiology and Surgical Critical Care, Hospital Universitario La Paz, Madrid, Spain.

Luise, Stéphane: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

MacGregor, Mark*: Ashford and St. Peters National Health Service Foundation Trust, Ashford, United Kingdom.

Machado, Humberto S.*: Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal; Instituto Ciências Biomédicas Abel Salazar, Universidade do Porto, Porto, Portugal; and Centro de Investigação Clínica em Anestesiologia, Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

Magalhães, Danielle*: Associação Hospitalar Beneficente São Vicente de Paulo, Passo Fundo, Brazil.

Maillard, Julien: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Malerbi, Patrizia: Anestesia e Rianimazione, IRCCS Policlinico San Martino, Genova, Italy.

Manimekalai, Natesan: University of Mississippi Medical Center, Jackson, Mississippi.

Margaron, Michael*: St. Richard's Hospital, Chichester, United Kingdom.

Markstaller, Klaus*: Department of Anesthesia, Critical Care and Pain Medicine, Medical University Vienna, Vienna, Austria.

Martin, Archer K.: Department of Anesthesiology and Perioperative Medicine, College of Medicine, Mayo Clinic, Jacksonville, Florida.

Martin, David P.: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota.

Martin, Yvette N.: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota.

Martínez-Ocon, Julia: Department of Anesthesiology, Hospital Clínic, Barcelona, Spain.

Martin-Loeches, Ignacio*: Department of Intensive Care Medicine, Multidisciplinary Intensive Care Research Organization (MICRO), St. James's Hospital, Dublin, Ireland.

Maseda, Emilio: Department of Anesthesiology and Surgical Critical Care, Hospital Universitario La Paz, Madrid, Spain.

Matot, Idit*: Division of Anesthesia, Pain and Critical Care, Tel-Aviv Medical Center affiliated with Sackler Medical School, Tel Aviv University, Tel Aviv, Israel.

McAuliffe, Niamh: Department of Anesthesiology, St. Michael's Hospital, University of Toronto, Toronto, Ontario, Canada.

McKenzie, Travis J.: Department of Surgery, Mayo Clinic, Rochester, Minnesota.

Medina, Paulina: Department of Anesthesiology, School of Medicine, Pontificia Universidad Catolica de Chile, Santiago, Chile.

Meersch, Melanie: Department of Anesthesiology, Intensive Care and Pain Medicine, University Hospital Münster, Münster, Germany.

Menzen, Angelika*: St. Marienhospital gGmbH Friesoythe, Friesoythe, Germany.

Mertens, Els*: University Hospital Antwerp, Antwerp, Belgium.

Meurer, Bernd: Marienhospital Wesel, Wesel, Germany.

Meyer-Treschan, Tanja*: Department of Anesthesiology, University Hospital Duesseldorf, Heinrich-Heine University Duesseldorf, Duesseldorf, Germany.

Miao, Changhong*: Department of Anesthesiology, Fudan University Shanghai Cancer Center, Shanghai, China; and Department of Oncology, Shanghai Medical College, Fudan University, Shanghai, China.

Micalizzi, Camilla: Dipartimento di Scienze Chirurgiche e Diagnostiche Integrate, Università degli Studi di Genova, Genova, Italy; and Anestesia e Terapia Intensiva, IRCCS Policlinico San Martino, Genova, Italy.

Milić, Morena: Clinic of Anesthesiology, Reanimation and Intensive Care Medicine, University Hospital Dubrava, Zagreb, Croatia.

Módoło, Norma Sueli Pinheiro: Department of Anesthesiology, Universidade Estadual Paulista, Botucatu, Brazil.

Moine, Pierre: Department of Anesthesiology, University of Colorado School of Medicine, Aurora, Colorado.

Mölders, Patrick: Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Universitätsklinikum Knappschaftskrankenhaus Bochum, Bochum, Germany.

Montero-Feijoo, Ana: Department of Anesthesiology and Surgical Critical Care, Hospital Universitario La Paz, Madrid, Spain.

Moret, Enrique: University Hospital Germans Trias i Pujol, Institut Català de la Salut, Universidad Autónoma de Barcelona, Barcelona, Spain.

Muller, Markus K.: Frauenfeld Cantonal Hospital, Frauenfeld, Switzerland.

Murphy, Zoe: York Teaching Hospitals National Health Service Foundation Trust, York, United Kingdom.

Nalwaya, Pramod*: Royal Stoke University Hospital, University Hospitals of North Midlands National Health Service Trust, Stoke-on-Trent, Staffordshire, United Kingdom.

Naumovski, Filip*: Department of Anesthesiology, Reanimation and Intensive Care Medicine, University Clinic for Traumatology, Orthopedics, Anesthesia, Reanimation, Intensive Care and Emergency Center, Skopje, Macedonia.

Navalesi, Paolo*: Anesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Graeca University, Catanzaro, Italy.

Navarro e Lima, Lais Helena: Department of Anesthesiology, Universidade Estadual Paulista, Botucatu, Brazil.

Nesek Adam, Višnja*: Clinical Hospital Sveti Duh, Faculty of Medicine, J.J. Strossmayer University, Zagreb, Croatia.

Neumann, Claudia: Department of Anesthesiology and Intensive Care Medicine, University Hospital of Bonn, Bonn, Germany.

Newell, Christopher*: Southmead Hospital, North Bristol National Health Service Trust, Bristol, United Kingdom.

Nisnevitch, Zoufira*: Montefiore Medical Center, Bronx, New York.

Nizamuddin, Junaid: Section of Critical Care Medicine, Department of Anesthesia and Critical Care, University of Chicago, Chicago, Illinois.

Novazzi, Cecilia: Dipartimento di Anestesia, Azienda Ospedaliera Asst Settlaghi Ospedale di Circolo e Fondazione Macchi, Università dell'Insubria, Varese, Italy.

Nunes, Catarina S.: Universidade Aberta, Departamento de Ciências e Tecnologia, Porto, Portugal and Centro de Investigação Clínica em Anestesiologia, Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

O'Connor, Michael: Section of Critical Care Medicine, Department of Anesthesia and Critical Care, University of Chicago, Chicago, Illinois.

Oprea, Günther: Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Universitätsklinikum Knappschaftskrankenhaus Bochum, Bochum, Germany.

Orhan Sungur, Mukadder: Istanbul University, Istanbul Faculty of Medicine, Istanbul, Turkey.

Özbilgin, Şule: Dokuz Eylül University Faculty of Medicine, Department of Anesthesiology and Intensive Care, Balçova, Turkey.

Pace, Maria Caterina: Department of Women, Child and General and Specialized Surgery, University of Campania "L. Vanvitelli," Naples, Italy.

Pacheco, Marcos: Department of Anesthesiology from Centro Hospitalar de Entre o Douro e Vouga, São Sebastião Hospital, Santa Maria da Feira, Portugal.

Packianathaswamy, Balaji*: Hull and East Yorkshire Hospitals National Health Service Trust, Kingston upon Hull, United Kingdom.

Palma Gonzalez, Estefania: Área de Salud Don Benito-Villanueva, Don Benito, Spain.

Papaspyros, Fotios*: Polyclinique Montier la Celle, Saint-André-les-Vergers, France.

Paredes, Sebastián: Department of Anesthesiology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile.

Passavanti, Maria Beatrice: Department of Women, Child and General and Specialized Surgery, University of Campania "L. Vanvitelli," Naples, Italy.

Pedemonte, Juan Cristobal: Department of Anesthesiology, School of Medicine, Pontificia Universidad Catolica de Chile, Santiago, Chile.

Pelosi, Paolo*: Dipartimento di Scienze Chirurgiche e Diagnostiche Integrate, Università degli Studi di Genova, Genova, Italy; and Anestesia e Terapia Intensiva, IRCCS Policlinico San Martino, Genova, Italy.

Peremin, Sanja: Clinic of Anesthesiology, Reanimatology and Intensive Care Medicine, University Hospital Dubrava, Zagreb, Croatia.

Philipsenburger, Christoph: Department of Anesthesiology, Heidelberg University Hospital, Heidelberg, Germany.

Pinho, Daniela: Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

Pinho, Silvia: Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

Posthuma, Linda M.: Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Pota, Vincenzo: Department of Women, Child and General and Specialized Surgery, University of Campania “L. Vanvitelli,” Naples, Italy.

Preckel, Benedikt: Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Priani, Paolo: Department of Morphology, Surgery and Experimental Medicine, Anesthesia and Intensive Care University Section, University of Ferrara, Ferrara, Italy.

Putensen, Christian*: Department of Anesthesiology and Intensive Care Medicine, University Hospital of Bonn, Bonn, Germany.

Rached, Mohamed Aymen: Département d’Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d’Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Radoeshki, Aleksandar: University Clinic of Surgery “Ss. Naum Ohridski,” Faculty of Medicine, University “Ss. Cyril and Methodius,” Skopje, Macedonia.

Ragazzi, Riccardo: Department Morphology, Surgery and Experimental Medicine, Anesthesia and Intensive Care University Section, University of Ferrara, Ferrara, Italy.

Rajamanickam, Tamilselvan: Royal Stoke University Hospital, University Hospitals of North Midlands National Health Service Trust, Stoke-on-Trent, Staffordshire, United Kingdom.

Rajamohan, Arthi: Department of Anesthesiology, St. Michael’s Hospital, University of Toronto, Toronto, Ontario, Canada.

Ramakrishna, Harish*: Mayo Clinic Arizona, Phoenix, Arizona.

Rangarajan, Desikan*: Homerton University Hospitals National Health Service Foundation Trust, London, United Kingdom.

Reiterer, Christian: Department of Anesthesia, Critical Care and Pain Medicine, Medical University Vienna, Vienna, Austria.

Renew, J. Ross: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Jacksonville, Florida.

Reynaud, Thomas: Département d’Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d’Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Rhys, Rhidian: Southmead Hospital, North Bristol National Health Service Trust, Bristol, United Kingdom.

Rivas, Eva*: Department of Anesthesiology, Hospital Clínic, Instituto de Investigación Biomédica August Pi y Sunyer, Universitat de Barcelona, Barcelona, Spain.

Robitzky, Luisa: Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Universitätsklinikum Knappschafts-Krankenhaus Bochum, Bochum, Germany.

Rossaint, Rolf*: Department of Anesthesiology, Medical Faculty, Rheinisch-Westfälische Technische Hochschule Aachen University, Aachen, Germany.

Rubulotta, Francesca*: Imperial College Healthcare National Health Service Trust, London, United Kingdom.

Sabbatini, Giovanni: Dipartimento di Scienze della Salute, Università degli Studi di Milano, Milan, Italy; and Struttura Complessa Anestesia e Rianimazione, Azienda Socio Sanitaria Territoriale Santi Paolo e Carlo, Centro Ricerca Coordinata Insufficienza Respiratoria, Milan, Italy.

Samuels, Jon D.: Weill Cornell Medicine, Department of Anesthesiology, New York–Presbyterian Hospital, New York, New York.

Sanahuja, Josep Martí: Department of Anesthesiology, Hospital Clínic, Barcelona, Spain.

Sansone, Pasquale: Department of Women, Child and General and Specialized Surgery, University of Campania “L. Vanvitelli,” Naples, Italy.

Santos, Alice: Department of Anesthesiology, Centro Hospitalar São João, Porto, Portugal.

Sayedlahl, Mohamed: Department of Anesthesiology, King Abdullah Medical City, Makkah, Saudi Arabia; and Department of Anesthesiology, Mansoura University, Mansoura, Egypt.

Schaefer, Maximilian S.: Department of Anesthesiology, University Hospital Duesseldorf, Heinrich-Heine University Duesseldorf, Duesseldorf, Germany.

Scharffenberg, Martin: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Schiffer, Eduardo: Département d’Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d’Anesthésiologie; Faculty of Medicine, Hôpitaux Universitaires de Genève, University of Geneva, Geneva, Switzerland.

Schliewe, Nadja: Department of Anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany.

Schorer, Raoul: Département d’Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d’Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Schultz, Marcus J.: Department of Anesthesiology, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

Schumann, Roman: Department of Anesthesiology and Perioperative Medicine, Tufts Medical Center, Boston, Massachusetts.

Selmo, Gabriele: Università dell'Insubria, Dipartimento di Anestesia, Azienda Ospedaliera Asst Settlaghi Ospedale di Circolo e Fondazione Macchi, Varese, Italy.

Sendra, Mar: University Hospital Germans Trias i Pujol, Institut Català de la Salut, Universidad Autónoma de Barcelona, Barcelona, Spain.

Senturk, Mert*: Istanbul University, Istanbul Faculty of Medicine, Turkey.

Severgnini, Paolo*: Università dell'Insubria, Dipartimento di Anestesia, Azienda Ospedaliera Asst Settlaghi Ospedale di Circolo e Fondazione Macchi, Varese, Italy.

Shaw, Kate: York Teaching Hospitals National Health Service Foundation Trust, York, United Kingdom.

Shosholcheva, Mirjana*: University Clinic of Surgery "Ss. Naum Ohridski," Faculty of Medicine, University "Ss. Cyril and Methodius," Skopje, Macedonia.

Sibai, Abdulrazak*: Anesthesia Department, Ministry of National Guard Health Affairs, King Abdulaziz Medical City, Riyadh, Saudi Arabia.

Simon, Philipp: Department of Anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany.

Simonassi, Francesca: Dipartimento di Scienze Chirurgiche e Diagnostiche Integrate, Università degli Studi di Genova, Genova, Italy; and Anestesia e Terapia Intensiva, IRCCS Policlinico San Martino, Genova, Italy.

Sinno, Claudia: Università dell'Insubria, Dipartimento di Anestesia, Azienda Ospedaliera Asst Settlaghi Ospedale di Circolo e Fondazione Macchi, Varese, Italy.

Sivriköz, Nuket: Istanbul University, Istanbul Faculty of Medicine, Istanbul, Turkey.

Skandalou, Vasiliki*: Alexandra General Hospital, Athens, Greece.

Smith, Neil: Hull and East Yorkshire Hospitals National Health Service Trust, Kingston upon Hull, United Kingdom.

Soares, Maria: Serviço de Anestesiologia, Centro Hospitalar do Porto, Porto, Portugal.

Socorro Artiles, Tania: University Clinic Hospital Valencia, Valencia, Spain.

Sousa Castro, Diogo*: Department of Anesthesiology from Centro Hospitalar de Entre o Douro e Vouga, São Sebastião Hospital, Santa Maria da Feira, Portugal.

Sousa, Miguel: Department of Anesthesiology from Centro Hospitalar de Entre o Douro e Vouga, São Sebastião Hospital, Santa Maria da Feira, Portugal.

Spadaro, Savino*: Department Morphology, Surgery and Experimental Medicine, Anesthesia and Intensive Care University Section, University of Ferrara, Ferrara, Italy.

Sprung, Juraj*: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota.

Stamatakis, Emmanouil: Alexandra General Hospital, Athens, Greece.

Steiner, Luzius A.: Department of Anesthesia, Surgical Intensive Care, Prehospital Emergency Medicine and Pain Therapy, University Hospital Basel, University of Basel, Basel, Switzerland.

Stevenazzi, Andrea: University Hospital Germans Trias i Pujol, Institut Català de la Salut, Universidad Autónoma de Barcelona, Barcelona, Spain.

Suarez-de-la-Rica, Alejandro*: Department of Anesthesiology and Surgical Critical Care, Hospital Universitario La Paz, Madrid, Spain.

Suppan, Mélanie: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Teichmann, Robert: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Tena Guerrero, José María*: Área de Salud Don Benito-Villanueva, Don Benito, Spain.

Thiel, Bram: Department of Anesthesiology, Onze Lieve Vrouwe Gasthuis, Amsterdam, The Netherlands.

Tolós, Raquel: University Hospital Germans Trias i Pujol, Institut Català de la Salut, Universidad Autónoma de Barcelona, Barcelona, Spain.

Tore Altun, Gulbin: Marmara University Pendik Training and Research Hospital, Istanbul, Turkey.

Tucci, Michelle: University of Mississippi Medical Center, Jackson, Mississippi.

Turnbull, Zachary A.: Weill Cornell Medicine, New York, New York.

Turudić, Žana: Clinic of Anesthesiology, Reanimation and Intensive Care Medicine, University Hospital Dubrava, Zagreb, Croatia.

Unterberg, Matthias: Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Universitätsklinikum Knappschaftskrankenhaus Bochum, Bochum, Germany.

Van Limmen, Jurgen: Ghent University Hospital, Ghent, Belgium.

Van Nieuwenhove, Yves: Ghent University Hospital, Ghent, Belgium.

Van Waesberghe, Julia: Department of Anesthesiology, Medical Faculty, Rheinisch-Westfälische Technische Hochschule Aachen University, Aachen, Germany.

Vidal Melo, Marcos Francisco*: Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, Massachusetts.

Vitković, Bibiana: Clinic of Anesthesiology, Reanimation and Intensive Care Medicine, University Hospital Dubrava, Zagreb, Croatia.

Vivona, Luigi: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Vizcaychipi, Marcela*: Magill Department of Anesthesia, Chelsea and Westminster National Health Service Foundation Trust, London, United Kingdom.

Volta, Carlo Alberto: Department Morphology, Surgery and Experimental Medicine, Anesthesia and Intensive Care University Section, University of Ferrara, Ferrara, Italy.

Weber, Anne: Département d'Anesthésiologie, Pharmacologie, Soins Intensifs et Urgences, Service d'Anesthésiologie, Hôpitaux Universitaires de Genève, Geneva, Switzerland.

Weingarten, Toby N.: Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota.

Wittenstein, Jakob: Department of Anesthesiology and Intensive Care, Pulmonary Engineering Group, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany.

Wrigge, Hermann*: Department of anesthesiology and Intensive Care Medicine, University of Leipzig, Leipzig, Germany and Department of Anesthesiology, Intensive Care and Emergency Medicine, Pain Therapy; Bergmannstrost Hospital Halle, Halle, Germany.

Wyffels, Piet: Ghent University Hospital, Ghent, Belgium.

Yagüe, Julio: Department of Anesthesiology and Surgical Critical Care, Hospital Universitario La Paz, Madrid, Spain.

Yates, David: York Teaching Hospitals National Health Service Foundation Trust, York, United Kingdom.

Yavru, Ayşen: Istanbul University, Istanbul Faculty of Medicine, Istanbul, Turkey.

Zac, Lilach: Division of Anesthesia, Pain and Critical Care, Tel-Aviv Medical Center affiliated with Sackler Medical School, Tel Aviv University, Tel Aviv, Israel.

Zhong, Jing: Department of Anesthesiology, Fudan University Shanghai Cancer Center, Shanghai, China; and Department of Oncology, Shanghai Medical College, Fudan University, Shanghai, China.

Information and Funding. The Clinical Trials Network of the European Society of Anesthesiology provided financial support for the steering committee meetings, for onsite visits to participating sites, for the building of the electronic data capture system, and for advertising the study. The Technische Universität Dresden provided logistical support for the coordinating site. The Conselho Nacional de Desenvolvimento Científico e Tecnológico provided financial support for insurance in Brazil. The Association of Anaesthetists of Great Britain and Ireland and the Northern Ireland Society of Anaesthetists provided financial support for the participating sites in the United Kingdom.